

## Calcineurin Inhibits *VCXI*-Dependent $H^+/Ca^{2+}$ Exchange and Induces $Ca^{2+}$ ATPases in *Saccharomyces cerevisiae*

KYLE W. CUNNINGHAM<sup>1,2\*</sup> AND GERALD R. FINK<sup>2</sup>

Department of Biology, The Johns Hopkins University, Baltimore, Maryland 21218,<sup>1</sup> and Whitehead Institute, Massachusetts Institute of Technology, Cambridge, Massachusetts 02142

Received 21 September 1995/Returned for modification 27 November 1995/Accepted 27 December 1995

The *PMCI* gene in *Saccharomyces cerevisiae* encodes a vacuolar  $Ca^{2+}$  ATPase required for growth in high- $Ca^{2+}$  conditions. Previous work showed that  $Ca^{2+}$  tolerance can be restored to *pmc1* mutants by inactivation of calcineurin, a  $Ca^{2+}$ /calmodulin-dependent protein phosphatase sensitive to the immunosuppressive drug FK506. We now report that calcineurin decreases  $Ca^{2+}$  tolerance of *pmc1* mutants by inhibiting the function of *VCXI*, which encodes a vacuolar  $H^+/Ca^{2+}$  exchanger related to vertebrate  $Na^+/Ca^{2+}$  exchangers. The contribution of *VCXI* in  $Ca^{2+}$  tolerance is low in strains with a functional calcineurin and is high in strains which lack calcineurin activity. In contrast, the contribution of *PMCI* to  $Ca^{2+}$  tolerance is augmented by calcineurin activation. Consistent with these positive and negative roles of calcineurin, expression of a *vcx1::lacZ* reporter was slightly diminished and a *pmc1::lacZ* reporter was induced up to 500-fold by processes dependent on calcineurin, calmodulin, and  $Ca^{2+}$ . It is likely that calcineurin inhibits *VCXI* function mainly by posttranslational mechanisms. Activities of *VCXI* and *PMCI* help to control cytosolic free  $Ca^{2+}$  concentrations because their function can decrease *pmc1::lacZ* induction by calcineurin. Additional studies with reporter genes and mutants indicate that *PMRI* and *PMR2A*, encoding P-type ion pumps required for  $Mn^{2+}$  and  $Na^+$  tolerance, may also be induced physiologically in response to high- $Mn^{2+}$  and  $-Na^+$  conditions through calcineurin-dependent mechanisms. In these situations, inhibition of *VCXI* function may be important for the production of  $Ca^{2+}$  signals. We propose that elevated cytosolic free  $Ca^{2+}$  concentrations, calmodulin, and calcineurin regulate at least four ion transporters in *S. cerevisiae* in response to several environmental conditions.

The immunosuppressive drugs cyclosporin A and FK506 block activation of human T cells by specifically inactivating calcineurin, which is necessary for induction of interleukin-2 and other genes (for a review, see reference 59). Though these drugs are structurally unrelated, they both require binding to cytosolic receptor proteins, cyclophilin and FKBP-12, respectively, to form inactive complexes with calcineurin. Calcineurin may have additional functions in other cell types, and the loss of these functions may contribute to the side effects of these drugs, which include nephrotoxicity, neurotoxicity, and osteoporosis (13). Recent studies have implicated calcineurin as part of a regulatory cascade leading to control of the  $Na^+/H^+$  exchanger in T cells (56), the  $Na^+/K^+$  ATPase in kidney cells (5), and the *N*-methyl-D-aspartate receptor in neurons (42, 63). A better understanding of the biological roles for calcineurin in different cell types may promote the development of improved strategies for immunosuppression.

In the budding yeast *Saccharomyces cerevisiae*, homologs of calcineurin catalytic (15, 39, 68) and regulatory (16, 35) subunits have been cloned and characterized. The protein phosphatase activity of yeast calcineurin is stimulated by binding  $Ca^{2+}$ /calmodulin and inhibited by binding FK506 and cyclosporin A complexes with the yeast FKBP-12 and cyclophilin A homologs (18, 51), which are encoded by *FPR1* (11, 29) and *CPR1* (27, 64), respectively. Mutants lacking calcineurin function are partially defective in the recovery from prolonged exposure to mating pheromones (15, 16, 18). Calcineurin-deficient mutants also exhibit decreased tolerance to  $Na^+$ ,  $Li^+$ ,

and other ions (40, 45). Recently, inactivation of calcineurin has been shown to increase  $Ca^{2+}$  tolerance of at least one  $Ca^{2+}$ -sensitive mutant (14) and to decrease  $Ca^{2+}$  tolerance of others (21, 62). In none of these situations has the target(s) for calcineurin been identified.

The yeast genes *PMCI* and *PMRI* encode membrane proteins related to mammalian plasma membrane  $Ca^{2+}$  ATPases and sarcoendoplasmic reticulum  $Ca^{2+}$  pumps, respectively (4, 14, 57). Although the Pmc1p and Pmr1p proteins are localized to different organelles (Pmc1p to the vacuole and Pmr1p to the Golgi complex), they both function in  $Ca^{2+}$  sequestration and promote growth in media containing high  $Ca^{2+}$  concentrations. Mutants lacking *PMCI* function grow poorly in media containing 200 mM  $Ca^{2+}$ , and *pmc1 pmr1* double mutants are inviable even at low  $Ca^{2+}$  concentrations. Both of these effects are reversed by the action of FK506/FKBP-12 and cyclosporin A/cyclophilin A, by null mutations in the calcineurin A or B subunit, or by point mutations in calmodulin (14) which destroy its high-affinity  $Ca^{2+}$  binding sites (22). One interpretation of these results is that although calcineurin activation by  $Ca^{2+}$ /calmodulin is not essential for vegetative growth, it may inhibit the function of another factor that is necessary for growth under high- $Ca^{2+}$  conditions.

This report describes the cloning and characterization of a low-affinity vacuolar  $H^+/Ca^{2+}$  exchanger (17, 47, 50), whose function in vivo is inhibited by calcineurin activation. *VCXI* functions in  $Ca^{2+}$  tolerance and  $Ca^{2+}$  sequestration much more efficiently when calcineurin is inactivated that when it is activated. Studies of *vcx1* mutants suggest that calcineurin also promotes the expression of *PMCI*, *PMRI*, and *PMR2A*, which increases their contribution to tolerance of  $Ca^{2+}$ ,  $Mn^{2+}$ , and  $Na^+$ . The intracellular  $Ca^{2+}$  transporters encoded by *VCXI*, *PMCI*, and *PMRI* help to control cytosolic free  $Ca^{2+}$  con-

\* Corresponding author. Mailing address: Department of Biology, The Johns Hopkins University, 3400 N. Charles St., Baltimore, MD 21218. Phone: (410) 516-7844. Fax: (410) 516-5213. Electronic mail address: kwc@jhunix.hcf.jhu.edu.

centrations ( $[Ca^{2+}]_c$ ) in high- $Ca^{2+}$  conditions and thereby modulate calcineurin function. We suggest that this feedback control mechanism is important to promote efficient  $Ca^{2+}$  signaling in yeast cells exposed to a wide range of environmental conditions.

#### MATERIALS AND METHODS

**Yeast culture media.** Standard yeast culture media (60) were prepared from reagents supplied by Difco or BBL. YPD pH 5.5 medium was prepared and supplemented with  $CaCl_2$  (Sigma catalog no. C 2536) and either Difco Noble Agar or BBL granulated agar as described previously (14). Samples of FK506 were generously provided by Vertex Pharmaceuticals, Inc. (Cambridge, Mass.) and Fujisawa Corp. (Tokyo, Japan) and added where indicated from 0.2-mg/ml stock solutions in dimethyl sulfoxide.

**Cloning of *VCX1* and recombinant DNA.** All procedures involving recombinant DNA in *S. cerevisiae* and *Escherichia coli* were performed by using standard techniques (58) as follows. The *pmc1::LEU2* strain K473 was transformed with a high-copy-number library of yeast genomic DNA carried on the vector pRS202 (13a) and plated on synthetic complete (SC)-Ura agar medium at a density of approximately 5,000 colonies per plate. After 2 days of incubation at 30°C, the colonies were replica plated to YPD pH 5.5 medium supplemented with 200 mM  $CaCl_2$  and incubated at 30°C for 3 days.  $Ca^{2+}$ -tolerant clones were recovered, and plasmid DNA was isolated and characterized by restriction mapping. Of 24 plasmids isolated independently by this approach, 1 plasmid contained *PMCI*, 1 plasmid contained *PMRI*, 2 plasmids contained overlapping inserts spanning *VCX1*, and 20 plasmids contained overlapping inserts spanning *CRP1*, a previously undescribed gene locus currently under evaluation. Deletion mapping revealed that the functional *VCX1* gene was carried on a 1.7-kb minimal fragment which hybridized to lambda-prime clone 2439 representing a new locus on chromosome IV (55). The 1.7-kb fragment in pRS202 (pKC159) was sequenced on both strands with full overlap, using a nested deletion strategy (Amersham). Sequencing and restriction mapping of adjacent sequences indicate that *VCX1* is adjacent to *CDC48* and *ORF-D* (19). To construct pKC72 used for creating *vcx1* null mutants, the 1.7-kb fragment containing *VCX1* was first subcloned into plasmid pBSII (Stratagene), and then the 559-bp segment between the *StuI* and *MseI* sites in *VCX1* was replaced with a 5.0-kbp fragment of pSE1076 containing *URA3* and kanamycin resistance flanked by 1.1-kbp direct repeats of *hisG* (1). The *VCX1::hemagglutinin* (HA) epitope-tagged variant (*VCX1::HA*) on pKC142 was constructed by a multistep process. A 7.5-kbp genomic DNA fragment spanning *VCX1* was first inserted into low-copy-number centromere-based plasmid pRS316 (61), forming pKC98. Site-directed mutagenesis of pKC98 was performed to create a unique *NheI* cleavage site at codon +3 (forming pKC140), into which was inserted a 96-bp *XbaI* fragment of pKC54 (14) coding for three repeats of the HA epitope (forming pKC142). Sequencing and complementation assays of pKC140 and pKC142 confirmed that the expected modifications were successful and functional in yeast strains. To rescue the *VCX1-D1* allele, a 0.78-kbp *SacI*-to-*Clal* fragment of pKC159 containing the 5' end of *VCX1* was inserted into pRS303 (forming pKC106), which was digested with *SauI* and integrated into the *VCX1* locus by transformation of strains K482 and K482-1 (14). Genomic DNA prepared from these transformed strains was digested with *SaI* and treated with T4 DNA ligase, and plasmids pKC129 (*VCX1*) and pKC130 (*VCX1-D1*) were recovered by transformation into *E. coli* DH5 $\alpha$  and then sequenced by using complementary oligonucleotides. Reintegration of pKC130 into *pmc1* mutants produced the original  $Ca^{2+}$ -tolerant phenotype associated with *VCX1-D1*. Plasmid pKC151 was isolated from a cDNA library (38) and contained a functional *PMCI* cDNA under control of the *GALI* promoter in pRS316-*gal*.

A series of *lacZ* reporter genes was constructed from the well-characterized *cyc1::lacZ* reporters pLG $\Delta$ 312 and pLG $\Delta$ 178 (24). The *vcx1::lacZ* reporter plasmid pKC200 was constructed by first removing the *cyc1* sequences from pLG $\Delta$ 178 with *XhoI* and *BamHI* and then inserting a 2.0-kbp *HindIII*-to-*NheI* segment of pKC142 along with flanking polylinker sequences from pBSII and YEp356R (44). The *pmr1::lacZ* reporter plasmid pKC199 was constructed by inserting a 1,180-bp *XhoI*-*BamHI* fragment of pKC11 (37) into *XhoI*-plus-*BamHI*-digested pLG $\Delta$ 178. The *pmr2A::lacZ* reporter plasmid pKC201 contained a 1.4-kbp *Sall*-*EcoRI* fragment from plasmid B1999 (57) inserted into *XhoI*-plus-*BamHI*-digested pLG $\Delta$ 178 along with flanking polylinker sequences from pBSII. The reporter plasmid pKC190 containing *pmc1::lacZ* was constructed by inserting a 585-bp *HindIII*-*XbaI* fragment of pKC45 (14) plus flanking polylinker sequences from YEp356R into *XhoI*-plus-*BamHI*-digested pLG $\Delta$ 178. pKC211 is a derivative of pKC190 in which 413 bp of *PMCI* untranslated DNA (from *HindIII* at -579 to *AflII* at -166 relative to the initiation codon) is deleted, leaving 172 bp of *PMCI* 5' untranslated DNA containing putative transcriptional and translational initiation sequences. pKC191 is a derivative of pKC190 and pLG $\Delta$ 178 in which *cyc1* transcriptional and translational initiation sequences (from *XhoI* to *BamHI*) replace the 172-bp *AflII*-*BamHI* fragment of pKC190, leaving 413 bp of *PMCI* 5' untranslated DNA (from *HindIII* to *AflII*). pLB178-43 contains three copies of oligonucleotide TCCACGAAAA, which confers cell cycle regulation on *HO*, inserted into the *XhoI* site of pLG $\Delta$ 178 (10).

**Construction of yeast strains.** All yeast strains listed in Table 1 are derivatives

TABLE 1. Yeast strains used in this study

Strain	Genotype <sup>a</sup>	Reference
W303-1A	+	65
JGY41	<i>cmd1-3</i>	22
K470	<i>PMCI::HA (pmc1::URA3)</i>	14
K473	<i>pmc1::LEU2</i>	14
K603	<i>cnb1::LEU2</i>	14
K605	<i>pmc1::TRP1</i>	14
K607	<i>cnb1::LEU2 pmc1::TRP1</i>	14
K609	<i>pmr1::HIS3</i>	14
K617	<i>vcx1::URA3</i>	
K619	<i>cnb1::LEU2 vcx1::URA3</i>	
K625	<i>pmr1::HIS3 vcx1::URA3</i>	
K633	<i>pmr2::HIS3</i>	
K635	<i>cnb1::LEU2 pmr2::HIS3</i>	
K641	<i>pmr2::HIS3 vcx1::URA3</i>	
K643	<i>cnb1::LEU2 pmr2::HIS3 vcx1::URA3</i>	
K661	<i>vcx1<math>\Delta</math></i>	
K663	<i>cnb1::LEU2 vcx1<math>\Delta</math></i>	
K665	<i>pmc1::TRP1 vcx1<math>\Delta</math></i>	
K667	<i>cnb1::LEU2 pmc1::TRP1 vcx1<math>\Delta</math></i>	
K669	<i>cmd1-3 vcx1<math>\Delta</math></i>	
K673	<i>cmd1-3 pmc1::TRP1 vcx1<math>\Delta</math></i>	
K685	<i>cmd1-3</i>	
K689	<i>cmd1-3 pmc1::TRP1</i>	
K697	<i>VCX1::HA cnb1::LEU2</i>	
K698	<i>VCX1::HA</i>	
K699	<i>PMCI::HA</i>	
K737	<i>pmc1::TRP1 VCX1-D1</i>	

<sup>a</sup> All strains contain additional mutations (*MATA ade2-1 can1-100 his3-11,15 leu2-3,112 trp1-1 ura3-1*) and are isogenic to W303-1A.

of strain W303-1A (65) which were constructed through transformation or isogenic crosses by using standard techniques (26). The *vcx1::URA3* null allele was originally created by transformation of diploid strain KXC178 (*MATA $\alpha$  cnb1::LEU2/+ pmc1::TRP1/+ pmr2::HIS3/+*) with the 5.7-kbp *KpnI*-to-*SacI* fragment of pKC72. Uracil prototrophs were screened by Southern blotting to confirm replacement of one *VCX1* allele with *vcx1::URA3* and then sporulated and subjected to tetrad analysis. The *vcx1 $\Delta$*  null allele was recovered by selection against *vcx1::URA3* on 5-fluoro-orotic acid medium (9). The *pmr2::HIS3* null mutation was first introduced into strain K473 by using plasmid B2000 (57) and confirmed by Southern blotting. A  $Ca^{2+}$ -tolerant and 5-fluoro-orotic acid-resistant transformant of K619 with a 7.5-kbp *KpnI*-to-*SacI* fragment of pKC142 was isolated to obtain strain K697 in which *vcx1::URA3* is replaced with *VCX1::HA*. Strain K699 was constructed by selecting for uracil auxotrophs of K470 and screening for retention of the triple HA epitopes by Western blotting (immunoblotting). All other strains were derived from crosses with other strains in the W303-1A background, and in each case no independently segregating suppressors or enhancers were evident.

**Immunological techniques.** Immunofluorescence microscopy was performed on log-phase yeast cells as described previously (14) except that ascites fluid containing anti-HA monoclonal antibody 12CA5 (Boehringer Mannheim) was used at a dilution of 1:400 and cyanine-3-conjugated goat anti-mouse immunoglobulin G (Rockland, Inc.) was used at a dilution of 1:250. For Western blot analysis (58), log-phase cultures were harvested, lysed with NaOH plus  $\beta$ -mercaptoethanol, and then precipitated with trichloroacetic acid (53). Precipitates were solubilized at 65°C for 10 min in 0.2 ml of sodium dodecyl sulfate (SDS) sample buffer, separated on polyacrylamide gels, transferred to Hybond Super-C (Amersham), stained with Ponceau S (Sigma), and imaged using with an enhanced chemiluminescence detection kit (Amersham) after probing with monoclonal antibody 12CA5 at a dilution of 1:4,000.

**<sup>45</sup>Ca<sup>2+</sup> transport in cell lysates.** Uptake of <sup>45</sup>Ca<sup>2+</sup> into particles of crude spheroplast lysates was quantitated by using a previously developed method (17), with slight modifications. Briefly, yeast strains were grown to mid-log phase in YPD medium or SCGal-Ura medium and then harvested, washed once in buffer TS (10 mM Tris-Cl [pH 8.0], 1 M sorbitol), and digested with 0.1 mg of Zymolyase 100T (Seikagaku) per ml for 45 min at 37°C in buffer TS plus 0.1%  $\beta$ -mercaptoethanol. The resulting spheroplasts were washed twice with 1 ml of buffer TS, resuspended in 1 ml of YPD medium supplemented with 0.6 M sorbitol and 50 mM potassium phosphate (pH 7.5), and incubated for 1 h at 37°C. The partially regenerated spheroplasts were cooled on ice, washed once with ice-cold buffer A [10 mM sodium piperazine-*N,N'*-bis(ethanesulfonic acid) (Na-PIPES), 10 mM sodium *N*-2-hydroxyethylpiperazine-*N'*-2-ethanesulfonic acid (Na-HEPES), 2 mM MgCl<sub>2</sub>, 100 mM KCl, 0.6 M sorbitol (pH 7.0)], and cell

pellets were resuspended with 20  $\mu$ l of buffer A on ice. Three minutes before the reactions were initiated, each cell suspension was lysed osmotically by diluting 10  $\mu$ l into 1.0 ml of prewarmed buffer B (10 mM Na-PIPES, 10 mM Na-HEPES, 2 mM MgCl<sub>2</sub>, 100 mM KCl [30°C, pH 7.0]) supplemented with 10 or 100  $\mu$ M CaCl<sub>2</sub>, 1 to 2  $\mu$ Ci of [<sup>45</sup>Ca]Cl<sub>2</sub> (New England Nuclear) per ml, and where indicated 10  $\mu$ M concanamycin B (generous gift of H. L. Ploegh, Massachusetts Institute of Technology). Reactions were initiated at time zero by the addition of ATP to 1 mM from a 100 mM pH 7.0 stock solution. At the indicated time intervals, 100- or 200- $\mu$ l aliquots were removed, diluted into 4 ml of ice-cold 20 mM MgCl<sub>2</sub>, rapidly filtered onto 2.4-cm GFF filters (Whatman), and washed three times with the same solution. Filters were dried, and radioactivity was quantitated by liquid scintillation counting. Ca<sup>2+</sup> uptake per 10<sup>9</sup> lysed cells was calculated from the measured counts, the specific activity, and culture turbidity.

**Measurement of nonexchangeable Ca<sup>2+</sup> pools in living cells.** Yeast strains were grown to mid-log phase (optical density at 600 nm [OD<sub>600</sub>] = 0.7 to 1.8) at 30°C in YPD medium, harvested, and then resuspended to a final OD<sub>600</sub> of 0.1 in 1.5 ml of fresh YPD pH 5.5 medium supplemented with 5 mM CaCl<sub>2</sub> and ~20  $\mu$ Ci of [<sup>45</sup>Ca]Cl<sub>2</sub> per ml. After growth at 30°C for an additional 7.25 h (four to five doublings in cell density), cultures were harvested, washed twice with medium lacking <sup>45</sup>Ca<sup>2+</sup>, resuspended in an equal volume, and incubated for an additional 40 min at 30°C to allow efflux of the exchangeable Ca<sup>2+</sup> pool. Total cell-associated radioactivity was determined by rapid filtration and liquid scintillation counting (14), and total cell number was determined by turbidity measurements. The average of duplicate samples was used to calculate the amount of Ca<sup>2+</sup> remaining in the nonexchangeable pool. The levels of Ca<sup>2+</sup> in the exchangeable pool was relatively constant in these strains at 2.6  $\pm$  0.7 (standard deviation) nmol/10<sup>9</sup> cells.

**Ion tolerance assays.** Yeast strains were inoculated into YPD pH 5.5 medium and grown overnight at 30°C. Saturated cell suspensions were then diluted 500-fold into 8 or 12 0.2-ml cultures in 96-well flat-bottom dishes containing YPD pH 5.5 medium supplemented with a wide range of CaCl<sub>2</sub>, MnCl<sub>2</sub>, or NaCl. Suspensions were mixed and incubated 16 to 20 h at 30°C without shaking, and the OD<sub>570</sub> was measured for each resuspended culture, using a microplate spectrophotometer (Dynatech Laboratories). All added salts were completely soluble under these conditions. The amount of each cation causing a 50% decrease in cell growth relative to the growth of unsupplemented cultures was interpolated from linear plots of the data.

**Miscellaneous procedures.** Total cellular  $\beta$ -galactosidase was determined at room temperature in cells permeabilized with chloroform and SDS (23), using the substrate *o*-nitrophenyl- $\beta$ -D-galactopyranoside (Sigma Chemical Co.).

**Nucleotide sequence accession number.** The DNA sequences of *VCXI* and *VCXI-D1* are listed in the GenBank and EMBL databases under accession number U36603.

## RESULTS

**Isolation and analysis of *VCXI*.** Calcineurin appears to inhibit a factor required for growth of *pmc1* mutants in media containing high Ca<sup>2+</sup> concentrations (14). To identify this target, we isolated genes from a high-dosage library which, when overexpressed, restored growth to *pmc1* mutants in YPD pH 5.5 medium containing 200 mM CaCl<sub>2</sub> (see Materials and Methods). Analysis of 24 independent clones yielded plasmids containing four different genes: *PMCI*, *PMRI*, and two novel genes termed *CRP1* and *VCXI*. The predicted 411-amino-acid product of *VCXI* (Fig. 1) indicated that it might be an appropriate candidate for inhibition by calcineurin. The Vcx1p protein contains 11 putative transmembrane domains, as determined from hydropathy plots (Fig. 2B), and shows the highest sequence similarity to the retinal rod Na<sup>+</sup>/Ca<sup>2+</sup>,K<sup>+</sup> exchanger (54) and to the cardiac Na<sup>+</sup>/Ca<sup>2+</sup> exchanger (46) (Fig. 2C). Vcx1p and these mammalian exchangers are predicted to be similar in transmembrane organization, although the mammalian enzymes also contain cleaved leader peptides and two large hydrophilic insertions. The regions with the greatest homology between these gene products encompass putative membrane spanning domains in the N- and C-terminal halves of the proteins. In a BLAST search of GenBank release 90 (2), open reading frames with unknown functions in the genomes of *E. coli* (accession number U18997) and *S. cerevisiae* (accession number P42839) were found to be significantly related to that of *VCXI*. The *VCXI* gene product exhibits little or no sequence similarity to a putative H<sup>+</sup>/Ca<sup>2+</sup> exchanger of *E. coli* (32). Thus, *VCXI* overexpression may prevent Ca<sup>2+</sup>-depen-

```

GATCAAATTCGCCGAAAAATAAAAAATTCCTCACGAGAATATGGAATTACTACTCGCCC 60
AGCATTATTTATCTCTCAGAGTAAAAAGAGTTTCCTCTTCTCAAATTAAGAACG 120
AGTCCATAAATATAGTTGACCTCCAGCGATATCAATTCATCGGCTGTAGCAAAATAA 180
AACACATAGATACAATGGATGCACTACCCCACTATTAACIGTGGCAACAGCATCCP 240
1 M D A T T P L L T V A N S H P
GCCCGCAATCCAAAGCACACTGCATGGAGAGCAGCTGTGTATGATTACAGATATTTTG 300
16 A R N P K H T A W R A A V Y D L Q Y I L
AAAGCGTCACCCCTGAATTTCCATTGGTATTGTCCTTTAGGCTGATTTGGGGACAC 360
36 K A S P L N F L L V F V P L G L I W G H
TTCCAACATATCATACACTGACATTTCTTTTAAATTTCTTGGCAATTAACCGTTGGCA 420
56 F Q L S H T L T F L F N F L A I I P L A
GCTATCTGGCTAATGCCAGGAAGAGTGGCTGATGAAGCTGGTAACACCATTGGGGGA 480
76 A I L A N A T E E L A D K A G N T I G G
CTGTAATAGCTACTTTTGGTAACCGTGGAACTAATGTTTCTATCATTTGCCCTGAAA 540
96 L L N A T F G N A V E L I V S I I A L K
AAAGTCAAGTGAGAATTTGTCAGGCTCGATGCTAGGTAGTCTTCTTCAATTTGCTG 600
116 K G Q V R I V Q A S M L G S L L S N L L
TTAGTCTTGGATTATGCTTCATATTCGGTGGATACAATAGAGTCCCAACAGACATCAAC 660
136 L V L G L C F I F G G Y N R V Q Q T F N
CAAACCGCGCTCAAAAATGCTCTTACTTGCCTTACTTGCCTGCGTGCATCCCTACTGATT 720
156 Q T A A Q T M S S L L A I A C A S L L
CCCCGTGCTTTAGAGGCCACCTACCTCATGGCAAGGAAGACCATTATCGATGGAAA 780
176 P A A F R A T L P H G K E D H F I D G K
ATATTGGAGTTATCCAGAGGCCCTCTATTGTTATTCCTACGTTTACGTTTGTCTCTTA 840
196 I L E L S R G T S I V I L I V Y V L F L
TATTTCCAGCTAGGAGCCATCACCGCTTGTGGCAACAGGAAGAGACCGATGAA 900
216 Y F Q L G S H H A L F E Q Q E E T D E
GTTATGAGCACCATTTCCAGGAATCCACATCACTCTTGGATGTCAGTCAATCATTTGGTG 960
236 V M S T I S R N P H S L S V K S S L V
ATACTTCTAGTACAACGTGATCATCTCTTTTGGCGGATTTCTAGTCCGATACGATA 1020
256 I L L G T T V I I S F C A D F L V G T I
GACAACGTTGTTGAATCACCAGGCTATCAAAAATTTATAGGTTGATTTGCATCTCT 1080
276 D N V V E S T G L S K T F I G L I V I P
ATTGTGGTAAATGCCGAGACATGCTACTTCAGTCTTGGTGGCCATGAAGGATAAAGATG 1140
296 I V G N A A E H V T S V L V A M K D K M
GATCTGGCGCTAGGTTGCTTCCATCGGTTCTCTTTACCAAGTTGCTTATTTGTTACCA 1200
316 D L A L G V A I G S S L Q V A L F V T P
TTCATGGTTCTTGTGGGCTGGATGATCGATGTTCCAAATGACGCTAAATTTCTCCACTTT 1260
336 F M V L V G W M I D V P M T L N F S T F
GAAACCGCTACTCTTTTATTTGCTGTTTCTTCAATTAATTAATCTCGATGGTGAG 1320
356 E T A T L F I A V F L S N Y L I L D G E
TCAAATGTTGGAGGGTGTATGCTCTAGCTATGATATGATTTGATGCAATGGCATTT 1380
376 S N W L E G V M S L A M Y I L I A M A F
TTCTATTCCAGACGAAAAACCCCTGACTCTATGGAAATAGTTTATGATGAAGAACA 1440
396 F Y Y P D E K T L D S I G N S L
GATGCGCAAAATTCACCATTTCTTTTTTTTATATTTTATCTCTTCGTAATTTCTAT 1500
GATGGACCAATACAAGATGATATGACTAGCAAGCTATTTATGATCGTTATATTATA 1560
CATATTATGTTACGCACTAATTTTTATATCGAATTTATCTGAAAAAAGTTTCAA 1620
ACCAACAAAATTTTGTACACTTTCGAAACTCAAATTTGACACCATTTCCATCTGAT 1680
AAAAGGTTAATATTTTCGAAGCTT 1704

```

FIG. 1. DNA sequence of the *VCXI* locus and predicted gene product. The DNA sequence (residues numbered at the right) and deduced protein sequence (residues numbered at the left) of the *VCXI* locus were determined as described in Materials and Methods and deposited in GenBank (accession number U36603). Sequencing of adjacent DNA showed that residues 1699 through 1704 are contiguous with residues 6485 through 6480 of a previously sequenced locus encoding *CDC48* (19). Sequencing of *VCXI-D1*, a spontaneous mutation which confers a dominant Ca<sup>2+</sup> tolerance phenotype in *pmc1* mutants, revealed only a single nucleotide substitution (underlined) in which the ATG (Met) codon at position 383 is converted to ATA (Ile).

dent growth inhibition of *pmc1* mutants by increasing Ca<sup>2+</sup> export from the cytosol.

***VCXI* encodes a vacuolar H<sup>+</sup>/Ca<sup>2+</sup> exchanger.** An oligonucleotide encoding three tandem repeats of the HA epitope was fused in frame near the predicted translational start site of *VCXI* (see Materials and Methods) in order to visualize the gene product in situ. The resulting *VCXI::HA* derivative was fully functional in assays for high-dosage suppression of *pmc1* and low-copy-number complementation of *vcx1* mutant phenotypes (see Fig. 6). Cells carrying *VCXI::HA* displayed epitopes exclusively near the vacuole membrane (Fig. 3A), whereas a control *VCXI* strain lacking the epitope tag displayed only weak background staining (Fig. 3B). The distribution of Vcx1p by immunofluorescence was similar to but much brighter than that of Pmc1p (14).

The major Ca<sup>2+</sup> uptake activity observed in crude cell ex-




FIG. 2. Predicted features of Vcx1p. (A) Predicted open reading frame and partial restriction map of the functional *VCX1* locus. (B) Kyte-Doolittle hydrophathy plot (window = 15 residues) of the predicted Vcx1p sequence. Putative transmembrane domains are numbered 1 to 11. (C) Multiple sequence alignment of Vcx1p, bovine retinal  $\text{Na}^+/\text{Ca}^{2+}, \text{K}^+$  exchanger (54), and canine  $\text{Na}^+/\text{Ca}^{2+}$  exchanger (46). Highlighted residues indicate identity in at least two of three sequences, and overlines correspond to predicted transmembrane domains. Numbers at the right refer to amino acid positions. No significant similarities were detected outside the aligned regions.

tracts and in purified vacuole membrane vesicles is a low-affinity  $\text{H}^+/\text{Ca}^{2+}$  exchange which utilizes a pH gradient formed by the vacuolar  $\text{H}^+$  ATPase (17, 47, 50). To test whether *VCX1* function is necessary for this activity, we assayed for  $^{45}\text{Ca}^{2+}$  uptake in total cell lysates (17) prepared from a wild-type strain (W303-1A) and a *vcx1* null mutant (K661) in which the chromosomal *VCX1* open reading frame was partially deleted and disrupted with foreign sequences (see Materials and Methods). Prior to the addition of ATP, the levels of  $^{45}\text{Ca}^{2+}$  accumulation in the particulate fraction were similar in the two lysates (Fig. 4A, time zero). However, after ATP was added, the *VCX1* lysate accumulated much more  $^{45}\text{Ca}^{2+}$  than the *vcx1* lysate, and the former activity was inhibited by addition of 10  $\mu\text{M}$  concanamycin B, a compound related to bafilomycin A that specifically inhibits vacuolar  $\text{H}^+$  ATPases (33). Both *VCX1* and *vcx1* strains accumulate the weak base quinacrine in their vacuoles in vivo, as detected by fluorescence microscopy (66), and this accumulation was blocked by 10  $\mu\text{M}$  concanamycin B (data not shown). These results show that vacuolar  $\text{H}^+/\text{Ca}^{2+}$  exchange activity is markedly reduced in *vcx1* mutants, but vacuole morphology and vacuolar acidification are not noticeably affected by the mutation.

The residual concanamycin B-resistant  $\text{Ca}^{2+}$  transport detected in *vcx1* mutants was not detected in *pmc1 vcx1* mutants (data not shown), suggesting that *PMC1* may encode a functional  $\text{Ca}^{2+}$  pump. To test this point directly, a *vcx1 pmc1* double mutant (K665) was transformed with an empty plasmid or pKC151, which expresses *PMC1* in galactose medium

through the strongly inducible *GAL1* promoter. After growth in galactose medium, cell lysates were prepared and assayed for  $^{45}\text{Ca}^{2+}$  uptake as described above except that  $\text{CaCl}_2$  in the buffer was reduced from 100 to 10  $\mu\text{M}$  and the ATP-independent transport activity in parallel samples was subtracted for each time point (Fig. 4B). Lysates of the strain expressing *PMC1* contained significant levels of ATP-dependent  $\text{Ca}^{2+}$  transport activity (Fig. 4B) that was resistant to concanamycin B (data not shown), whereas lysates of the strain lacking *PMC1* contained very little of this activity (Fig. 4B). These data show that *VCX1* encodes a vacuolar  $\text{H}^+/\text{Ca}^{2+}$  exchanger and *PMC1* encodes a vacuolar  $\text{Ca}^{2+}$  ATPase. Additionally, the cell-free  $\text{Ca}^{2+}$  transport assay shows that *VCX1*-dependent activity is much greater than residual *PMC1*-dependent activity (Fig. 4A) under these optimized assay conditions (containing 100  $\mu\text{M}$   $\text{Ca}^{2+}$ ). However, the following results indicate that in growing cells, *PMC1* functions are greater than *VCX1* functions as a result of both lower  $[\text{Ca}^{2+}]_c$  and regulation by calcineurin.

***VCX1*-dependent  $\text{Ca}^{2+}$  sequestration is enhanced when calcineurin is inactive.** To examine the role of *VCX1* in vivo,  $\text{Ca}^{2+}$  sequestration into a nonexchangeable intracellular pool was quantitated in log-phase cells containing various combinations of the *vcx1*, *pmc1*, and *cnb1* null mutations (Fig. 5). Each strain was labeled uniformly with  $^{45}\text{Ca}^{2+}$  in duplicate cultures during four to five generations of logarithmic growth in YPD pH 5.5 medium supplemented with 5 mM  $\text{CaCl}_2$ . The cells were collected, washed, and subjected to a 30-min chase period in medium lacking radioactivity to allow efflux of the exchangeable  $\text{Ca}^{2+}$  pool. The cells were then rapidly collected on filters, washed, and counted to estimate the nonexchangeable  $\text{Ca}^{2+}$  pool. The wild-type cells accumulated approximately four times more  $\text{Ca}^{2+}$  in the nonexchangeable pool than *pmc1* mutants (Fig. 5, bars 1 and 3) as observed previously (14), and these levels were only slightly affected by the additional *vcx1* null mutations (bars 2 and 4). Thus, relative to *PMC1*, *VCX1* plays only a small role in vacuolar  $\text{Ca}^{2+}$  sequestration in proliferating cells. However, the large increase in  $\text{Ca}^{2+}$  sequestration observed in *pmc1 cnb1* double mutants (bar 5) relative to *pmc1* mutants is completely dependent on *VCX1* function because *vcx1 pmc1 cnb1* triple mutants accumulate only low levels of nonexchangeable  $\text{Ca}^{2+}$  (bar 6). Qualitatively similar results were obtained in experiments in which extracellular  $\text{Ca}^{2+}$  was decreased to about 0.2 mM (YPD medium) or increased to 20 mM (data not shown), although the sizes of exchangeable and nonexchangeable pools varied considerably under these conditions as observed previously (17). Therefore, *VCX1* plays a much larger role in  $\text{Ca}^{2+}$  sequestration in *cnb1* mutants than in strains in which calcineurin is functional.

***VCX1* functions in  $\text{Ca}^{2+}$  tolerance when calcineurin is inactive.** The role of *VCX1* in  $\text{Ca}^{2+}$  tolerance was addressed by monitoring growth of *vcx1* mutants in liquid media supplemented with increasing concentrations of  $\text{CaCl}_2$ . After a 16-h incubation, the levels of added  $\text{CaCl}_2$  that caused a 50% decrease in relative growth (the  $\text{IC}_{50}$ s) were 360 and 320 nM for wild-type and *vcx1* strains, whereas the  $\text{IC}_{50}$ s for *pmc1* mutants and *pmc1 vcx1* double mutants were approximately 65 and 38 nM, respectively (Fig. 6). In five similar experiments, *VCX1* was always found to confer a relatively low level of  $\text{Ca}^{2+}$  tolerance, although the measured  $\text{IC}_{50}$ s varied somewhat because of variations in inoculum size and time of incubation.

In contrast to the small role of *VCX1* in strains in which calcineurin is functional, *VCX1* confers a much larger degree of  $\text{Ca}^{2+}$  tolerance when calcineurin has been inactivated. The high  $\text{Ca}^{2+}$  tolerance of *cnb1 pmc1* double mutants ( $\text{IC}_{50}$ , ~320 nM  $\text{CaCl}_2$ ) is completely dependent on *VCX1* function because *cnb1 pmc1 vcx1* triple mutants are extremely sensitive to




FIG. 3. Immunofluorescence localization of epitope-tagged Vcx1p. A centromere plasmid containing a functional epitope-tagged *VCX1::HA* derivative was constructed as described in Materials and Methods. Strain W303-1A harboring *VCX1::HA* (A) or *VCX1* (B) was grown to early exponential phase in SC-Ura medium, harvested, fixed with formaldehyde, and processed for immunofluorescence microscopy using the anti-HA monoclonal antibody 12CA5 (Boehringer) and cyanine-9-conjugated goat anti-mouse immunoglobulin G antibody as described in Materials and Methods. The same field of cells was photographed by Nomarski optics (left) and by indirect immunofluorescence to detect Vcx1p (center) and to detect DNA after staining with 4,6'-diamidino-2-phenylindole (DAPI; right).

added  $\text{Ca}^{2+}$  ( $\text{IC}_{50}$ ,  $\sim 8$  mM). The *cnb1 vcx1* double mutant is also more sensitive to  $\text{Ca}^{2+}$  ( $\text{IC}_{50}$ ,  $\sim 100$  mM) than a *cnb1* single mutant ( $\text{IC}_{50}$ ,  $\sim 370$  mM), which indicates that *VCX1* performs a significant role in  $\text{Ca}^{2+}$  tolerance in calcineurin-deficient mutants, independently of *PMCI*. Similar results were obtained in cultures in which calcineurin is inhibited by addition of  $0.2 \mu\text{g}$  of FK506 per ml (not shown) or inactivated

by the *cmd1-3* mutation (Fig. 6), which produces a mutant calmodulin that is both unable to bind  $\text{Ca}^{2+}$  with high affinity (22) and unable to activate calcineurin in vivo (14). Expression of a calmodulin-independent derivative of the calcineurin catalytic subunit *CNA1 $\Delta$ C* decreases  $\text{Ca}^{2+}$  tolerance of *cmd1-3 pmc1* mutants (14) and increases  $\text{Ca}^{2+}$  tolerance of *cmd1-3 vcx1* mutants (data not shown), suggesting that the major role




FIG. 4. Ca<sup>2+</sup> transport assays in yeast cell extracts. (A) Total spheroplast lysates were prepared from mid-log-phase YPD cultures of the wild-type (WT) strain W303-1A and the *vcx1* mutant K661 and assayed for uptake of <sup>45</sup>Ca<sup>2+</sup> with and without the inhibitor of vacuolar H<sup>+</sup> ATPase concanamycin B (CB). After 3 min of preincubation at 30°C in buffer containing an optimal level of <sup>45</sup>Ca<sup>2+</sup> (100 μM), 1 mM ATP was added at 0 min, and aliquots were then removed at the indicated times, rapidly diluted in cold buffer, filtered, and processed as described in Materials and Methods. Ca<sup>2+</sup> accumulation in nanomoles per 10<sup>9</sup> cell equivalents was calculated (without subtracting the prior ATP-independent Ca<sup>2+</sup> accumulation). (B) Total cell lysates were prepared from a *vcx1 pmc1* double mutant (K665) containing plasmid pRS316-*gal* (open squares) or pKC151-*GAL::PMC1* (filled squares). Extracts were preincubated with <sup>45</sup>Ca<sup>2+</sup> (10 μM) for 5 min at 30°C and split into two equal portions, and then reactions were initiated at 0 min by addition of 1 mM ATP to one half. Samples were collected and processed as described above, and ATP-dependent Ca<sup>2+</sup> uptake was calculated by subtracting the ATP-independent Ca<sup>2+</sup> accumulation from the total Ca<sup>2+</sup> accumulation.

of calmodulin in Ca<sup>2+</sup> tolerance is to activate calcineurin. The simplest interpretation of all of these results is that calcineurin activation by Ca<sup>2+</sup>/calmodulin diminishes the contribution of *VCX1* in Ca<sup>2+</sup> tolerance and Ca<sup>2+</sup> sequestration.

Previous studies showed that *pmc1 pmr1* double mutants were inviable at all levels of Ca<sup>2+</sup> but were viable and very Ca<sup>2+</sup> tolerant when calcineurin was inactivated by low levels of FK506 or by *cnb1* mutations (14). If calcineurin-dependent inhibition of *VCX1* is responsible for the lethality of *pmc1 pmr1* double mutants, then deletion of *VCX1* should be lethal in the *cnb1 pmc1 pmr1* strain. Tetrad analysis of heterozygous diploids showed that all combinations of these mutations were efficiently recovered as viable clones, except for *pmc1 pmr1* double mutants, *pmc1 pmr1 vcx1* triple mutants, and *cnb1 pmc1 pmr1 vcx1* quadruple mutants. This experiment was reproducible over a wide range of conditions. Thus, *VCX1* function is required for the viability of the *cnb1 pmc1 pmr1* triple mutant, and *VCX1* cannot perform its essential function when calcineurin is active. Attempts to reconstitute calcineurin-de-




FIG. 5. Nonexchangeable Ca<sup>2+</sup> pools in yeast mutants. Log-phase yeast cultures were uniformly labeled in YPD pH 5.5 medium supplemented with 5 mM <sup>45</sup>Ca<sup>2+</sup> and subjected to a brief chase period in nonradioactive medium to allow efflux of the exchangeable Ca<sup>2+</sup> pool. The remaining nonexchangeable pool was averaged from duplicate cultures as described in Materials and Methods. Standard deviations from the duplicate samples ranged from 0.1 to 0.3 nmol/10<sup>9</sup> cells. Bars 1 to 6 show data for strains W303-1A, K661, K605, K665, K607, and K667, respectively. +, wild-type alleles.

pendent inactivation of vacuolar H<sup>+</sup>/Ca<sup>2+</sup> exchange in vitro have not yet succeeded. However, evidence presented below suggests that calcineurin may slightly decrease the expression of *VCX1* as well as the activity of Vcx1p in vivo.

**Production of Vcx1p and Pmc1p.** Our results indicate that calcineurin not only diminishes the role of *VCX1* in Ca<sup>2+</sup> tolerance but also can promote the role of other Ca<sup>2+</sup> tolerance factors (for example, *cnb1 vcx1* double mutants are less tolerant of Ca<sup>2+</sup> than *vcx1* mutants). To determine whether calcineurin differentially modulates expression of *VCX1* and *PMC1*, total cell extracts prepared after growth in different conditions were analyzed by Western blotting. The predicted




FIG. 6. Ca<sup>2+</sup> tolerance in yeast mutants. The IC<sub>50</sub> for each strain was determined after 16 h of growth in supplemented YPD pH 5.5 medium as described in Materials and Methods. Similar results were obtained in independent experiments and when FK506 (0.2 μg/ml) was used in lieu of *cnb1* mutations. All strains listed grew at similar rates in standard unsupplemented media (YPD and SC). Additionally, the epitope-tagged *VCX1::HA* allele integrated at the chromosomal locus was indistinguishable from *VCX1* in similar situations. +, wild-type allele.




FIG. 7. Expression of Vcx1p and Pmc1p in response to  $\text{Ca}^{2+}$  and FK506. (A) Western blot analysis of *PMCI::HA* (lanes 1 to 5, strain K699) and *PMCI* (lanes 6 and 7, strain W303-1A) grown for 6 h in YPD pH 5.5 medium (lanes 1 and 6) or in medium supplemented with 50 mM  $\text{CaCl}_2$  (lane 2), 100 mM  $\text{CaCl}_2$  (lane 3), 200 mM  $\text{CaCl}_2$  (lane 4), or 200 mM  $\text{CaCl}_2$  plus 0.4  $\mu\text{g}$  of FK506 per ml (lanes 5 and 7). Total cell extracts were prepared as described in Materials and Methods, and samples corresponding to 1.25  $\text{OD}_{600}$  units were analyzed on 8% polyacrylamide gels adjacent to molecular weight markers (positions shown at the left in kilodaltons). Pmc1p and cross-reacting polypeptides p180 and p170 are indicated at the right. (B) Western blot analysis of *VCXI::HA* (strain K698) grown for 6 h in YPD pH 5.5 medium (lane 1) or medium supplemented with 200 mM  $\text{CaCl}_2$  (lane 2), 0.4  $\mu\text{g}$  of FK506 per ml (lane 3), or 200 mM  $\text{CaCl}_2$  plus 0.4  $\mu\text{g}$  of FK506 per ml (lane 4). Samples corresponding to 0.3  $\text{OD}_{600}$  unit were run on 10% polyacrylamide gels as described above. No cross-reacting polypeptides other than Vcx1p were evident under these conditions.

49-kDa product of *VCXI::HA* was observed as a series of bands migrating at 39 to 43 kDa during growth in YPD pH 5.5 medium (Fig. 7B, lane 1), which decreased very slightly (two-fold or less) during growth in 200 mM  $\text{Ca}^{2+}$  (lane 2) but did not noticeably change in abundance or mobility in response to FK506 or  $\text{Ca}^{2+}$  plus FK506 (lanes 3 and 4). The predicted 135-kDa product of *PMCI::HA* increased markedly in abundance as  $\text{Ca}^{2+}$  in the growth medium was increased from 0 to 200 mM (Fig. 7A, lanes 1 to 4), and this effect was blocked when FK506 was present (lane 5). High  $\text{Ca}^{2+}$  conditions also caused the FK506-sensitive accumulation of a 180-kDa polypeptide in both *PMCI::HA* and wild-type strains, whereas a second cross-reacting polypeptide of  $\sim 170$  kDa was unchanged (Fig. 7A). High- $\text{Ca}^{2+}$  conditions therefore decrease Vcx1p accumulation only slightly and greatly increase Pmc1p accumulation, both through FK506-sensitive mechanisms.

**Expression of *VCXI*, *PMCI*, *PMRI*, and *PMR2A* reporter genes is modulated by calcineurin activation in response to elevated  $[\text{Ca}^{2+}]_c$ .** *PMCI* gene expression was also quantitated by using a series of plasmid-based reporter genes in which *lacZ* coding sequences from *E. coli* were placed downstream of the promoter and predicted N-terminal codons of these genes. All yeast strains containing the *pmc1::lacZ* reporter plasmid (pKC190) accumulated very low levels of  $\beta$ -galactosidase activity (0.1 to 0.2 U) when grown in standard media (Fig. 8A). After 4.25 h of incubation in medium supplemented with  $\text{Ca}^{2+}$ , expression of *pmc1::lacZ* increased approximately 150-fold in wild-type strains, 200-fold in *vcx1* mutants, 500-fold in *pmc1* mutants, and 650-fold in *pmc1 vcx1* mutants (Fig. 8A), whereas expression in *cnb1 vcx1* double mutants did not increase during growth in  $\text{Ca}^{2+}$ -supplemented medium. Similarly, increased  $\beta$ -galactosidase accumulation was blocked by addition of 0.2  $\mu\text{g}$  of FK506 per ml (Table 2). A *cyc1::lacZ* reporter used previously to analyze expression of cytochrome *c* (24) was slightly repressed by these conditions ( $\sim 1.1$ -fold [Table 2]). To help rule out the possibility of  $\text{Ca}^{2+}$  and calcineurin-responsive elements in the parent vector, we analyzed derivatives of *cyc1::lacZ* that lack all upstream activation sequences (24) or have those sequences replaced with a cell cycle-regulated upstream activation sequence (UAS) derived from the *HO* gene (10). There was no detectable expression of the UAS-deficient

*cyc1::lacZ* derivative under any of our conditions, and there was only  $\sim 1.3$ -fold repression of the *ho::cyc1::lacZ* reporter by added  $\text{Ca}^{2+}$  (Table 2). Finally, a 413-bp fragment of the *PMCI* upstream region (from  $-579$  to  $-166$  relative to initiator codon) was found to be necessary and sufficient for calcineurin-dependent induction by  $\text{Ca}^{2+}$  because deletion of this region from *pmc1::lacZ* abolished all expression and insertion of this segment upstream of the UAS-deficient *cyc1::lacZ* reporter restored full  $\text{Ca}^{2+}$ - and calcineurin-dependent expression of  $\beta$ -galactosidase (Table 2). These results suggest that the *PMCI* gene can be regulated at the transcriptional level through upstream activation sequences which depend on calcineurin activation.

Expression of *vcx1::lacZ*, *pmr1::lacZ*, and *pmr2A::lacZ* reporters was also examined. The *vcx1::lacZ* reporter was repressed 2.2-fold and *pmr1::lacZ* was induced 2.3- to 2.9-fold by calcineurin-dependent mechanisms in response to high- $\text{Ca}^{2+}$  conditions (Fig. 8C and Table 2). Lastly, a *pmr2A::lacZ* reporter was induced over 25-fold by the actions of  $\text{Ca}^{2+}$  and calcineurin (Table 2) and up to 500-fold in some conditions (Fig. 8B). As for *pmc1::lacZ*, calcineurin-dependent expression of the *pmr2A::lacZ* reporter was exacerbated in the *pmc1* and *pmc1 vcx1* mutants; however, the range of added  $\text{Ca}^{2+}$  necessary to induce *pmr2A::lacZ* expression was much lower than that of *pmc1::lacZ* (Fig. 8). Other differences between the two reporters in the different yeast strains can be observed in the highest- $\text{Ca}^{2+}$  conditions. These results suggest that Pmc1p and to a lesser degree Vcx1p serve to lower  $[\text{Ca}^{2+}]_c$  in high- $\text{Ca}^{2+}$  conditions and thereby decrease calcineurin activation. There was no detectable induction of either *pmc1::lacZ*, *pmr1::lacZ*, or *pmr2A::lacZ* in *cmd1-3* mutants (data not shown). Thus, activation of calcineurin by calmodulin and elevated  $[\text{Ca}^{2+}]_c$  can differentially affect the expression of at least four reporter genes. No conserved sequence elements are obvious in the promoter regions of these genes. Calcineurin-dependent repression of *VCXI* and induction of *PMCI* and *PMRI* provide at least partial explanations for all the observed effects of different mutations on  $\text{Ca}^{2+}$  tolerance (Fig. 6).

**Calmodulin and calcineurin mediate responses to  $\text{Mn}^{2+}$  and  $\text{Na}^+$ .** Because *PMRI* and *PMR2A* have been implicated in tolerance to added  $\text{Mn}^{2+}$  and  $\text{Na}^+$ , respectively (28, 37), we tested whether calcineurin may be involved in tolerance to these ions. Inactivation of calcineurin with mutations or FK506 also causes a significant decrease in tolerance to  $\text{Mn}^{2+}$  and  $\text{Na}^+$  (Fig. 9 and references 40 and 45). To test whether the consequences of calcineurin inactivation are dependent on *PMRI* and *PMR2A* function, we compared  $\text{Mn}^{2+}$  and  $\text{Na}^+$  tolerances in *pmr1* and *pmr2* null mutants both with and without a functional calcineurin. Addition of FK506 to the growth medium markedly decreased  $\text{Mn}^{2+}$  tolerance of wild-type and *vcx1* mutants but had no significant effect on the tolerance of *pmr1* and *pmr1 vcx1* mutants (Fig. 9A). The nonadditive effect of FK506 plus *pmr1* mutations suggests that the role of calcineurin in  $\text{Mn}^{2+}$  tolerance is dependent on *PMRI* function. A similar experiment to assess  $\text{Na}^+$  tolerance of *pmr2* mutants (which lack *PMR2A* and three additional repeats at this locus) with and without the *cnb1* null mutation (Fig. 9B) showed that there was only a slight additive effect of the two mutations. The introduction of *vcx1* mutations had only small quantitative effects on the  $\text{Mn}^{2+}$  and  $\text{Na}^+$  tolerance of these strains, indicating that *VCXI* contributes little to  $\text{Mn}^{2+}$  and  $\text{Na}^+$  tolerance with or without calcineurin function. These results are consistent with a model whereby calcineurin promotes  $\text{Mn}^{2+}$  and  $\text{Na}^+$  tolerance through effects on *PMRI* and *PMR2A* (and not *VCXI*), but other possible targets of calcineurin are not ruled out.




FIG. 8. Expression of *lacZ* reporters for *PMCI*, *PMR2A*, *PMR1*, and *VCX1* in response to added  $\text{Ca}^{2+}$  and in different mutant strains. Yeast strains were transformed with plasmids pKC190 (A) and pKC201 (B), grown to mid-log phase in SC-Ura medium, collected, and suspended in YPD pH 5.5 medium plus  $\text{CaCl}_2$  as indicated. After 4.25 h of incubation at 30°C with shaking, total cellular  $\beta$ -galactosidase activity was determined for each culture. Strains: WT (wild type), W303-1A; *pmc1*, K605; *vcx1*, K661; *cnb1 vcx1*, K663; *pmc1 vcx1*, K665; *pmc1 VCX1-D1*, K737. (C) Plasmids pKC199 (*pmr1::lacZ*) and pKC200 (*vcx1::lacZ*) were transformed into the *vcx1* mutant K661, incubated, and processed as described above except that

TABLE 2. Expression of reporter genes in *vcx1* mutants

Plasmid	Reporter	$\beta$ -Galactosidase (U) <sup>a</sup>			Ratio <sup>b</sup> (+Ca/ +Ca+FK)
		YPD	+Ca	+Ca+FK	
pKC190	<i>pmc1::lacZ</i>	<0.2	13.8	<0.2	69 ± 10
pLGΔ312	<i>cyc1::lacZ</i>	494.2	391.1	445.8	0.88 ± 0.07
pLGΔ178	( $\Delta$ UAS) <i>cyc1::lacZ</i>	<0.2	<0.2	<0.2	
pLB178-43	<i>ho::cyc1::lacZ</i>	47.4	41.4	53.0	0.78 ± 0.07
pKC211	( $\Delta$ 13) <i>pmc1::lacZ</i>	<0.2	<0.2	<0.2	
pKC191	(413) <i>pmc1::cyc1::lacZ</i>	0.5	13.5	0.3	52 ± 7
pKC200	<i>vcx1::lacZ</i>	12.9	9.7	21.9	0.44 ± 0.07
pKC199	<i>pmr1::lacZ</i>	22.3	41.9	18.3	2.3 ± 0.1
pKC201	<i>pmr2A::lacZ</i>	0.9	59.2	2.2	27 ± 4

<sup>a</sup> Total cellular  $\beta$ -galactosidase activity was assayed as described in Materials and Methods after 4 h of growth in YPD pH 5.5 medium supplemented as indicated with 100 mM  $\text{CaCl}_2$  (Ca) and 0.2  $\mu\text{g}$  of FK506 (FK) per ml. Data are averages of two independent experiments; 0.2 U was the limit of detection.

<sup>b</sup> Average of two independent experiments ( $\pm$  standard deviation).

Is calcineurin required for induction of the *PMR1* and *PMR2A* reporter genes in response to  $\text{Mn}^{2+}$  and  $\text{Na}^+$ ? Expression of the *pmr1::lacZ* reporter was induced significantly in *vcx1* mutants, but not *cmd1-3 vcx1* mutants, after a shift to media supplemented with 1 mM  $\text{MnCl}_2$  (Fig. 10A). Addition of FK506 to the  $\text{MnCl}_2$ -supplemented cultures decreased expression by approximately 2.5-fold in *vcx1* mutants but had no significant effect in *vcx1 cmd1-3* mutants. There was also no significant FK506-sensitive induction of *ho::lacZ* in *vcx1* mutants, indicating that calmodulin and calcineurin are required for the specific induction of *pmr1::lacZ* in response to  $\text{Mn}^{2+}$ . Similarly, *pmr2A::lacZ* expression was induced by 500 mM NaCl in *pmc1* mutants, and induction was markedly decreased either by FK506 addition (Fig. 10B) or by introduction of *cmd1-3* or *cnb1* mutations (data not shown). Qualitatively similar results were obtained when wild-type strains, *vcx1* mutants, and *pmc1 vcx1* mutants were each analyzed, although the relative levels of calcineurin-dependent induction appeared to increase as  $\text{Ca}^{2+}$  transporters were inactivated (data not shown). Significant calcineurin-dependent induction of  $\text{Na}^+$  stress has been shown previously to increase accumulation of *PMR2A* mRNAs in wild-type yeast strains and to a lesser degree in *cnb1* mutants (20). Thus, functions of calmodulin and calcineurin may be involved in the induction of *PMR1* and *PMR2A* gene expression in response to elevated extracellular  $\text{Mn}^{2+}$  and  $\text{Na}^+$ . These findings provide at least one plausible explanation for the decreased  $\text{Mn}^{2+}$  and  $\text{Na}^+$  tolerance of calcineurin mutants.

**Activated variants of Vcx1p decrease  $\text{Ca}^{2+}$  signaling.** Our results suggest that  $[\text{Ca}^{2+}]_c$  becomes elevated in response to added  $\text{Ca}^{2+}$ ,  $\text{Mn}^{2+}$ , or  $\text{Na}^+$  in order to promote the calcineurin-dependent induction of specific tolerance factors. The latter two responses may require inactivation of Vcx1p for efficient production of  $\text{Ca}^{2+}$  signals. To test this idea, we isolated *VCX1-D* mutants which are active despite the function of calcineurin and tested whether such mutants would diminish the calcineurin-dependent induction of *pmr2A::lacZ* in response to  $\text{Na}^+$ . In our original screen for spontaneous  $\text{Ca}^{2+}$ -tolerant revertants of *pmc1* strains, recessive mutations comprising three complementation groups were isolated (14). Six additional revertants were dominant over wild-type *VCX1* for  $\text{Ca}^{2+}$  tolerance, segregated 2:2 in backcrosses, and segregated

FK506 (0.5  $\mu\text{g}/\text{ml}$ ) was included in half of the YPD cultures (+FK). All patterns of expression were reproducible in independent experiments.






FIG. 9. Roles of calcineurin in  $Mn^{2+}$  and  $Na^{+}$  tolerance. The indicated yeast strains were inoculated into YPD pH 5.5 medium supplemented with  $0.5 \mu\text{g}$  of FK506 (FK) per ml,  $MnCl_2$ , or NaCl as indicated. After 16 h of growth at  $30^\circ\text{C}$ , culture density and  $IC_{50}$  were determined by interpolation as described in Materials and Methods. +, wild-type allele. Similar results were obtained in three independent experiments.

4:0 in crosses with each other. Crosses between these strains and *vcx1::URA3* demonstrated tight linkage of the dominant mutations to the *VCX1* locus. We recovered all of the dominant mutations on plasmids by using an allele rescue strategy (26) with *StuI*-digested plasmid pKC106. Subcloning and sequencing demonstrated that point mutations within the *VCX1* open reading frame, all of which resulted in amino acid substitutions, were responsible for the  $Ca^{2+}$  tolerance phenotype. A *pmc1 VCX1-D1* double mutant was also found to sequester high levels of  $Ca^{2+}$  in the nonexchangeable pool (approximately 10 times that of *pmc1* mutants; data not shown) and to block partially the induction of *pmc1::lacZ* and *pmr2A::lacZ* in response to added  $Ca^{2+}$  (Fig. 8). Complete sequencing of *VCX1-D1* revealed a single nucleotide change, G-1344→A, which changes codon 383 from Met to Ile in the 11th putative transmembrane domain. These results suggest that *VCX1-D1* produces a variant vacuolar  $H^{+}/Ca^{2+}$  exchanger that is very active in strains in which calcineurin is nominally functional.

Calcineurin-dependent induction of *pmr2A::lacZ* by  $Na^{+}$  was also found to be much less in *pmc1 VCX1-D1* mutants than in *pmc1* mutants (Fig. 10B). There was no significant change, however, in the calcineurin-independent (FK506-resistant) induction of *pmr2A::lacZ* by  $Na^{+}$ . Thus, inappropriately high activity of the vacuolar  $H^{+}/Ca^{2+}$  exchanger decreases a number of calcineurin-dependent responses. *PMCI* function also appears to modulate calcineurin-dependent gene expression (Fig. 8) presumably through effects on  $[Ca^{2+}]_c$ . We conclude that calcineurin activation by  $Ca^{2+}$ /calmodulin is a physiological response to high  $Ca^{2+}$ ,  $Mn^{2+}$ , and  $Na^{+}$  concentrations which regulates the appropriate detoxification mechanisms. Feedback interactions between calcineurin and vacuolar  $Ca^{2+}$

transporters may ensure that such processes occur efficiently over a wide range of conditions.

## DISCUSSION

**$Ca^{2+}$  homeostasis in *S. cerevisiae*.** Our previous work suggested that calcineurin might inhibit a factor required for  $Ca^{2+}$  tolerance of *pmc1* mutants (14). Here we have identified *VCX1*, encoding the major vacuolar  $H^{+}/Ca^{2+}$  exchanger, as a direct or indirect target of calcineurin required for this effect. The simplest model consistent with the new results is that  $Ca^{2+}$  homeostasis in yeast cells is achieved by a sophisticated feedback mechanism involving at least calmodulin, calcineurin, and three intracellular  $Ca^{2+}$  transporters encoded by *VCX1*, *PMCI*, and *PMRI* (Fig. 11). Together these  $Ca^{2+}$  transporters control  $[Ca^{2+}]_c$  and appear to prevent the toxic accumulation of  $Ca^{2+}$  in the cytosol, which is especially important in conditions of high extracellular  $Ca^{2+}$  concentrations. We propose that calcineurin activation by calmodulin and elevated  $[Ca^{2+}]_c$  leads to increased expression of *PMCI* and possibly *PMRI* and to slightly decreased expression of *VCX1*. It is more likely that calcineurin inactivates Vcx1p by a posttranslational mechanism because the repression of Vcx1p expression seems insufficient




FIG. 10. Calcineurin mediates responses to  $Mn^{2+}$  and  $Na^{+}$ . (A) Expression of *pmc1::lacZ* or *ho::lacZ* reporters in yeast strains K661 (*vcx1*) and K669 (*vcx1 cmd1-3*) after 4 h of growth in YPD pH 5.5 medium (+0) or in the same medium supplemented with 1 mM  $MnCl_2$  (+Mn) and  $0.2 \mu\text{g}$  of FK506 per ml (+Mn +FK).  $\beta$ -Galactosidase activity from independent cultures was determined, and all values were normalized to that of the unsupplemented cultures (arbitrarily set at 100 relative units), averaged, and plotted as bars. Error markers indicate standard deviations from  $n$  independent experiments. (B) Expression of *pmr2A::lacZ* in *pmc1* (K605) and *pmc1 VCX1-D1* (K737) strains was determined after 4 h of growth in YPD pH 5.5 medium supplemented with the indicated amounts of NaCl (filled symbols, solid lines) and with  $0.4 \mu\text{g}$  of FK506 (FK) per ml (open symbols, dashed lines).




FIG. 11. Working model of  $\text{Ca}^{2+}$  homeostasis in *S. cerevisiae*. Lines with arrowheads indicate positive functions or activating interactions, and lines with bars indicate inhibitory interactions. Question marks denote unknown factors or interactions that may be indirect.  $[\text{Ca}^{2+}]_o$ , extracellular free  $\text{Ca}^{2+}$  concentration.

to account for the dramatic effects of calcineurin on *VCX1*-dependent  $\text{Ca}^{2+}$  tolerance and sequestration. Although the exact mechanisms by which calcineurin accomplishes all these functions are not yet clear, this model provides a plausible explanation for why calcineurin function decreases  $\text{Ca}^{2+}$  tolerance of *pmc1* mutants (14) and increases  $\text{Ca}^{2+}$  tolerance of *vcx1* mutants and *pmc1 vcx1* double mutants (Fig. 6).

Mutant strains lacking all three  $\text{Ca}^{2+}$  transporters (*VCX1*, *PMC1*, and *PMR1*) are inviable in all growth media tested to date. However, mutants containing only a single functional  $\text{Ca}^{2+}$  transporter gene are viable and provide insight into the relative roles of each  $\text{Ca}^{2+}$  transporter and its regulators. For example, *PMR1* provides functions essential for the viability of *cnb1 pmc1 vcx1* triple mutants, which implies that *PMR1* retains significant activity in the absence of calcineurin activation. This finding is consistent with the relatively high basal expression of *pmr1::lacZ* (Fig. 8) and with the known secretory roles of *PMR1* in low- $\text{Ca}^{2+}$  conditions (4, 57). The viability of *cnb1 pmr1 vcx1* triple mutants suggests that basal expression of *PMC1* is also sufficient to carry out essential functions. Furthermore, the inviability of *pmc1 pmr1* double mutants and viability and high  $\text{Ca}^{2+}$  tolerance of *cnb1 pmc1 pmr1* triple mutants (14) and of *pmc1 pmr1 VCX1-D1* triple mutants (data not shown) can be interpreted to indicate that *VCX1* can provide the essential functions when the inhibitory effects of calcineurin are removed. Inhibition of *VCX1* function by calcineurin is somewhat incomplete because residual activity of *VCX1* was detected in *pmc1* mutants (Fig. 6). Thus, all three transporters appear to function to some degree with or without calcineurin activation. In  $\text{Ca}^{2+}$  tolerance assays, *PMC1* provides the largest contribution whereas *VCX1* and *PMR1* ordinarily contribute much less.

In *S. cerevisiae*,  $\text{Ca}^{2+}$ /calmodulin regulates the intracellular  $\text{Ca}^{2+}$  transporters indirectly through effects on calcineurin. Animal cells, in contrast, regulate the plasma membrane  $\text{Ca}^{2+}$  ATPases and both the retinal and cardiac  $\text{Na}^+/\text{Ca}^{2+}$  exchangers by direct binding of  $\text{Ca}^{2+}$ /calmodulin and also by phosphorylation (for reviews, see references 12 and 52); regulation by calcineurin has not been reported. Sequence alignments indi-

cate that the calmodulin-binding regulatory domains of mammalian plasma membrane  $\text{Ca}^{2+}$  ATPases and the retinal  $\text{Na}^+/\text{Ca}^{2+}$ ,  $\text{K}^+$  exchangers are not conserved in *Pmc1p* and *Vcx1p* (Fig. 2 and reference 14). We observed that calcineurin-dependent inhibition of *VCX1* function and promotion of *PMC1* function are both abolished by *cmd1-3* mutations except when the calmodulin-independent *CNA1ΔC* derivative of calcineurin is expressed (14). Thus, yeast cells differ from mammalian cells in that the vacuolar  $\text{Ca}^{2+}$  transporters are regulated predominantly by calcineurin activation rather than direct binding of  $\text{Ca}^{2+}$ /calmodulin. Although our model fits the extant data, it does not rule out other possible interactions among these factors. For example, all *cmd1-3* strains and *cmd1-3 cnb1* double mutants are less  $\text{Ca}^{2+}$  tolerant than corresponding *cnb1* mutants (Fig. 6 and data not shown), suggesting that  $\text{Ca}^{2+}$ /calmodulin may have some other roles in  $\text{Ca}^{2+}$  tolerance that are independent of calcineurin, *PMC1* and *VCX1*.

Kinetic analyses of the vacuolar  $\text{H}^+/\text{Ca}^{2+}$  exchanger from *S. cerevisiae* suggested that despite a relatively high  $K_m$  for  $\text{Ca}^{2+}$ , this enzyme would be sufficient to account for the levels of  $\text{Ca}^{2+}$  sequestration observed over a wide range of environmental conditions (17). Similar conclusions have been reached for vacuolar  $\text{H}^+/\text{Ca}^{2+}$  exchangers of filamentous fungi and higher plants (8, 41). These considerations, taken together with the finding that *vma1* mutants (and other mutants deficient in the vacuolar  $\text{H}^+/\text{Ca}^{2+}$  exchange activity) are extremely sensitive to added  $\text{Ca}^{2+}$ , have led to the prediction that mutants lacking the  $\text{H}^+/\text{Ca}^{2+}$  exchanger would similarly be unable to grow in high- $\text{Ca}^{2+}$  conditions (3). Our results show that *vcx1* mutants lack the major vacuolar  $\text{H}^+/\text{Ca}^{2+}$  exchanger but are only slightly less tolerant of  $\text{Ca}^{2+}$  than wild-type cells and much more  $\text{Ca}^{2+}$  tolerant than *pmc1* mutants (Fig. 6) and *vma1* mutants (49). Furthermore, *VCX1* contributes little to  $\text{Ca}^{2+}$  sequestration in the range of 0.2 to 20 mM extracellular  $\text{Ca}^{2+}$  (Fig. 5 and data not shown). We presume that the optimized assay for  $\text{H}^+/\text{Ca}^{2+}$  exchange activity (Fig. 4A) does not effectively reconstitute calcineurin-dependent inhibition of *VCX1* function because wild-type and *cnb1* extracts were found to be indistinguishable (data not shown). The model described above does not explain why *vma1* mutants are sensitive to  $\text{Ca}^{2+}$ , but it predicts that *PMC1* and *PMR1* would be important for the residual  $\text{Ca}^{2+}$  tolerance of *vma1* mutants. Consistent with this prediction, *pmr1 vma1* double mutants are not viable (62) and *cnb1 vma1* double mutants are either extremely  $\text{Ca}^{2+}$  sensitive or inviable (21, 30, 62) as a result of decreased expression of *PMC1* and *PMR1* or other effects.

Tanida et al. propose that calcineurin may inhibit an intracellular  $\text{Ca}^{2+}$  transporter distinct from the vacuolar  $\text{H}^+/\text{Ca}^{2+}$  exchanger and that its increased function after FK506 addition may cause decreased  $\text{Ca}^{2+}$  tolerance in *vma* mutants (62). They observed that FK506 addition increased  $\text{Ca}^{2+}$  sequestration into the nonexchangeable pool and rapidly decreased  $[\text{Ca}^{2+}]_c$ . These results do not rule out *VCX1* as the target of calcineurin inhibition because a small pH gradient persists in the vacuoles of *vma* mutants (43), which may be sufficient to promote vacuolar  $\text{H}^+/\text{Ca}^{2+}$  exchange activity. That FK506 or *cnb1* mutations cause decreased  $\text{Ca}^{2+}$  tolerance of *vma* mutants (21, 30, 62) can also be explained by a variety of secondary defects in cell physiology and not necessarily changes in  $\text{Ca}^{2+}$  transport. For example, a drastic reduction of vacuolar polyphosphate levels in *vma* mutants (67) could lead to decreased  $\text{Ca}^{2+}$  tolerance simply by lowering the  $\text{Ca}^{2+}$  buffering in this organelle (17) irrespective of effects on  $\text{Ca}^{2+}$  transport. Furthermore, *vma* mutants display pleiotropic defects in ion

homeostasis, sorting of vacuolar proteins, general metabolism, and other processes (reviewed in references 3 and 34) which might affect  $\text{Ca}^{2+}$  tolerance indirectly. Additional studies are necessary to determine how *vma* mutants and other  $\text{Ca}^{2+}$ -sensitive mutants (6, 48) affect the vacuolar  $\text{Ca}^{2+}$  transporters encoded by *VCX1* and *PMCI*.

Increased expression of  $\text{Ca}^{2+}$  ATPases would benefit the cell in high- $\text{Ca}^{2+}$  conditions to offset increased  $\text{Ca}^{2+}$  influx. At present, it is more difficult to appreciate how the cell benefits from *VCX1* when its function seems to be strongly inhibited by calcineurin activation. Because calcineurin and *VCX1* appear to regulate each other in a positive feedback loop, it is not possible to conclude whether *VCX1-D1* produces a  $\text{H}^+/\text{Ca}^{2+}$  exchanger that is specifically resistant to inhibition by calcineurin or whether some other aspect of its function or expression is augmented. The decreased function of calcineurin in *VCX1-D1* strains is consistent with the hypothesis that inhibition of *VCX1* is important to promote the formation of  $\text{Ca}^{2+}$  signals in *S. cerevisiae*. Presumably *VCX1* functions more significantly during conditions which have yet to be identified.

**$\text{Ca}^{2+}$  signaling in *S. cerevisiae*.** We report here the calcineurin-dependent induction of *pmr1::lacZ* and *pmr2A::lacZ* in response to added  $\text{Mn}^{2+}$  and  $\text{Na}^+$  as well as to added  $\text{Ca}^{2+}$  (Fig. 8 and 10). The *cnb1* and *cmd1-3* mutations each block these regulatory effects, suggesting that calcineurin activation by calmodulin may be occurring in response to an increase in  $[\text{Ca}^{2+}]_c$ . Because *PMR1* and *PMR2A* encode  $\text{Mn}^{2+}$  and  $\text{Na}^+$  tolerance factors (28, 37), their calmodulin- and calcineurin-dependent induction in response to increased environmental levels of these ions may indicate more physiological means of generating  $\text{Ca}^{2+}$  signals in yeast cells. Their induction by moderate- $\text{Ca}^{2+}$  conditions, as opposed to the high- $\text{Ca}^{2+}$  conditions needed for maximal *PMCI* induction, does not necessarily imply roles in  $\text{Ca}^{2+}$  tolerance but may reflect responsiveness to relatively small increases in  $[\text{Ca}^{2+}]_c$ . Introduction of *pmr2* null mutations into *pmc1*, *cnb1*, *vcx1*, and *cnb1 pmc1 vcx1* mutants does not significantly affect  $\text{Ca}^{2+}$  tolerance (data not shown). Possible means of generating  $\text{Ca}^{2+}$  signals in response to  $\text{Na}^+$  include the gated opening of putative  $\text{Ca}^{2+}$ -selective channels in the vacuole membrane (7) or plasma membrane (25, 31).

**Relevance to other cell types.** There is an impressive degree of conservation among *S. cerevisiae* and animals in the mechanisms of calcineurin activation by  $\text{Ca}^{2+}$  and calmodulin and inhibition by immunosuppressive drugs and their immunophilin receptors (36). Our finding that yeast calcineurin can regulate expression of specific genes has parallels with the major role of calcineurin in the regulation of gene expression in lymphocytes and other mammalian cells. It will be interesting to determine whether the similarities between *S. cerevisiae* and animals in  $\text{Ca}^{2+}$  signaling mechanisms extend to additional components, such as calcineurin-dependent transcription factors and  $\text{Ca}^{2+}$  channels. This work provides a number of useful tools for these future studies.

#### ACKNOWLEDGMENTS

We thank Kendal Hirschi for comments on the manuscript and Steve Kron for advice. We are grateful to H. L. Ploegh (Massachusetts Institute of Technology) for providing concanamycin B and to Ihor Bekersky (Fujisawa USA, Inc.) and Vicki Sato (Vertex Pharmaceuticals) for generous gifts of FK506. Tabitha Barr, Jaline Han, and Karen Wood provided excellent technical assistance in portions of this study.

K.W.C. is supported by American Cancer Society grant PF-3472. G.R.F. is an American Cancer Society Professor of Genetics. This work was funded in part by a grant from the National Institutes of Health (GM-40266 to G.R.F.) and by private donations from Myco Pharmaceuticals Inc. and The Johns Hopkins University.

#### REFERENCES

- Alani, E., L. Cao, and N. Kleckner. 1987. A method for gene disruption that allows repeated use of *URA3* selection in the construction of multiply disrupted yeast strains. *Genetics* **115**:541–545.
- Altschul, S. F., W. Gish, W. Miller, E. W. Myers, and D. J. Lipman. 1990. Basic local alignment search tool. *J. Mol. Biol.* **215**:403–410.
- Anraku, Y., N. Umemoto, R. Hirata, and Y. Ohya. 1992. Genetic and cell biological aspects of the yeast vacuolar  $\text{H}^+$ -ATPase. *J. Bioeng. Biomembr.* **24**:395–405.
- Antebi, A., and G. R. Fink. 1992. The yeast  $\text{Ca}^{2+}$ -ATPase homologue, *PMR1*, is required for normal Golgi function and localizes in a novel Golgi-like distribution. *Mol. Biol. Cell* **3**:633–654.
- Aperia, A., F. Ibarra, L.-B. Svensson, C. Klee, and P. Greengard. 1992. Calcineurin mediates alpha-adrenergic stimulation of  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase activity in renal tubule cells. *Proc. Natl. Acad. Sci. USA* **89**:7394–7397.
- Beeler, T., K. Gable, C. Zhao, and T. Dunn. 1994. A novel protein, Csg2p, is required for  $\text{Ca}^{2+}$  regulation in *Saccharomyces cerevisiae*. *J. Biol. Chem.* **269**:7279–7284.
- Belde, P. J., J. H. Vossen, G. W. Borst-Pauwels, and A. P. Theuvsenet. 1993. Inositol 1,4,5-triphosphate releases  $\text{Ca}^{2+}$  from vacuolar membrane vesicles of *Saccharomyces cerevisiae*. *FEBS Lett.* **323**:113–118.
- Blackford, S., P. A. Rea, and D. Sanders. 1990. Voltage sensitivity of  $\text{H}^+/\text{Ca}^{2+}$  antiport in higher plant tonoplast suggests a role in vacuolar calcium accumulation. *J. Biol. Chem.* **265**:9617–9620.
- Boeke, J. D., and G. R. Fink. 1984. A positive selection for mutants lacking orotidine-5'-phosphate decarboxylase activity in yeast: 5-fluoro-orotic acid resistance. *Mol. Gen. Genet.* **197**:345–346.
- Breedon, L., and K. Nasmyth. 1987. Cell cycle control of the yeast *HO* gene: *cis*- and *trans*-acting regulators. *Cell* **48**:389–397.
- Brizuela, L., G. Chrebet, K. A. Bostian, and S. A. Parent. 1991. Antifungal properties of the immunosuppressant FK506: identification of an FK506-responsive yeast gene distinct from *FKB1*. *Mol. Cell. Biol.* **11**:4616–4626.
- Carafoli, E. 1992. The  $\text{Ca}^{2+}$  pump of the plasma membrane. *J. Biol. Chem.* **267**:2115–2118.
- Collier, S. J. 1990. Immunosuppressive drugs. *Curr. Opin. Immunol.* **2**:854–858.
- Connelly, C., and P. Hieter. Unpublished data.
- Cunningham, K. W., and G. R. Fink. 1994. Calcineurin-dependent growth control in *Saccharomyces cerevisiae* mutants lacking *PMCI*, a homolog of plasma membrane  $\text{Ca}^{2+}$  ATPases. *J. Cell Biol.* **124**:351–363.
- Cyert, M. S., R. Kunisawa, D. Kaim, and J. Thorner. 1991. Yeast has homologs (*CNA1* and *CNA2* gene products) of mammalian calcineurin, a calmodulin-regulated phosphoprotein phosphatase. *Proc. Natl. Acad. Sci. USA* **88**:7376–7380.
- Cyert, M. S., and J. Thorner. 1992. Regulatory subunit (*CNB1* gene product) of yeast  $\text{Ca}^{2+}$ /calmodulin-dependent phosphoprotein phosphatase is required for adaptation to pheromone. *Mol. Cell. Biol.* **12**:3460–3469.
- Dunn, T., K. Gable, and T. Beeler. 1994. Regulation of cellular  $\text{Ca}^{2+}$  by yeast vacuoles. *J. Biol. Chem.* **269**:7273–7278.
- Foor, F., S. A. Parent, N. Morin, A. M. Dahl, N. Ramadan, G. Chrebet, K. A. Bostian, and J. B. Nielsen. 1992. Calcineurin mediates inhibition by FK506 and cyclosporin of recovery from alpha-factor arrest in yeast. *Nature (London)* **360**:682–684.
- Frohlich, K. U., H. W. Fries, M. Rudiger, R. Erdmann, D. Botstein, and D. Mecke. 1991. Yeast cell cycle protein CDC48p shows full-length homology to the mammalian protein VCP and is a member of a protein family involved in secretion, peroxisome formation, and gene expression. *J. Cell Biol.* **114**:443–453.
- Garcia-deblas, B., F. Rubio, F. J. Quintero, M. A. Banuelos, R. Haro, and A. Rodriguez-Navarro. 1993. Differential expression of two genes encoding isoforms of the ATPase involved in sodium efflux in *Saccharomyces cerevisiae*. *Mol. Gen. Genet.* **236**:363–368.
- Garrett-Engle, P., B. Moilanen, and M. S. Cyert. 1995. Calcineurin, the  $\text{Ca}^{2+}$ /calmodulin-dependent protein phosphatase, is essential in yeast mutants with cell integrity defects and in mutants that lack a functional vacuolar  $\text{H}^+$  ATPase. *Mol. Cell. Biol.* **15**:4103–4114.
- Geiser, J. R., D. van Tuinen, S. E. Brockerhoff, M. M. Neff, and T. N. Davis. 1991. Can calmodulin function without binding calcium? *Cell* **65**:949–959.
- Guarente, L. 1983. Yeast promoters and *lacZ* fusions designed to study expression of cloned genes in yeast. *Methods Enzymol.* **101**:181–191.
- Guarente, L., and T. Mason. 1983. Heme regulates transcription of the *CYC1* gene of *S. cerevisiae* via an upstream activation site. *Cell* **32**:1279–1286.
- Gustin, M. C., X. L. Zhou, B. Martinac, and C. Kung. 1988. A mechanosensitive ion channel in the yeast plasma membrane. *Science* **242**:762–765.
- Guthrie, C., and G. R. Fink (ed.). 1991. *Methods in enzymology*, vol. 194. Guide to yeast genetics and molecular biology. Academic Press, New York.
- Haendler, B., R. Keller, P. C. Hiestand, H. P. Kocher, G. Wegman, and N. R. Movva. 1989. Yeast cyclophilin: isolation and characterization of the protein, cDNA and gene. *Gene* **83**:39–46.
- Haro, R., B. Garcia-deblas, and A. Rodriguez-Navarro. 1991. A novel P-type ATPase from yeast involved in sodium transport. *FEBS Lett.* **291**:189–191.

29. Heitman, J., N. R. Movva, P. C. Hiestand, and M. N. Hall. 1991. FK506-binding protein proline rotamase is a target for the immunosuppressive agent FK506 in *Saccharomyces cerevisiae*. *Proc. Natl. Acad. Sci. USA* **88**: 1948–1952.
30. Hemenway, C. S., K. Dolinski, M. E. Cardenas, M. A. Hiller, E. W. Jones, and J. Heitman. 1995. *VPH6* mutants of *Saccharomyces cerevisiae* require calcineurin for growth and are defective in vacuolar H<sup>+</sup> ATPase assembly. *Genetics* **141**:833–844.
31. Iida, H., H. Nakamura, T. Ono, M. S. Okumura, and Y. Anraku. 1994. *MID1*, a novel *Saccharomyces cerevisiae* gene encoding a plasma membrane protein, is required for Ca<sup>2+</sup> influx and mating. *Mol. Cell. Biol.* **14**:8259–8271.
32. Ivey, D. M., A. A. Guffanti, J. Zemsky, E. Pinner, R. Karpel, E. Padan, S. Schuldiner, and T. A. Krulwich. 1993. Cloning and characterization of a putative Ca<sup>2+</sup>/H<sup>+</sup> antiporter gene from *Escherichia coli* upon functional complementation of Na<sup>+</sup>/H<sup>+</sup> antiporter-deficient strains by the overexpressed gene. *J. Biol. Chem.* **268**:11296–11303.
33. Kinashi, H., K. Someno, and K. Sakaguchi. 1984. Isolation and characterization of concanamycins A, B and C. *J. Antibiot.* **3**:1333–1343.
34. Klionsky, D. J., P. K. Herman, and S. D. Emr. 1990. The fungal vacuole: composition, function, and biogenesis. *Microbiol. Rev.* **54**:266–292.
35. Kuno, T., H. Tanaka, H. Mukai, C. D. Chang, K. Hiraga, T. Miyakawa, and C. Tanaka. 1991. cDNA cloning of a calcineurin B homolog in *Saccharomyces cerevisiae*. *Biochem. Biophys. Res. Commun.* **180**:1159–1163.
36. Kunz, J., and M. N. Hall. 1993. Cyclosporin A, FK506 and rapamycin: more than just immunosuppression. *Trends Biochem. Sci.* **18**:334–338.
37. Lapinskas, P. J., K. W. Cunningham, X. F. Liu, G. R. Fink, and V. C. Culotta. 1995. Mutations in *PMR1* suppress oxidative damage in yeast lacking superoxide dismutase. *Mol. Cell. Biol.* **15**:1382–1388.
38. Liu, H., J. Krizek, and A. Bretscher. 1992. Construction of a GAL1-regulated yeast cDNA expression library and its application to the identification of genes whose overexpression causes lethality in yeast. *Genetics* **132**:665–673.
39. Liu, Y., S. Ishii, M. Tokai, H. Tsutsumi, O. Ohki, R. Akada, K. Tanaka, E. Tsuchiya, S. Fukui, and T. Miyakawa. 1991. The *Saccharomyces cerevisiae* genes (*CMP1* and *CMP2*) encoding calmodulin-binding proteins homologous to the catalytic subunit of mammalian protein phosphatase 2B. *Mol. Gen. Genet.* **227**:52–59.
40. Mendoza, I., F. Rubio, A. Rodriguez-Navarro, and J. M. Pardo. 1994. The protein phosphatase calcineurin is essential for NaCl tolerance of *Saccharomyces cerevisiae*. *J. Biol. Chem.* **269**:8792–8796.
41. Miller, A. J., G. Vogg, and D. Sanders. 1990. Cytosolic calcium homeostasis in fungi: roles of plasma membrane transport and intracellular sequestration of calcium. *Proc. Natl. Acad. Sci. USA* **87**:9348–9352.
42. Mulkey, R. M., S. Endo, S. Shenolikar, and R. C. Malenka. 1994. Involvement of a calcineurin/inhibitor-1 phosphatase cascade in hippocampal long-term depression. *Nature (London)* **369**:486–488.
43. Munn, A. L., and H. Riezman. 1994. Endocytosis is required for the growth of vacuolar H<sup>+</sup> ATPase-defective yeast: identification of six new *END* genes. *J. Cell Biol.* **127**:373–386.
44. Myers, A. M., A. Tzagoloff, D. M. Kinney, and C. J. Lusty. 1986. Yeast shuttle and integrative vectors with multiple cloning sites suitable for construction of *lacZ* fusions. *Gene* **45**:299–310.
45. Nakamura, T., Y. Liu, D. Hirata, H. Namba, S. Harada, T. Hirokawa, and T. Miyakawa. 1993. Protein phosphatase type 2B (calcineurin)-mediated, FK506-sensitive regulation of intracellular ions in yeast is an important determinant for adaptation to high salt stress conditions. *EMBO J.* **12**:4063–4071.
46. Nicoll, D. A., S. Longoni, and K. D. Philipson. 1990. Molecular cloning and functional expression of the cardiac sarcolemmal Na<sup>+</sup>/Ca<sup>2+</sup> exchanger. *Science* **250**:562–565.
47. Ohsumi, Y., and Y. Anraku. 1983. Calcium transport driven by a proton motive force in vacuolar membrane vesicles of *Saccharomyces cerevisiae*. *J. Biol. Chem.* **258**:5614–5617.
48. Ohya, Y., Y. Ohsumi, and Y. Anraku. 1986. Isolation and characterization of Ca<sup>2+</sup>-sensitive mutants of *Saccharomyces cerevisiae*. *J. Gen. Microbiol.* **132**: 979–988.
49. Ohya, Y., N. Umemoto, I. Tanida, A. Ohta, H. Iida, and Y. Anraku. 1991. Calcium-sensitive *cls* mutants of *Saccharomyces cerevisiae* showing a Pet-phenotype are ascribable to defects of vacuolar membrane H<sup>+</sup> ATPase activity. *J. Biol. Chem.* **266**:13971–13977.
50. Okorokov, L. A., T. V. Kulakovskaya, L. P. Lichko, and E. V. Polorotova. 1985. H<sup>+</sup>/ion antiport as the principal mechanism of transport systems in the vacuolar membrane of the yeast *Saccharomyces carlsbergensis*. *FEBS Lett.* **192**:303–306.
51. Parent, S. A., J. B. Nielsen, N. Morin, G. Chrebet, N. Ramadan, A. M. Dahl, J.-J. Hsu, K. A. Bostian, and F. Foor. 1993. Calcineurin-dependent growth of an FK506 and CsA hypersensitive mutant of *Saccharomyces cerevisiae*. *J. Gen. Microbiol.* **139**:2973–2984.
52. Philipson, K. D., and D. A. Nicoll. 1992. Sodium-calcium exchange. *Curr. Opin. Cell Biol.* **4**:678–683.
53. Reid, G. A., and G. Schatz. 1982. Import of proteins into mitochondria. Yeast cells grown in the presence of carbonyl cyanide m-chlorophenylhydrazone accumulate massive amounts of some mitochondrial precursor polypeptides. *J. Biol. Chem.* **257**:13056–13061.
54. Reilander, H., A. Achilles, U. Friedel, G. Maul, F. Lottspeich, and N. J. Cook. 1992. Primary structure and functional expression of the Na/Ca-K-exchanger from bovine rod photoreceptors. *EMBO J.* **11**:1689–1695.
55. Riles, L., J. E. Dutchik, A. Baktha, B. K. McCauley, E. C. Thayer, M. P. Leckie, V. V. Braden, J. E. Depke, and M. V. Olson. 1993. Physical maps of the six smallest chromosomes of *Saccharomyces cerevisiae* at a resolution of 2.6 kilobase pairs. *Genetics* **134**:81–150.
56. Rosoff, P. M., and G. Terres. 1986. Cyclosporin A inhibits Ca<sup>2+</sup>-dependent stimulation of the Na<sup>+</sup>/H<sup>+</sup> antiport in human T cells. *J. Cell Biol.* **103**:457–463.
57. Rudolph, H. K., A. Antebi, G. R. Fink, C. M. Buckley, T. E. Dorman, J. LeVitre, L. S. Davidow, J. I. Mao, and D. T. Moir. 1989. The yeast secretory pathway is perturbed by mutations in *PMR1*, a member of a Ca<sup>2+</sup> ATPase family. *Cell* **58**:133–145.
58. Sambrook, J., E. F. Fritsch, and T. Maniatis. 1989. *Molecular cloning: a laboratory manual*, 2nd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.
59. Schreiber, S. L., and G. R. Crabtree. 1992. The mechanism of action of cyclosporin A and FK506. *Immunol. Today* **13**:136–142.
60. Sherman, F., J. B. Hicks, and G. R. Fink. 1986. *Methods in yeast genetics*. Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.
61. Sikorski, R. S., and P. Hieter. 1989. A system of shuttle vectors and yeast host strains designed for efficient manipulation of DNA in *Saccharomyces cerevisiae*. *Genetics* **122**:19–27.
62. Tanida, I., A. Hasegawa, H. Iida, Y. Ohya, and Y. Anraku. 1995. Cooperation of calcineurin and vacuolar H<sup>+</sup>-ATPase in intracellular Ca<sup>2+</sup> homeostasis of yeast cells. *J. Biol. Chem.* **270**:10113–10119.
63. Tong, G., D. Shepherd, and C. E. Jahr. 1995. Synaptic desensitization of NMDA receptors by calcineurin. *Science* **267**:1510–1512.
64. Tropschug, M., I. B. Barthelmess, and W. Neupert. 1989. Sensitivity to cyclosporin A is mediated by cyclophilin in *Neurospora crassa* and *Saccharomyces cerevisiae*. *Nature (London)* **342**:953–955.
65. Wallis, J. W., G. Chrebet, G. Brodsky, M. Rolfe, and R. Rothstein. 1989. A hyper-recombination mutation in *S. cerevisiae* identifies a novel eukaryotic topoisomerase. *Cell* **58**:409–419.
66. Weisman, L. S., R. Bacallao, and W. Wickner. 1987. Multiple methods of visualizing the yeast vacuole permit evaluation of its morphology and inheritance during the cell cycle. *J. Cell Biol.* **105**:1539–1547.
67. Wurst, H., T. Shiba, and A. Kornberg. 1995. The gene for a major exopolyphosphatase of *Saccharomyces cerevisiae*. *J. Bacteriol.* **177**:898–906.
68. Ye, R. R., and A. Bretscher. 1992. Identification and molecular characterization of the calmodulin-binding subunit gene (*CMP1*) of protein phosphatase 2B from *Saccharomyces cerevisiae*. An alpha-factor inducible gene. *Eur. J. Biochem.* **204**:713–723.