

Glycogen Synthase Kinase 3 Is an Insulin-Regulated C/EBP α Kinase

SARAH E. ROSS, ROBIN L. ERICKSON, NAHID HEMATI, AND ORMOND A. MACDOUGALD*

Department of Physiology, University of Michigan Medical School, Ann Arbor, MI 48109-0622

Received 8 July 1999/Returned for modification 25 August 1999/Accepted 8 September 1999

CCAAT/enhancer binding protein α (C/EBP α) is a transcription factor involved in creating and maintaining the adipocyte phenotype. We have shown previously that insulin stimulates dephosphorylation of C/EBP α in 3T3-L1 adipocytes. Studies to identify the insulin-sensitive sites of phosphorylation reveal that a C/EBP α peptide (amino acids H215 to K250) is phosphorylated on T222, T226, and S230 *in vivo*. The context of these phosphoamino acids implicates glycogen synthase kinase 3 (GSK3), whose activity is known to be repressed in response to insulin, as a potential kinase for phosphorylation of T222 and T226. Accordingly, GSK3 phosphorylates the predicted region of C/EBP α on threonine *in vitro*, and GSK3 uses C/EBP α as a substrate *in vivo*. In addition, the effect of pharmacological agents on GSK3 activity correlates with regulation of C/EBP α phosphorylation. Treatment of 3T3-L1 adipocytes with the phosphatidylinositol 3-kinase inhibitor wortmannin results in phosphorylation of C/EBP α , whereas treatment with the GSK3 inhibitor lithium results in dephosphorylation of C/EBP α . Collectively, these data indicate that insulin stimulates dephosphorylation of C/EBP α on T222 and T226 through inactivation of GSK3. Since dephosphorylation of C/EBP α in response to lithium is blocked by okadaic acid, strong candidates for the T222 and T226 phosphatases are protein phosphatases 1 and 2a. Treatment of adipocytes with insulin alters the protease accessibility of widespread sites within the N terminus of C/EBP α , consistent with phosphorylation causing profound conformational changes. Finally, phosphorylation of C/EBP α and other substrates by GSK3 may be required for adipogenesis, since treatment of differentiating preadipocytes with lithium inhibits their conversion to adipocytes.

In view of the prevalence of obesity and obesity-related diseases, such as type II diabetes, it is important to understand the molecular basis for adipose tissue development and metabolism. Traditionally, adipocytes were known to play an important role in lipid homeostasis through their ability to store triacylglycerol and release free fatty acids and glycerol in response to changing energy needs. It is now recognized that adipocytes play a more central role in metabolism through their secretion of factors that regulate food intake and metabolic efficiency (46). The negative health consequences of excess white adipose tissue have been well documented (49), but the effects of an absence of fat have only been studied recently. Mice without white adipose tissue were created by directing expression to adipocytes of a dominant-negative protein that forms dimers with members of the C/EBP and AP-1 families, but which does not bind DNA (37). These mice have abnormal growth rates and enlarged internal organs and die prematurely. In addition, they exhibit a number of metabolic defects, including diabetes, with reduced leptin and increased serum glucose, insulin, free fatty acids, and triacylglycerols. The widespread effects of blocking C/EBP and AP-1 activities in adipose tissue highlight the importance of studying these transcription factors in adipocyte differentiation and metabolism.

The molecular events associated with preadipocyte differentiation have been most thoroughly studied in 3T3-L1 cells (reviewed in references 9, 13, 32, 34, and 46). Treatment of preadipocytes with inducers of differentiation stimulates a rapid and transient increase in C/EBP β and C/EBP δ , which in turn mediates the transcriptional activation of peroxisomal proliferator-activated receptor γ (PPAR γ) during the 2 days following stimulation of differentiation. Expression of PPAR γ ,

probably in conjunction with C/EBP β , induces expression of C/EBP α to its maximal level 4 to 5 days after initiation of differentiation. Together, C/EBP α and PPAR γ activate the transcription of genes involved in creating and maintaining the adipocyte phenotype (e.g., adipocyte lipid binding protein [422/aP2], stearoyl coenzyme A desaturase I, insulin-responsive glucose transporter [GLUT4], and leptin) (6, 24, 25, 31). The expression of PPAR γ and C/EBP α remains elevated in adipocytes through positive self- and cross-regulation.

C/EBP α not only plays an important role in preadipocyte differentiation, but also regulates gene expression in fully differentiated adipocytes. Although C/EBP α -deficient fibroblasts acquire the morphological characteristics of adipocytes upon ectopic expression and activation of PPAR γ , these C/EBP α -/- adipocytes are resistant to insulin. Further analyses revealed that these cells fail to increase glucose uptake in response to insulin because of impaired expression of insulin receptor and insulin receptor substrate 1 (50). Likewise, NIH-3T3 cells, which differentiate into adipocytes without expression of C/EBP α , are insulin resistant. In this case, it was found that the cells do not acquire the ability to transport glucose in response to insulin because of impaired expression of GLUT4 (17). Thus, C/EBP α is essential for the acquisition of insulin sensitivity by adipocytes. Given the requirement of C/EBP α for insulin responsiveness, it is not surprising that insulin feeds back to inhibit this transcription factor. In 3T3-L1 adipocytes, insulin suppresses transcription of the C/EBP α gene and stimulates dephosphorylation of C/EBP α protein (22, 31). These events closely correlate with suppression of GLUT4 gene expression, perhaps as part of the mechanism of desensitization to insulin. In this study, we investigate further the dephosphorylation of C/EBP α by insulin.

C/EBP α is a member of the basic region-leucine zipper (bZIP) family of transcription factors (reviewed in reference 35). The C-terminal zipper domain mediates the formation of homodimers as well as heterodimers with family members and perhaps a subset of other bZIP transcription factors (e.g.,

* Corresponding author. Mailing address: Department of Physiology, University of Michigan Medical School, 1301 E. Catherine Rd., Ann Arbor, MI 48109-0622. Phone: (734) 647-4880. Fax: (734) 936-8813. E-mail: macdouga@umich.edu.

TABLE 1. Oligonucleotides synthesized and then subcloned into newly created *PstI* and *SphI* sites of C/EBP α

Oligonucleotide	Sequence
Wild type; TTS	GCCTGGCCACCCACACCGCCGCCACGCCGTGCCAGCCGCATG
Mutant	
AAA	GCCTGGCCACCCCGCACCGCCGCCGCCGCCGTGCCGCCACATG
SSS	GCCTGGCCACCCCGCCCGCCGCCAGCCCGTGCCAGCCACATG
STS	GCCTGGTCACCCAGCCCGCCGCCAGCCCGTGCCAGCCACATG
ATS	GCCTGGTCACCCCGCCCGCCGCCAGCCCGTGCCAGCCGCATG
AAS	GCCTGGCCACCCCGCACCGCCGCCGCCGCCGTGCCAGCCACATG

ATF2) (44). The basic region is adjacent to the zipper domain and binds specific DNA sequences. The N terminus of C/EBP α contains multiple transactivation domains that work synergistically to transactivate C/EBP α -dependent promoters. The region N terminal to bZIP contains four regions that are highly conserved throughout evolution and that are separated by linker regions enriched in glycines and prolines (18). Conserved regions 1, 2, and 3 loosely correspond to transactivation domains identified previously (19, 39–41). Each conserved region is capable of transactivating the leptin promoter when fused to the C/EBP α bZIP domain (18). Conserved region 4 is the most highly conserved region outside the bZIP and probably plays a regulatory role in C/EBP α function. In the present study, three sites of phosphorylation were identified within this region, T222, T226, and S230, two of which are regulated by insulin. We present evidence that glycogen synthase kinase 3 (GSK3) phosphorylates T222 and T226, causing a conformational change in C/EBP α . Upon treatment with insulin, T222 and T226 are dephosphorylated through inactivation of GSK3 and activation of protein phosphatase 1 (PP1) or PP2A. Finally, we present evidence that GSK3 is required for adipocyte development, since inhibition of GSK3 activity with lithium blocks, not only C/EBP α phosphorylation, but also preadipocyte differentiation.

MATERIALS AND METHODS

Cell culture and transfection. Mouse 3T3-L1 preadipocytes (20) and human embryonic kidney 293T cells (a kind gift from Mitchell Lazar) were maintained in Dulbecco's modified Eagle's medium (DMEM; Gibco-BRL) supplemented with 10% calf serum, as described previously (22). Cells were transfected by calcium phosphate coprecipitation, as described previously (24). Expression plasmids, at the amounts indicated, were supplemented with pcDNA3.1(-) (Invitrogen), such that the total DNA was 20 μ g/10-cm-diameter plate. Cells were lysed 24 to 48 h following transfection. Constitutively active GSK3 (S9A) was obtained from Peter J. Roach (Indiana University School of Medicine) (16).

C/EBP α expression plasmids. To increase the ease of manipulating the mouse C/EBP α gene (5), four unique restriction sites were created through introduction of silent mutations. C/EBP α from the *NruI* site (+5 nucleotide [nt]) to the *EcoRV* site (+2111 nt) was subcloned into pBluescript (KS+) [pBS(KS+)] (Stratagene), and the following silent mutations were made by site-directed mutagenesis (QuickChange; Stratagene): a *KpnI* site was generated by mutation of C531 to G, a *PstI* site was generated by mutation of A774 to G, an *SphI* site was generated by mutation of C822 to T, and an *XhoI* site was generated by mutation of G1074 to C. In addition, the translational start site for C/EBP α was optimized, as described previously (29), to maximize expression of full-length p42C/EBP α , while minimizing translation from internal methionines. A p30C/EBP α expression vector was constructed by excising the first 180 nt of the gene, which includes the start sites of translation for p42C/EBP α and p40C/EBP α , but not the start site for p30C/EBP α .

The C/EBP α gene was also engineered to encode proteins in which T222, T226, and S230 have been mutated to either alanines or serines. To create these mutations, the TTS (wild type), AAA, SSS, STS, ATS, and AAS oligonucleotides were synthesized and then subcloned into the newly created *PstI* and *SphI* sites of C/EBP α (Table 1). These C/EBP α mutants were subcloned from pBS(KS+) into pcDNA 3.1(+) (Invitrogen) by using *BamHI* and *HindIII* sites, and the resultant expression vectors were used in transient transfections.

To make His-tagged, full-length C/EBP α (His-p42C/EBP α), the *NruI-HindIII* fragment (containing C/EBP α +5 nt to +2079 nt) was excised from C/EBP α , and the *HindIII* site was filled in with Klenow fragment prior to subcloning into the

PvuII site of pEBVHisA (Invitrogen). Similarly, His-tagged p18C/EBP α (His-p18C/EBP α ; also referred to as His-p18TTS), His-tagged p12C/EBP α (His-p12C/EBP α), and His-tagged p10C/EBP α (His-p10C/EBP α) were constructed by inserting the *MluI-EcoRV* fragment (+698 nt to +2111 nt) into the *PvuII* site of pEBVHisA, the *SmaI-SmaI* fragment (+861 nt to +1293 nt) into the *XhoI* site of pEBVHisC, and the *BanI-SmaI* fragment (+931 nt to +1293 nt) into the *PvuII* site of pEBVHisB, respectively. His-p18SSS and His-p18AAA were constructed with the p42C/EBP α SSS and p42C/EBP α AAA mutants, respectively. Briefly, after the *MluI* site was filled in, the *MluI-HindIII* fragment was subcloned into the *EcoRV* and *HindIII* sites of pcDNA3.1(-). This fragment was then excised by using *PmeI* and *HindIII* and subcloned into the *PvuII* and *HindIII* sites of pEBVHisA. His-p18STS, His-p18ATS, and His-p18AAS were created as described above for His-p18TTS with the primers indicated.

In vivo phosphorylation. 293T cell monolayers were preincubated in phosphate-free DMEM (Gibco-BRL) for 30 min. These cells were then incubated in phosphate-free DMEM that was supplemented with 32 P_i (Amersham; 0.5 mCi/ml; 4 ml/10-cm-diameter plate) for 3 h. After being rinsed twice with phosphate-buffered saline, the cells were lysed in urea lysis buffer (8 M urea, 0.5 M NaCl, 45 mM Na₂HPO₄, 5 mM NaH₂PO₄, 10 mM imidazole [pH 8.0]) for protein purification.

Purification of His-tagged proteins. To identify sites of phosphorylation in C/EBP α , 100 10-cm-diameter plates of 293T cells were transiently transfected with expression plasmids encoding either His-p42C/EBP α or His-p18C/EBP α . These plates were rinsed with phosphate-buffered saline, and then each was lysed in 0.5 ml of urea lysis buffer and sonicated. Pooled lysates were incubated with 2 ml of ProBond nickel resin (Invitrogen) for 2 h at room temperature. Nickel resin was washed three times with 10 bed volumes of wash buffer (8 M urea, 0.4 M NaCl, 17.6 mM Na₂HPO₄, 32.4 mM NaH₂PO₄, 10 mM imidazole [pH 6.75]), and bound proteins were eluted by being washed two times in 5 bed volumes of elution buffer (8 M urea, 0.4 M NaCl, 6 mM Na₂HPO₄, 44 mM NaH₂HPO₄, 10 mM imidazole [pH 5.30]). The solution containing these eluted proteins was adjusted to pH 8.0, and the proteins were incubated a second time with nickel resin. This purification process was repeated twice with the following changes: for each of the subsequent rounds, the bed volume was decreased by half, and for the final wash, the stringency was increased by raising the concentration of imidazole from 10 to 50 mM. This protocol was also used in a scaled-down version to purify His-C/EBP α from 12 plates of transfected 293T cells labeled with 32 P_i. Unlabeled and labeled products from this purification procedure were pooled and concentrated by centrifugation (Centricon-10; Amicon). Purification products were then separated by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE). To optimize protein purification and monitor protein purity, proteins were visualized by silver staining (2), and C/EBP α was detected by immunoblot analysis. For protein recovery, SDS-PAGE gels were zinc stained according to the manufacturer's instructions (Pierce). His-C/EBP α was identified by its mobility and excised from the gel. Samples were then sent to the Protein Structure Core Facility at the University of Nebraska Medical Center for analyses. In-gel cleavage was performed by using vapor cyanogen bromide followed by trypsin. Peptides were separated on a Vydac C₁₈ column and sequenced with an ABI 477 protein sequencer, as described previously (48).

Immunoblot analysis. Cell lysis and immunoblotting for C/EBP α were performed as described previously (22) with a polyclonal C/EBP α antibody that was raised against a synthetic peptide (amino acids 253 to 265) (29).

Phosphoamino acid analysis. Purified His-tagged forms of truncated or full-length C/EBP α that had been phosphorylated either in vitro or in vivo were separated by SDS-PAGE. After transfer onto PVDF (polyvinylidene difluoride)-Plus membrane (Micron Separations, Inc.), and visualization by autoradiography, His-tagged C/EBP α was excised and hydrolyzed in 6 M HCl for 60 min at 110°C. Acid was removed by evaporation in a Speedvac and with two 0.5-ml washes of water. Samples were resuspended in 10 μ l of water containing 1 ng of phosphoserine, phosphotyrosine, and phosphothreonine (Sigma) and were spotted onto cellulose plates (Kodak). Amino acids were separated by thin-layer electrophoresis for 45 min at 20 mA in pH 2.5 buffer (5.9% [vol/vol] glacial acetic acid, 0.8% [vol/vol] formic acid [88%], 0.3% [vol/vol] pyridine, 0.3 mM EDTA) (21). Phosphoamino acid standards were visualized by ninhydrin staining, and 32 P-labeled phosphoamino acids were detected by autoradiography.

In vitro kinase reactions. Protein purification for in vitro kinase assays of His-p42C/EBP α , His-p18C/EBP α , His-p12C/EBP α , and His-p10C/EBP α was performed as described above, except that His-tagged forms of C/EBP α were not eluted from nickel resin following the last purification. Rather, purified C/EBP α proteins were washed with kinase buffer and then incubated in kinase buffer with or without 5 U of GSK3 (New England Biolabs) for 60 min at 37°C. The nickel resin was washed extensively with wash buffer, and proteins were eluted in elution buffer and separated by SDS-PAGE as described above. Proteins were transferred to PVDF-Plus membrane (22), and the blot was subjected to autoradiography and immunoblot analyses.

Preparation of nuclei. Nuclei were purified from 3T3-L1 preadipocytes or adipocytes by a procedure modified from that of Dignam et al. (15). Briefly, 3T3-L1 cells were washed with 5 ml of phosphate-buffered saline, and then 2 ml of hypotonic lysis buffer (20 mM Tris [pH 7.5], 10 mM NaCl, 3 mM MgCl₂, 1 mM dithiothreitol [DTT]) with 2- μ l/ml protease inhibitors (PIC 1, which is 1-mg/ml leupeptin, 1-mg/ml antipain, and 10-mg/ml benzamide in aprotinin; and PIC 2, which is 1-mg/ml chymostatin and 1-mg/ml pepstatin A in dimethyl sulfoxide [DMSO]), and phosphatase inhibitors (30 mM β -glycerophosphate, 1 mM sodium orthovanadate, and 1-mg/ml *p*-nitrophenylphosphate). Cells were scraped in 1 ml of hypotonic lysis buffer, and Igepal CA-630 was added (1:100 [vol/vol] for preadipocytes, 1:67 [vol/vol] for adipocytes) prior to Dounce homogenization (25 strokes). Disruption of plasma membranes was verified by trypan blue staining and light microscopy. Nuclei were pelleted in a microcentrifuge at 5,000 \times *g* for 1 min. The supernatant was removed, the nuclei-containing pellet was resuspended in 1 ml of hypotonic lysis buffer, and the sample was centrifuged as described above. The process of resuspension and centrifugation was repeated. For quantification of nuclei, 1.5- μ l aliquots were lysed in 0.5% SDS and vortexed, and *A*₂₆₀ was measured. Nuclear pellets were lysed in isoelectric focusing buffer or immunoblot lysis buffer or used for the protease accessibility assay.

Protease accessibility assay. To assess the sensitivity of nuclear C/EBP α to protease, nuclei from 3T3-L1 adipocytes were resuspended in hypotonic lysis buffer at 60 *A*₂₆₀/ml. Increasing amounts of trypsin in 20 μ l of phosphate-buffered saline were added to 30- μ l aliquots of nuclei on ice for 30 min. The final concentrations of trypsin were 0, 8, 24, 80, and 240 μ g/ml. Reactions were terminated with 50 μ l of 2 \times immunoblot lysis buffer (2% SDS and 120 mM Tris [pH 6.8]) and heated to 100°C for 10 min. Samples were subjected to SDS-PAGE and immunoblot analysis for C/EBP α .

Alkaline phosphatase treatment of samples. Nuclei were lysed in an SDS-containing buffer (1% SDS, 60 mM Tris [pH 6.8]) at a concentration of approximately 20 mg of protein/ml. Nuclear proteins were incubated with alkaline phosphatase (Boehringer Mannheim) at a concentration of approximately 1 U/mg of protein in the SDS-containing buffer for 1 h at 37°C. Samples were mixed with an equal volume of isoelectric focusing buffer prior to electrophoretic separation.

Isoelectric focusing. Nuclei were lysed in isoelectric focusing buffer (9 M urea, 1% Igepal CA-630, 1% DTT) at a concentration of approximately 10 mg of protein/ml. This nuclear lysate was sonicated and then centrifuged at 1000 *g* for 15 min, and the supernatant was collected for further analysis. Polyacrylamide-urea mini gels were made according to the manufacturer's instructions (Bio-Rad) with a 1:1 mixture of pH 3 to 10 and pH 5 to 8 ampholytes (Bio-Rad). Protein samples (approximately 20 μ g) were loaded and focused for 15 min at 100 V, 15 min at 200 V, and then 1 h at 450 V. Proteins were then transferred onto PVDF-plus membrane for immunoblot analysis.

Preadipocyte differentiation and Oil Red-O staining. 3T3-L1 preadipocytes were induced to differentiate as described previously (47), except that the medium was not supplemented with insulin on day 4 or thereafter (30). Oil Red-O staining was performed essentially as described previously (42). Briefly, cell monolayers were washed with phosphate-buffered saline, fixed in 3.7% formaldehyde for 2 min, washed with H₂O, incubated with Oil Red-O solution for 1 h at room temperature, and then washed with H₂O.

RESULTS

Identification of T222, T226, and S230 as phosphoamino acids. We have shown previously that C/EBP α contains amino acids that become dephosphorylated in response to insulin (22, 31). To determine which amino acids in C/EBP α are phosphorylated, we labeled the cellular phosphate pool and purified ³²P-labeled C/EBP α for analyses. In these experiments, 293T cells were transiently transfected with expression vectors encoding His-tagged C/EBP α constructs. Transfected cells were incubated with ³²P_i, and His-tagged forms of C/EBP α were purified on a nickel resin. Eluted proteins were separated by SDS-PAGE, and C/EBP α was excised from the gel. C/EBP α was subjected to in-gel cleavage by treatment with cyanogen bromide and trypsin. The C/EBP α peptides were separated by high-performance liquid chromatography. Following this separation, labeled peptides were sequenced by Edman degrada-




FIG. 1. Identification of phosphorylation sites in C/EBP α . (A) A schematic diagram of C/EBP α . Regions that are highly conserved across species (conserved regions 1 to 4 and bZIP domain) are shaded. Bent arrows denote translational start sites for two predominant C/EBP α species (p42C/EBP α and p30C/EBP α). A phosphopeptide identified in three independent labeling experiments is indicated by the solid bar. Amino acids within a region of this phosphopeptide are shown for mouse, human, chicken, *Xenopus*, and *Aplysia* proteins. The consensus sequence for GSK3 is given. (B) Schematic representation of the His-tagged C/EBP α fusion proteins used in subsequent labeling experiments. The C/EBP α amino acid sequence from T222 to P231 is given, and the C/EBP α mutants in which T222, T226, and S230 were mutated to either alanines or serines are illustrated. (C) His-p18TTS (lane 1), His-p18AAA (lane 2), His-p18SSS (lane 3), His-p18STS (lane 4), His-p18ATS (lane 5), and His-p18AAS (lane 6) were expressed in 293T cells, labeled with ³²P in vivo, purified, and separated by SDS-PAGE. These samples were subjected to immunoblot analysis for C/EBP α (top panel; IB) and autoradiography (middle panel; ³²P). Phosphoamino acid analysis was performed with the remainder of the His-tagged C/EBP α mutants (bottom panels). The positions of phosphoserine (pS), phosphothreonine (pT), and phosphotyrosine (pY) are indicated. His-p18AAA contained no detectable ³²P.

tion to identify the phosphoamino acids. Using this approach, in three independent experiments, the peptide spanning amino acids H215 through K250 was identified as a phosphopeptide (solid bar in Fig. 1A). Amino acid analysis of this phosphopeptide revealed that it contains both phosphoserine and phosphothreonine. Edman degradation demonstrated that this peptide is phosphorylated on T222 and S230.

The identified amino acids are within a consensus sequence of phosphorylation by GSK3 (Fig. 1A) (26). In addition, since insulin inhibits GSK3 activity in adipocytes (11), GSK3 is a




FIG. 2. GSK3 phosphorylates C/EBP α in vitro. (A) Schematic representation of His-tagged C/EBP α fusion proteins used as substrates in kinase assays in vitro. A putative region of GSK3 phosphorylation is denoted by the solid bar. (B) His-tagged p42, p18, p12, and p10 C/EBP α were expressed in 293T cells and purified. These proteins were mixed together, and kinase reactions were performed in the absence (–) or presence (+) of recombinant GSK3. Proteins were separated by SDS-PAGE (15% polyacrylamide gel) and subjected to immunoblot analysis with antibody specific for C/EBP α . The migration of His-tagged p42, p18, p12, and p10 C/EBP α is indicated. (C) The immunoblot in panel B was subjected to autoradiography. (D) 32 P-labeled His-p42C/EBP α was excised, and phosphoamino acid analysis was performed. Phosphoamino acid standards (pS, pT, and pY) are indicated. Similar results were obtained in three independent experiments.

particularly good candidate for modulating sites that are dephosphorylated upon insulin treatment. We therefore considered the possibility that GSK3 is a C/EBP α kinase. For many GSK3 substrates, phosphorylation of a priming site at the *n* position is required for subsequent phosphorylation by GSK3 at the *n* – 4 position (43). Applying this hierarchical model to C/EBP α , we predicted that phosphorylation of S230 by an unspecified kinase would be necessary for phosphorylation at T226 by GSK3. Thereafter, phosphorylation of T226 would serve as the priming site for GSK3-mediated phosphorylation of T222. We identified phosphate on both T222 and S230, but we were unable to conclusively identify T226 as a phosphoamino acid by using Edman degradation. 32 P at this site may be difficult to detect because slower phosphate turnover at T226 relative to T222 and S230 could result in lower specific activity upon metabolic labeling, or because more rapid dephosphorylation of T226 than of the other sites could result in lower relative stoichiometry of phosphorylation.

To determine whether C/EBP α is phosphorylated on T226, we designed a series of truncated C/EBP α mutants for use in phosphoamino acid analysis experiments. Specifically, we created a His-tagged expression vector for an N-terminally truncated form of C/EBP α , His-p18TTS, and ones in which T222, T226, and S230 were converted to combinations of alanines and serines (His-p18SSS, His-p18AAA, His-p18STS, His-p18ATS, and His-p18AAS), as shown schematically in Fig. 1B. 293T cells were transiently transfected with expression vectors for His-p18C/EBP α proteins. Two days later, cells were labeled with 32 P_i, and His-tagged C/EBP α proteins were purified on a nickel resin. Immunoblot analysis of the purified products revealed that the C/EBP α proteins were expressed and purified (Fig. 1C, top panel). Autoradiography of these immunoblots revealed that each of these C/EBP α proteins, except His-p18AAA, is phosphorylated in vivo (Fig. 1C, middle panel). His-p18AAA (lane 2) contains no detectable phosphate, indicating that p18TTS is not phosphorylated on any sites other

than T222, T226, and S230. Upon phosphoamino acid analysis, we found that His-p18TTS (lane 1), His-p18STS (lane 4), and His-p18ATS (lane 5) are phosphorylated on serine and threonine, whereas His-p18SSS (lane 3) and His-p18AAS (lane 6) are phosphorylated on serine alone (Fig. 1C, bottom panel). Loss of phosphothreonine upon conversion of either His-p18STS to His-p18SSS or His-p18ATS to His-p18AAS indicates that T226 is phosphorylated in vivo. This finding, together with results from our initial mapping studies (above), demonstrates that C/EBP α is phosphorylated on T222, T226, and S230.

GSK3 phosphorylates C/EBP α in vitro. To investigate further whether GSK3 phosphorylates C/EBP α , we used purified C/EBP α as a substrate for GSK3 in vitro. For these assays, His-tagged versions of full-length or truncated forms of C/EBP α (Fig. 2A) were purified from 293T cells. Wild-type p42C/EBP α and p18C/EBP α contain the GSK3 consensus sites of phosphorylation, whereas p12C/EBP α and p10C/EBP α do not. Purified His-tagged forms of C/EBP α were combined, and duplicate kinase assays were performed in either the absence or presence of GSK3 prior to separation by SDS-PAGE. Immunoblot analysis revealed that the amounts of C/EBP α proteins in each reaction varied by less than a factor of 2 (Fig. 2B). Autoradiography revealed that p42C/EBP α and p18C/EBP α , but not p12C/EBP α and p10C/EBP α , are substrates for phosphorylation by GSK3 (Fig. 2C). Phosphoamino acid analysis demonstrated that phosphorylation of p42C/EBP α by GSK3 in vitro occurs only on threonine (Fig. 2D). These findings indicate that GSK3 phosphorylates threonine specifically in a region between amino acids R192 and G246 in vitro and are consistent with GSK3 phosphorylating T222 and T226.

GSK3 phosphorylates C/EBP α in vivo. Since the steady-state stoichiometry of protein phosphorylation depends upon the balance of kinase and phosphatase activities, we reasoned that overexpression of a C/EBP α kinase would increase the proportion of C/EBP α that is phosphorylated. To assess whether GSK3 is a C/EBP α kinase in a cellular context, 293T cells were transfected with expression vectors encoding either p30C/EBP α alone (Fig. 3, lanes 1 to 4) or p30C/EBP α and constitutively active GSK3 (Fig. 3, lanes 5 to 8). p30C/EBP α is an alternate translation product which arises from initiation at the third start codon (29). This form was used because phosphorylation-induced shifts in p30C/EBP α can be visualized by SDS-PAGE (22). Two days after transfection, cells were not treated or treated with the GSK3 inhibitor lithium for 1 h prior to lysis. Samples were separated by SDS-PAGE, and C/EBP α was detected by immunoblot analysis. When expressed alone, p30C/EBP α exists predominantly as a single high-mobility form (Fig. 3, lanes 1 and 3). Inhibition of GSK3 activity by treatment of these cells with lithium results in the loss of the weak band that crowns the p30C/EBP α species (Fig. 3, lanes 2 and 4). This observation suggests that, under these conditions, a small proportion of p30C/EBP α is phosphorylated due to endogenous GSK3 activity in 293T cells. Overexpression of constitutively




FIG. 3. p30C/EBP α is a substrate for GSK3 in vivo. 293T cells were transfected with p30C/EBP α alone or both p30C/EBP α and constitutively active GSK3. Forty-eight hours later, these cells were not treated (–) or treated with 25 mM LiCl (+) for 1 h prior to lysis. Samples were separated by SDS-PAGE (11.5% polyacrylamide gel) and subjected to immunoblot analyses for C/EBP α . Similar results were obtained in three independent experiments.




FIG. 4. Regulation of C/EBP α phosphorylation by pharmacological agents. (A) 3T3-L1 adipocytes were not treated (Con; lanes 1 and 2) or were treated with 167 nM insulin for 60 min (Ins; lanes 3 and 4), 200 nM wortmannin for 60 min (Wort; lanes 5 and 6), or 25 mM LiCl for 30 min (Li; lanes 7 and 8). Whole-cell lysates were separated by SDS-PAGE (11.5% polyacrylamide gel) and subjected to immunoblot analysis for C/EBP α . The migration of p42C/EBP α and p30C/EBP α is indicated. p30C/EBP α migrates as three bands (a to c). (B) 3T3-L1 adipocytes were treated as in panel A, except that nuclei were prepared. Nuclear proteins were separated by isoelectric focusing and subjected to immunoblot analysis for C/EBP α . Acidic (+) and basic (-) ends of the blot are indicated. Similar results were obtained in three independent experiments. Since p30C/EBP α is more basic (predicted pI of 9.7) than p42C/EBP α (predicted pI of 7.6), p30C/EBP α is not resolved into discrete bands when focused to equilibrium with standard ampholytes (pH range of 3 to 10). p42C/EBP α can be resolved into five bands (a to e) by this technique. (C) 3T3-L1 adipocytes were not treated (Con; lane 1) or were treated with 25 mM LiCl (Li; lane 2), 1.5 μ M okadaic acid (OA; lane 3), or both 25 mM LiCl and 1.5 μ M okadaic acid (lane 4) for 1 h. Whole-cell lysates were separated by SDS-PAGE (11.5% polyacrylamide gel) and subjected to immunoblot analysis for C/EBP α . (D) Nuclear lysates were prepared from 3T3-L1 adipocytes and either not treated (Con; lane 1) or treated with alkaline phosphatase (AP; lane 2) prior to isoelectric focusing and immunoblot analysis for C/EBP α . (E) Model of signaling pathway through which insulin, wortmannin, lithium, and okadaic acid regulate the phosphorylation of C/EBP α .

active GSK3 dramatically increases the proportion of the upper species (Fig. 3, lanes 5 and 7). Inhibition of GSK3 activity by 1 h of lithium treatment results in the loss of this upper species. These data suggest that GSK3 phosphorylates C/EBP α in vivo.

Phosphorylation of C/EBP α correlates with the regulation of GSK3 activity by insulin, wortmannin, and lithium. We have demonstrated that GSK3 phosphorylates C/EBP α in vitro and in vivo. We next tested whether the ability of insulin to inhibit GSK3 (38) is responsible for insulin-dependent dephosphorylation of C/EBP α in 3T3-L1 adipocytes. In many cell types, including adipocytes, activation of the insulin receptor is known to stimulate phosphatidylinositol (PI) 3-kinase activity, which, through the actions of other signaling molecules (possibly Akt [10] or PDK1 [14]), causes phosphorylation and inhibition of GSK3 (11). If GSK3 is the insulin-sensitive C/EBP α kinase, then we would expect inhibition of PI 3-kinase with wortmannin to stimulate GSK3 and therefore cause phosphorylation of C/EBP α . Conversely, we would expect inhibition of GSK3 activity with lithium to result in dephosphorylation of C/EBP α .

In these experiments, we treated 3T3-L1 adipocytes with wortmannin or lithium and then assessed the phosphorylation of the two predominant C/EBP α translation products, p30C/EBP α and p42C/EBP α , by immunoblot analysis. In the absence of treatment (Fig. 4A, lanes 1 and 2), p30C/EBP α has three distinct mobilities upon SDS-PAGE (bands a to c). We have shown previously that these differences in mobility are due to phosphorylation. Treatment of samples with alkaline phosphatase in vitro causes the top two bands to disappear with a proportionate increase in the bottom band (22). Treatment with either insulin (Fig. 4A, lanes 3 and 4) or lithium (Fig. 4A, lanes 7 and 8) causes dephosphorylation of p30C/

EBP α (loss of bands a and b with an increase in band c), whereas treatment with wortmannin (Fig. 4A, lanes 5 and 6) results in net phosphorylation of C/EBP α (loss of band c). Therefore, insulin appears to stimulate dephosphorylation of two sites within p30C/EBP α through inhibition of GSK3.

Treatment of adipocytes with insulin, wortmannin, or lithium appears to cause analogous changes in the phosphorylation of p42C/EBP α . However, phosphorylation-induced changes in the mobility of this larger protein are difficult to resolve by SDS-PAGE. Thus, to determine whether p42C/EBP α is regulated similarly to p30C/EBP α , a parallel experiment was performed in which 3T3-L1 adipocyte lysates were analyzed by isoelectric focusing (Fig. 4B). p42C/EBP α from unstimulated 3T3-L1 adipocytes is resolved into at least five bands by this technique (Fig. 4B, lanes 1 and 2 [forms a to e]). Treatment of adipocytes with either insulin (Fig. 4B, lanes 3 and 4) or lithium (Fig. 4B, lanes 7 and 8) results in a decrease in the two most acidic forms of p42C/EBP α (bands a and b). In contrast, treatment of cells with wortmannin (Fig. 4B, lanes 5 and 6) causes loss of the most basic form of p42C/EBP α (band e). To confirm that the presence of multiple p42C/EBP α bands arises from the phosphorylation of p42C/EBP α , adipocyte lysates were treated with alkaline phosphatase in vitro prior to isoelectric focusing. Upon this treatment, the acidic forms of C/EBP α were lost and there was accumulation of its most basic form (Fig. 4D, lane 2). This observation indicates that the most basic species is C/EBP α without phosphate and suggests that the acidic bands represent C/EBP α with increasing amounts of phosphate. The simplest interpretation of these data is that p42C/EBP α , like p30C/EBP α , is dephosphorylated at two sites following insulin treatment. In addition, the regulation of p30C/EBP α and p42C/EBP α phosphorylation by wortmannin

and lithium is consistent with the hypothesis that GSK3 is the insulin-sensitive C/EBP α kinase.

Phosphorylation of C/EBP α correlates with the regulation of PP1 and PP2A by okadaic acid. Dephosphorylation of C/EBP α requires not only inhibition of GSK3, but also activity of a C/EBP α phosphatase. PP1 is a good candidate, since the activity of this phosphatase is induced by insulin in 3T3-L1 adipocytes, and since treatment of adipocytes with okadaic acid, which inhibits PP1 and PP2A, results in accumulation of hyperphosphorylated C/EBP α (31). To determine whether PP1 or PP2A is directly responsible for dephosphorylation of the sites within p30C/EBP α , 3T3-L1 adipocytes were not treated or treated with lithium, okadaic acid, or lithium and okadaic acid (Fig. 4C, lanes 1 to 4). Since lithium stimulates only minimal dephosphorylation in the presence of okadaic acid, PP1 or PP2A is likely responsible for dephosphorylation of C/EBP α at these sites. Moreover, since induction of PP1 activity by insulin is inhibited by wortmannin (3), our finding that wortmannin stimulates phosphorylation of C/EBP α in the presence of lithium is consistent with wortmannin also inhibiting C/EBP α phosphatase activity (not shown). Taken together, these data are consistent with the model proposed in Fig. 4E, in which stimulation of PI 3-kinase by insulin results in simultaneous inhibition of the C/EBP α kinase (GSK3) and activation of the C/EBP α phosphatase (PP1 or PP2A) to cause dephosphorylation of C/EBP α .

Phosphorylation alters C/EBP α conformation. To determine whether phosphorylation alters the three-dimensional structure of C/EBP α , we compared the accessibilities to protease of specific C/EBP α residues after treatment of adipocytes with insulin or wortmannin. The sensitivity to proteolysis depends on the tertiary structure of the protein and the extent to which the protease-sensitive sites are protected or exposed through protein-protein or protein-DNA interactions. For instance, interaction of C/EBP α with DNA protects the bZIP domain from tryptic digestion (data not shown and reference 45). Trypsin cleaves R-X and K-X and is predicted to cut C/EBP α at 36 sites, of which 24 are in the bZIP domain and are therefore resistant to proteolysis. Although partial digestion of the 10 sites outside the bZIP domain by trypsin likely gives a large number of different C/EBP α fragments, the anti-peptide C/EBP α antibody used in our experiments only recognizes those that contain amino acids 253 to 265, a region just N terminal to the bZIP domain. Thus, only C-terminal proteolytic fragments of C/EBP α are observed in this protease accessibility assay.

Nuclei were isolated from 3T3-L1 adipocytes that had been treated for 1 h with insulin to induce dephosphorylation or wortmannin to induce phosphorylation (Fig. 3) (22). Nuclei were incubated with increasing concentrations of trypsin on ice for 30 min, and then nuclear proteins were separated by SDS-PAGE for immunoblot analysis (Fig. 5). Endogenous C/EBP α proteins, p42C/EBP α and p30C/EBP α , were observed in undigested samples from insulin-treated (Fig. 5, lane 1) or wortmannin-treated (Fig. 5, lane 2) adipocytes. The effects of phosphorylation on mobility of C/EBP α are not as well resolved as in Fig. 3, because the proportion of acrylamide in the gel was higher. Treatment of nuclei with increasing concentrations of trypsin resulted in C/EBP α digestion products of decreasing size. Of interest, the trypsin fingerprints from insulin- and wortmannin-treated cells, at any given trypsin concentration, are substantially different. For example, dephosphorylated C/EBP α from insulin-treated adipocytes shows cleavage at three distinct sites in the N terminus (Fig. 5, bands a to c) that are not observed with phosphorylated C/EBP α from wortmannin-treated adipocytes. In addition, more widespread changes




FIG. 5. Insulin and wortmannin alter the trypsin sensitivity of C/EBP α . 3T3-L1 adipocytes were treated for 1 h with 167 nM insulin or 250 nM wortmannin, as indicated. Nuclei were purified from cells and incubated for 30 min on ice in the absence (lanes 1 and 2) or presence of various concentrations of trypsin (lanes 3 and 4, 8 μ g/ml; lanes 5 and 6, 24 μ g/ml; lanes 7 and 8, 80 μ g/ml; lanes 9 and 10, 240 μ g/ml). Samples were separated by SDS-PAGE (15% polyacrylamide gel) and subjected to immunoblot analysis for C/EBP α . Because the C/EBP α antibody used in these experiments was raised against an epitope located near the bZIP domain, which is protected from trypsinization, these partially digested C/EBP α fragments are N-terminal truncation products.

in sensitivity to protease were observed in two other regions (Fig. 5, complex d and e). Although some differences in band pattern of complex d and e can be explained through differential phosphorylation altering mobility of C/EBP α upon SDS-PAGE, the loss of bands a to c after wortmannin treatment suggests that phosphorylation of C/EBP α alters the sensitivity of these sites to trypsin. It is conceivable that altered cleavage of some sites is due to phosphorylation per se (i.e., by inhibiting the interactions between trypsin and C/EBP α). However, the magnitude and widespread nature of the differential sensitivity suggest that phosphorylation causes a change in the three-dimensional structure of C/EBP α , thereby altering the accessibility of protease to many sites of cleavage. Since the accessibility of C/EBP α to trypsin was assessed in nuclei, changes in protease accessibility could also reflect differential interactions with transcriptional coactivators or other nuclear proteins.

Lithium inhibits preadipocyte differentiation. Given the importance of C/EBP α in preadipocyte differentiation (13) and the established role of GSK3 in the development of tissues in other species (4), we considered the possibility that GSK3 activity is required for adipogenesis. To assess this putative role for GSK3, we investigated whether the GSK3 inhibitor lithium could block preadipocyte differentiation (Fig. 6A). 3T3-L1 preadipocytes were induced to differentiate under standard conditions, with isobutylmethylxanthine, dexamethasone, insulin, and fetal calf serum, in the presence of either 25 mM LiCl or 25 mM NaCl as a control. Thereafter, cells were continuously incubated in the presence of LiCl or NaCl over the course of differentiation. Eight days after the induction of differentiation, the degree of adipogenesis was assessed qualitatively by staining cellular lipid droplets with Oil Red-O. Upon induction, 3T3-L1 preadipocytes differentiated almost completely, irrespective of the presence of NaCl (Fig. 6A, top). In contrast, lithium treatment almost completely inhibited adipocyte differentiation (Fig. 6A, bottom). Expression of adipocyte markers C/EBP α and 422/aP2 was also inhibited by lithium (data not shown). In addition to GSK3, lithium is known to inhibit inositol monophosphatase (27) and may inhibit other enzymatic activities as well. Nevertheless, the observed inhibition of adipogenesis by lithium is consistent with the hypothesis that GSK3 activity is required for adipocyte differentiation.

To address potential targets of GSK3 activity in preadipo-




FIG. 6. (A) Lithium inhibits preadipocyte differentiation. 3T3-L1 preadipocytes were induced to differentiate as described in Methods and Materials, except that on days 1, 2, 4, and 6, 25 mM NaCl or LiCl was added to the medium. On day 8, cells were fixed and stained with Oil Red-O to visually assess the accumulation of lipid droplets. (B) Schematic representation of a generalized C/EBP transcription factor. The region of C/EBP α that is phosphorylated by GSK3 and putative GSK3 recognition sequences in C/EBP β and C/EBP δ are given.

cyte differentiation, we considered known regulators of adipogenesis. The current model of 3T3-L1 differentiation involves coordinated expression of four transcription factors, C/EBP β , C/EBP δ , PPAR γ , and C/EBP α (13). Our work shows that at least one of these, C/EBP α , is phosphorylated by GSK3. As a first step to determine whether the others are likewise regulated by GSK3, we inspected their sequences for putative GSK3 sites. C/EBP β and C/EBP δ , but not PPAR γ , contain sequence elements which resemble GSK3 recognition sequences (Fig. 6B). Next, we determined whether the phosphorylation of C/EBP β is sensitive to lithium. Consistent with the hypothesis that C/EBP family members are regulated by GSK3, lithium treatment of cells caused dephosphorylation of C/EBP β as assessed by immunoblot analysis (41a). For each of these C/EBPs, the putative sites of GSK3 phosphorylation are located just N terminal to the bZIP. In addition to this conservation among C/EBP family members, within C/EBP α , C/EBP β , and C/EBP δ , the region of putative GSK3 phosphorylation is highly conserved across species. Collectively, these data raise the possibility that GSK phosphorylation of C/EBPs may be required for preadipocyte differentiation.

DISCUSSION

Our analyses have led to the discovery that several sites within conserved region 4 of C/EBP α are phosphorylated in vivo (Fig. 1). Two sites, T222 and T226, are phosphorylated by GSK3, whereas S230 is phosphorylated by an unknown kinase. In addition, this work delineates a part of the signaling mechanism through which C/EBP α phosphorylation is regulated.

Previous work (38) has established that insulin, acting through signaling intermediates, including PI 3-kinase, causes inhibition of GSK3 leads to dephosphorylation of two sites, T222 and T226 (Fig. 4). Furthermore, since okadaic acid treatment of adipocytes blocks the lithium-induced dephosphorylation of C/EBP α , it is likely that PP1 or PP2A is responsible for dephosphorylation of the insulin-sensitive sites (Fig. 4D). Our work also shows that insulin and wortmannin alter the protease accessibility of C/EBP α , thereby implying that phosphorylation at T222 and T226 causes a conformational change in the structure of C/EBP α (Fig. 5). Finally, we report here that lithium inhibits preadipocyte differentiation (Fig. 6), suggesting that GSK3-mediated phosphorylation of C/EBP α and other transcription factors, such as C/EBP β , is required for adipogenesis.

A hierarchical model of phosphorylation has been described for several GSK3 substrates (7, 43). If this model holds true for C/EBP α , then phosphorylation of S230 is required for subsequent phosphorylation of T226 and then T222. Results from two experiments indicate that phosphorylation by GSK3 need not occur in this requisite order. First, a glutathione *S*-transferase-C/EBP α fusion protein purified from bacteria (and therefore not phosphorylated) is phosphorylated by GSK3 in vitro on threonine alone (not shown). Second, based on the hierarchical model, we would expect neither the p30TTA mutant nor the p30AAA mutant to be phosphorylated by GSK3. However, we observed that while p30AAA exists as a single dephosphorylated band upon SDS-PAGE, p30TTA migrates as a doublet whose low-mobility band is rapidly lost after lithium treatment (not shown). Together, these experiments suggest that T222 and T226 are substrates for GSK3 even when S230 is mutated to alanine. Despite this negative evidence, it remains possible that phosphorylation of S230 changes the affinity of C/EBP α for recognition by GSK3 and may therefore serve a modulatory role.

Phylogenetic analysis of C/EBP α has allowed us to define four highly conserved regions in the N-terminal transactivation domain, and we believe that the regions of greatest functional importance are likely to be found therein (18). Each of conserved regions 1, 2, and 3 has intrinsic transactivation ability, suggesting that these regions may directly interact with coactivators or with the basal transcriptional machinery. In contrast, the fourth conserved region, although the most highly conserved, has no known function. Our finding that GSK3 phosphorylates three sites within conserved region 4 suggests that C/EBP α activity may be regulated through this mechanism. However, the precise role of this modification has proved difficult to uncover. Mutation of T222 and T226 to alanines does not change the ability of C/EBP α to transactivate leptin or C/EBP α promoters in reporter gene assays (data not shown). Furthermore, altering the phosphorylation status of C/EBP α by cotransfection of GSK3 also does not influence the ability of C/EBP α to transactivate in these experiments (data not shown). Finally, ectopic expression of the T222A, T226A mutant, like its wild-type counterpart, is sufficient to induce spontaneous differentiation of preadipocytes (data not shown), demonstrating that phosphorylation of these sites is not required for activation of chromatin-embedded genes. Thus, GSK3-mediated phosphorylation does not, in itself, dramatically alter the activity of C/EBP α in our assays. Since the protease accessibility of C/EBP α was determined in nuclei, the effects of phosphorylation on sensitivity of C/EBP α to trypsin (Fig. 5) may be the result of conformational changes or differential interactions with nuclear proteins.

It has been reported that protein kinase C can phosphorylate specific sites within the basic region of C/EBP α in vitro

(33). This modification is an attractive mechanism for the regulation of C/EBP α activity, since addition of a negative charge to the basic region profoundly reduces its affinity for DNA. However, we have no evidence from labeling experiments that this phosphorylation occurs *in vivo*. Phosphate incorporation into His-p18AAA was undetectable in our studies, suggesting that T222, T226, and S230 are the only phosphorylated residues in the C-terminal 18 kDa of C/EBP α (Fig. 1C).

We have shown that phosphorylation of C/EBP α by GSK3 is regulated by insulin. In addition to its role in insulin signaling, GSK3 is a mediator in the Wnt signaling pathway, where it has been shown to be important in the specification of cell fate (reviewed in references 4 and 12). Wnts are a family of secreted glycoproteins which, probably through activation of frizzled receptors, stimulate a signaling cascade resulting in the inactivation of GSK3. Improper expression of Wnts has severe developmental consequences. For instance, overexpression of Wnt-1 results in the formation of two-headed tadpoles (36). Wnts are also important for the formation of mesodermal derivatives such as *Xenopus* myoblasts, in which dominant-negative expression of Wnt blocks MyoD expression and impairs skeletal muscle formation (23). It is tempting to speculate that Wnt signaling may also be important in the differentiation of mesodermal derivatives into adipocytes by regulating the activity of transcription factors such as the C/EBP family. As shown in Fig. 6B, C/EBP β and C/EBP δ contain GSK3 consensus sequences and may, like C/EBP α , be phosphorylated by GSK3. Our finding that lithium prevents adipogenesis supports the hypothesis that GSK3 activity is required for the differentiation of 3T3-L1 preadipocytes. Although the presence of receptors for Wnt on 3T3-L1 preadipocytes has not been reported, other cell models with adipogenic potential, NIH 3T3 cells and CH310T1/2, respond to Wnts (1, 8). We propose that regulation by GSK3 of C/EBP α phosphorylation and possibly preadipocyte differentiation is controlled not only by insulin, but also by Wnts and other ligands.

Insulin has rapid effects on the flow of carbon through metabolic pathways by regulating the activity of metabolic enzymes through stimulating their phosphorylation or dephosphorylation. Insulin also has longer-term effects on metabolism by altering gene expression to regulate the amount of enzyme or regulatory proteins available for metabolism. GSK3 acts as a node through which insulin signals to regulate both carbohydrate metabolism and adipocyte gene expression. The best-characterized role of GSK3 is in mediating the effects of insulin on glycogen metabolism through its phosphorylation and inhibition of glycogen synthase (28). It is now apparent that GSK3 also mediates the effects of insulin on adipocyte gene expression, since C/EBP α , a transcription factor required for acquisition of insulin sensitivity, is phosphorylated by GSK3. Dephosphorylation of the GSK3 sites in both glycogen synthase and C/EBP α appears to be mediated by PP1, since both activities are sensitive to okadaic acid and wortmannin. Thus, insulin appears to use reciprocal regulation of GSK3 and PP1 activities to coordinately regulate short- and long-term effects on adipocyte metabolism.

ACKNOWLEDGMENTS

This work is supported by a research grant to O.A.M. from the NIDDK, National Institutes of Health (RO1-DK51563). S.E.R. and R.L.E. are supported by predoctoral fellowships from the Natural Sciences and Engineering Research Council of Canada.

We thank Lawrence Argetsinger, Christin Carter-Su, Susanne Mandrup, Barbara Nicke, Jessica Schwartz, and John Williams for critical review of the manuscript. In addition, we thank Robert Lewis for phosphoamino acid analysis of peptide H215-K250, Mitchell Lazar for

suggesting that we treat differentiating preadipocytes with lithium, and M. Daniel Lane for providing antiserum to C/EBP α .

REFERENCES

- Bafico, A., A. Gazit, S. S. Wu-Morgan, A. Yaniv, and S. A. Aaronson. 1998. Characterization of Wnt-1 and Wnt-2 induced growth alterations and signaling pathways in NIH 3T3 fibroblasts. *Oncogene* **16**:2819–2825.
- Blum, H., H. Beier, and H. J. Gross. 1987. Improved silver staining of plant proteins, RNA and DNA in polyacrylamide gels. *Virology* **8**:93–99.
- Brady, M. J., F. J. Bourbonais, and A. R. Saltiel. 1998. The activation of glycogen synthase by insulin switches from kinase inhibition to phosphatase activation during adipogenesis in 3T3-L1 cells. *J. Biol. Chem.* **273**:14063–14066.
- Cadigan, K. M., and R. Nusse. 1997. Wnt signaling: a common theme in animal development. *Genes Dev.* **11**:3286–3305.
- Christy, R. J., K. H. Kaestner, D. E. Geiman, and M. D. Lane. 1991. CCAAT/enhancer binding protein gene promoter: binding of nuclear factors during differentiation of 3T3-L1 preadipocytes. *Proc. Natl. Acad. Sci. USA* **88**:2593–2597.
- Christy, R. J., V. W. Yang, J. M. Ntambi, D. E. Geiman, W. H. Landschulz, A. D. Friedman, Y. Nakabeppu, T. J. Kelly, and M. D. Lane. 1989. Differentiation-induced gene expression in 3T3-L1 preadipocytes: CCAAT/enhancer binding protein interacts with and activates the promoters of two adipocyte-specific genes. *Genes Dev.* **3**:1323–1335.
- Chu, B., F. Soncin, B. D. Price, M. A. Stevenson, and S. K. Calderwood. 1996. Sequential phosphorylation by mitogen-activated protein kinase and glycogen synthase kinase 3 represses transcriptional activation by heat shock factor-1. *J. Biol. Chem.* **271**:30847–30857.
- Cook, D., M. J. Fry, K. Hughes, R. Sumathipala, J. R. Woodgett, and T. C. Dale. 1996. Wingless inactivates glycogen synthase kinase-3 via an intracellular signalling pathway which involves a protein kinase C. *EMBO J.* **15**:4526–4536.
- Cornelius, P., O. A. MacDougald, and M. D. Lane. 1994. Regulation of adipocyte development. *Annu. Rev. Nutr.* **14**:99–129.
- Cross, D. A. E., D. R. Alessi, P. Cohen, M. Andjelkovich, and B. A. Hemmings. 1995. Inhibition of glycogen synthase kinase-3 by insulin mediated by protein kinase B. *Nature* **378**:785–789.
- Cross, D. A. E., P. W. Watt, M. Shaw, J. van der Kaay, C. P. Downes, J. C. Holder, and P. Cohen. 1997. Insulin activates protein kinase B, inhibits glycogen synthase kinase-3 and activates glycogen synthase by rapamycin-insensitive pathways in skeletal muscle and adipose tissue. *FEBS Lett.* **406**:211–215.
- Dale, T. C. 1998. Signal transduction by the Wnt family of ligands. *Biochem. J.* **329**:209–223.
- Darlington, G. J., S. E. Ross, and O. A. MacDougald. 1998. The role of C/EBP genes in adipocyte differentiation. *J. Biol. Chem.* **273**:30057–30060.
- Delcommenne, M., C. Tan, V. Gray, L. Rue, J. Woodgett, and S. Dedhar. 1998. Phosphoinositide-3-OH kinase-dependent regulation of glycogen synthase kinase 3 and protein kinase B/AKT by the integrin-linked kinase. *Proc. Natl. Acad. Sci. USA* **95**:11211–11216.
- Dignam, J. D., R. M. Lebovitz, and R. G. Roeder. 1983. Accurate transcription initiation by RNA polymerase II in a soluble extract from isolated mammalian nuclei. *Nucleic Acids Res.* **11**:1475–1489.
- Eldar-Finkelman, H., G. M. Argast, O. Foord, E. H. Fischer, and E. G. Krebs. 1996. Expression and characterization of glycogen synthase kinase-3 mutants and their effect on glycogen synthase activity in intact cells. *Proc. Natl. Acad. Sci. USA* **93**:10228–10233.
- El-Jack, A. K., J. K. Hamm, P. F. Pilch, and S. R. Farmer. 1999. Reconstitution of insulin-sensitive glucose transport in fibroblasts requires expression of both PPAR γ and C/EBP α . *J. Biol. Chem.* **274**:7946–7951.
- Erickson, R. L., K. A. Longo, S. E. Ross, N. Hemati, and O. A. MacDougald. Structure and function of C/EBP α . *In* Proceedings of the Steenbock Symposium. IOS Press, Amsterdam, The Netherlands, in press.
- Friedman, A. D., and S. L. McKnight. 1990. Identification of two polypeptide segments of CCAAT/enhancer-binding protein required for transcriptional activation of the serum albumin gene. *Genes Dev.* **4**:1416–1426.
- Green, H., and M. Meuth. 1974. An established pre-adipose cell line and its differentiation in culture. *Cell* **3**:127–133.
- Groblewski, G. E., M. J. Wishart, M. Yoshida, and J. A. Williams. 1996. Purification and identification of a 28-kDa calcium-regulated heat-stable protein. *J. Biol. Chem.* **271**:31502–31507.
- Hemati, N., S. E. Ross, R. L. Erickson, G. E. Groblewski, and O. A. MacDougald. 1997. Signaling pathways through which insulin regulates CCAAT/enhancer binding protein α (C/EBP α) phosphorylation and gene expression in 3T3-L1 adipocytes: correlation with GLUT4 gene expression. *J. Biol. Chem.* **272**:25913–25919.
- Hoppler, S., J. D. Brown, and R. T. Moon. 1996. Expression of a dominant-negative Wnt blocks induction of MyoD in *Xenopus* embryos. *Genes Dev.* **10**:2805–2817.
- Hwang, C.-S., S. Mandrup, O. A. MacDougald, D. E. Geiman, and M. D. Lane. 1996. Transcriptional activation of the *obese* gene by CCAAT/enhancer binding protein α . *Proc. Natl. Acad. Sci. USA* **93**:873–877.

25. **Kaestner, K. H., R. J. Christy, and M. D. Lane.** 1990. Mouse insulin-responsive glucose transporter gene: characterization of the gene and trans-activation by the CCAAT/enhancer binding protein. *Proc. Natl. Acad. Sci. USA* **87**:251–255.
26. **Kennelly, P. J., and E. G. Krebs.** 1991. Consensus sequences as substrate specificity determinants for protein kinases and protein phosphatases. *J. Biol. Chem.* **266**:15555–15558.
27. **Klein, P. S., and D. A. Melton.** 1996. A molecular mechanism for the effect of lithium on development. *Proc. Natl. Acad. Sci. USA* **93**:8455–8459.
28. **Lawrence, J. C., and P. J. Roach.** 1997. New insights into the role and mechanism of glycogen synthase activation by insulin. *Diabetes* **46**:541–547.
29. **Lin, F.-T., O. A. MacDougald, A. M. Diehl, and M. D. Lane.** 1993. A 30 kilodalton alternative translation product of the CCAAT/enhancer binding protein α message: transcriptional activator lacking antimetabolic activity. *Proc. Natl. Acad. Sci. USA* **90**:9606–9610.
30. **MacDougald, O. A., P. Cornelius, F.-T. Lin, S. S. Chen, and M. D. Lane.** 1994. Glucocorticoids reciprocally regulate expression of the CCAAT/enhancer-binding protein α and δ genes in 3T3-L1 adipocytes and white adipose tissue. *J. Biol. Chem.* **269**:19041–19047.
31. **MacDougald, O. A., P. Cornelius, R. Liu, and M. D. Lane.** 1995. Insulin regulates transcription of the CCAAT/enhancer binding protein (C/EBP) α , β , and δ genes in fully-differentiated 3T3-L1 adipocytes. *J. Biol. Chem.* **270**:647–654.
32. **MacDougald, O. A., and M. D. Lane.** 1995. Transcriptional regulation of gene expression during adipocyte differentiation. *Annu. Rev. Biochem.* **64**:345–373.
33. **Mahoney, C. W., J. Shuman, S. L. McKnight, H.-C. Chen, and K.-P. Huang.** 1992. Phosphorylation of CCAAT-enhancer binding protein by protein kinase C attenuates site-selective DNA binding. *J. Biol. Chem.* **267**:19396–19403.
34. **Mandrup, S., and M. D. Lane.** 1997. Regulating adipogenesis. *J. Biol. Chem.* **272**:5367–5370.
35. **McKnight, S. L.** 1992. CCAAT/enhancer binding protein, p. 771–795. *Transcriptional regulation.* Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.
36. **McMahon, A. P., and R. T. Moon.** 1989. Ectopic expression of the proto-oncogene int-1 in *Xenopus* embryos leads to duplication of the embryonic axis. *Cell* **58**:1075–1084.
37. **Moitra, J., M. M. Mason, M. Olive, D. Krylov, O. Gavrilova, B. Marcus-Samuels, L. Feigenbaum, E. Lee, T. Aoyama, M. Eckhaus, M. L. Reitman, and C. Vinson.** 1998. Life without white fat: a transgenic mouse. *Genes Dev.* **12**:3168–3181.
38. **Moule, S. K., G. I. Welsh, N. J. Edgell, E. J. Foulstone, C. G. Proud, and R. M. Denton.** 1997. Regulation of protein kinase B and glycogen synthase kinase-3 by insulin and β -adrenergic agonists in rat epididymal fat cells. *J. Biol. Chem.* **272**:7713–7719.
39. **Nerlov, C., and E. B. Ziff.** 1995. CCAAT/enhancer binding protein- α amino acid motifs with dual TBP and TFIIB binding ability co-operate to activate transcription in both yeast and mammalian cells. *EMBO J.* **14**:4318–4328.
40. **Nerlov, C., and E. B. Ziff.** 1994. Three levels of functional interaction determine the activity of CCAAT/enhancer binding protein- α on the serum albumin promoter. *Genes Dev.* **8**:350–362.
41. **Pei, D., and C. Shih.** 1991. An “attenuator domain” is sandwiched by two distinct transactivation domains in the transcription factor C/EBP. *Mol. Cell. Biol.* **11**:1480–1487.
- 41a. **Piwien de Pilipuk, G., S. E. Ross, O. A. MacDougald, and J. Schwartz.** Unpublished data.
42. **Preece, A.** 1972. *A manual for histologic technicians.* Little Brown, Boston, Mass.
43. **Roach, P. J.** 1990. Control of glycogen synthase by hierarchical protein phosphorylation. *FASEB J.* **4**:2961–2968.
44. **Shuman, J. D., J. Cheong, and J. E. Coligan.** 1997. ATF-2 and C/EBP α can form a heterodimeric DNA binding complex *in vitro*. *J. Biol. Chem.* **272**:12793–12800.
45. **Shuman, J. D., C. R. Vinson, and S. L. McKnight.** 1990. Evidence of changes in protease sensitivity and subunit exchange rate on DNA binding by C/EBP. *Science* **249**:771–778.
46. **Spiegelman, B. M., and J. S. Flier.** 1996. Adipogenesis and obesity: rounding out the big picture. *Cell* **87**:377–389.
47. **Student, A. K., R. Y. Hsu, and M. D. Lane.** 1980. Induction of fatty acid synthetase synthesis in differentiating 3T3-L1 preadipocytes. *J. Biol. Chem.* **255**:4745–4750.
48. **Volle, D. J., J. A. Fulton, O. V. Chaika, K. McDermott, H. Luang, L. A. Steinke, and R. E. Lewis.** 1999. Phosphorylation of the kinase suppressor of Ras by associated kinases. *Biochemistry* **38**:5130–5137.
49. **Wickelgren, I.** 1998. Obesity: how big a problem? *Science* **280**:1364–1367.
50. **Wu, Z., E. D. Rosen, R. Brun, S. Hauser, G. Adelmant, A. E. Troy, C. McKeon, G. J. Darlington, and B. M. Spiegelman.** 1999. Cross-regulation of C/EBP α and PPAR γ controls the transcriptional pathway of adipogenesis and insulin sensitivity. *Mol. Cell* **3**:151–158.