

Interactions of Pex7p and Pex18p/Pex21p with the Peroxisomal Docking Machinery: Implications for the First Steps in PTS2 Protein Import

Katharina Stein, Annette Schell-Steven, Ralf Erdmann,* and Hanspeter Rottensteiner

Freie Universität Berlin, Fachbereich Biologie, Chemie, Pharmazie, D-14195 Berlin, Germany

Received 19 February 2002/Returned for modification 28 March 2002/Accepted 23 May 2002

Peroxisomal PTS2-dependent matrix protein import starts with the recognition of the PTS2 targeting signal by the import receptor Pex7p. Subsequently, the formed Pex7p/cargo complex is transported from the cytosol to the peroxisomal docking complex, consisting of Pex13p and Pex14p. In *Saccharomyces cerevisiae*, the latter event is thought to require the redundant Pex18p and Pex21p. Here we mapped the Pex7p interaction domain of Pex13p to its N-terminal 100 amino acids. Pex18p and Pex21p also interacted with this region, albeit only in the presence of Pex7p. Expression of an N-terminally deleted version of Pex13p in a *pex13Δ* mutant failed to restore growth on fatty acids due to a specific defect in the import of PTS2-containing proteins. We further show by yeast two-hybrid analysis, coimmunoprecipitation, and in vitro binding assays that Pex7p can bind Pex13p and Pex14p in the absence of Pex18p/Pex21p. The PTS2 protein thiolase was shown to interact with Pex14p but not with Pex13p in a Pex7p- and Pex18p/Pex21p-dependent manner, suggesting that only Pex14p binds cargo-loaded PTS2 receptor. We also found that the cytosolic Pex7p/thiolase-containing complex includes Pex18p. This complex accumulated in docking mutants but was absent in cells lacking Pex18p/Pex21p, indicating that Pex18p/Pex21p are required already before the docking event.

Peroxisomes are ubiquitous organelles of eucaryotic cells and fulfill a number of biochemical functions, including the β -oxidation of fatty acids (51). Peroxisome biogenesis is dependent on a class of genes that is conserved among species, the *PEX* genes (19, 40). The corresponding gene products, the peroxins, are involved in specific aspects of peroxisome formation, peroxisome proliferation, membrane protein insertion, and matrix protein import (14, 21). Most of the peroxins are required for matrix protein import, which can occur via two different pathways, the peroxisomal targeting signal type 1 (PTS1)-dependent one and the PTS2-dependent one. The majority of matrix proteins possess a PTS1 signal, which is located at the extreme C terminus and is composed of the tripeptide SKL or conservative variants thereof (18, 24). PTS2-containing proteins include 3-ketoacyl-coenzyme A thiolase, which in *Saccharomyces cerevisiae* is encoded by the *FOX3* gene. PTS2 targeting signals are located close to the N terminus and were proposed to follow the consensus sequence H/R-L-X₅-H-L (47, 48).

The targeting signals are recognized in the cytosol by the PTS1- and PTS2-specific factors Pex5p and Pex7p, respectively, which in turn deliver the cargo proteins to a docking complex at the peroxisomal membrane. Upon docking, the two pathways seem to merge, as the components of the docking complex, Pex14p and Pex13p, provide binding sites for both Pex5p and Pex7p (1, 16, 32). In the case of Pex5p, interaction with Pex13p and Pex14p has been shown to be direct (2, 41, 50). Since Pex13p and Pex14p are nonredundant peroxins, it follows that they act either together in the same complex or in

a consecutive fashion upon protein translocation. Evidence for Pex13p being in a complex that is distinct from the Pex14p-containing complex was recently provided upon purification of both proteins from rat liver peroxisomes (35). In *S. cerevisiae*, Pex13p is required for the targeting of Pex14p to the peroxisomal membrane (16), and thus, some of the defects in PTS2-dependent import that are observed in *pex13Δ* mutant cells may in fact be due to the absence of peroxisomal Pex14p from these cells.

PTS2 protein import specifically requires cytosolic factors in addition to Pex7p. These include the redundant Pex18p/Pex21p in *S. cerevisiae* (34); Pex5pL, the long isoform of Pex5p in higher eukaryotes (5, 31); and Pex20p of *Yarrowia lipolytica* (49). These PTS2-specific proteins possess only a weak overall sequence similarity, but recent reports suggest that they might perform a similar function. Pex5pL and Pex18p/Pex21p are capable of interacting with Pex7p via a conserved short motif and are thought to deliver cargo-loaded receptor to the peroxisomal membrane, whereas the binding of the PTS2 signal is accomplished by Pex7p (7, 27, 30). Support for a conserved function of these proteins was presented recently by the partial complementation of a *pex18Δ pex21Δ* mutant with Pex20p (8). However, Pex7p has not yet been identified in *Y. lipolytica*, and thus, Pex20p may contain the activities of both Pex7p and Pex18p/Pex21p. For *Y. lipolytica*, it has been shown that Pex20p is involved in the cytosolic oligomerization of thiolase, which seems to be a prerequisite for import. However, Pex20p might also be involved in later events, as it can directly interact with the intraperoxisomal Pex8p (46).

Here we present a functional analysis of the PTS2-specific peroxins Pex7p, Pex18p, and Pex21p and their docking factors Pex13p and Pex14p. The Pex7p-binding domain in Pex13p was mapped to the extreme N terminus of Pex13p, which fails to interact with Pex5p and Pex14p. The physiological relevance of

* Corresponding author. Mailing address: Freie Universität Berlin, Institut für Chemie/Biochemie, Thielallee 63, D-14195 Berlin, Germany. Phone: 43-30-838 52935. Fax: 43-30-838 52936. E-mail: ralferdman@zedat.fu-berlin.de.

TABLE 1. *S. cerevisiae* strains used

Strain	Description	Source or reference
UTL-7A	MAT α <i>leu2-3,112 ura3-52 trp1</i>	10
HF7c	MAT α <i>ura3-52 his3-200 lys2-801 ade2-101 trp1-901 leu2-3,112 gal4-542 gal80-538 LYS2::GAL1-HIS3 URA3::(GAL4 17-mers)₃-CYC1-LacZ</i>	11
PJ69-4A	MAT α <i>ura3-52 his3-200 trp1-901 leu2-3,112 gal4Δ gal80Δ LYS2::GAL1-HIS3 met2::GAL7-lacZ GAL2-ADE2</i>	P. James ^a
UTL-7Apex7 Δ	<i>pex7Δ::LEU2</i>	26
UTL-7Apex13 Δ	<i>pex13Δ::kanMX4</i>	23
UTL-7Apex14 Δ	<i>pex14Δ::LEU2</i>	1
UTL-7Apex18 Δ	<i>pex18Δ::kanMX4</i>	W. Kunau ^b
UTL-7Apex21 Δ	<i>pex21Δ::loxP</i>	W. Kunau
HF7cpex7 Δ	<i>pex7Δ::LEU2</i>	W. Kunau
HF7cpex21 Δ	<i>pex21Δ::loxP</i>	W. Kunau
yKat36	UTL-7Apex18 Δ :: <i>kanMX4 pex21Δ::loxP</i>	This study
yKat110	UTL-7Apex14 Δ :: <i>LEU2 pex18Δ::kanMX4 pex21Δ::loxP</i>	This study
yKat12	HF7cpex18 Δ :: <i>kanMX4 pex21Δ::loxP</i>	This study
yHPR251	UTL-7A [pHPR131]	This study
yHPR252	UTL-7Apex7 Δ :: <i>LEU2</i> [pHPR131]	This study
yHPR255	UTL-7A [pHPR132]	This study
yKat92	UTL-7Apex13 Δ :: <i>kanMX4</i> [pHPR131]	This study
yKat111	UTL-7Apex18 Δ :: <i>kanMX4 pex21Δ::loxP</i> [pHPR131]	This study
yAS3	UTL-7Apex7 Δ <i>PEX18-TAP TRP1</i>	This study
yAS4	UTL-7Apex21 Δ <i>PEX18-TAP TRP1</i>	This study

^a Madison, Wis.^b Bochum, Germany.

this domain for PTS2-dependent protein import was addressed. We further demonstrate that Pex7p and not Pex18p/Pex21p actually docks to the peroxisomal docking complex and provide evidence that Pex14p represents the initial docking protein. The function of Pex18p and Pex21p will be addressed, and our findings will be discussed in terms of a sequence of events for the early steps in PTS2 protein import.

MATERIALS AND METHODS

Strains, media, and general methods. *Escherichia coli* strain DH5 α was used for all plasmid amplifications and isolations, and *E. coli* strain C41 (DE3) (obtained from J. Walker, Medical Research Council, Cambridge, United Kingdom) was used for heterologous expression of recombinant fusion proteins. The yeast strains used in this study are listed in Table 1. *PEX18* was deleted from strains HF7cpex21 Δ and UTL-7Apex21 Δ by using a *kanMX4*-based disruption cassette that had been amplified from genomic DNA of strain HF7cpex18 Δ with primer pair RE221/222. *PEX14* was similarly deleted from strain yKat110 by amplifying a *LEU2*-based cassette with primer pair RE87/88 and genomic DNA of strain UTL-7Apex14 Δ as a template. The authenticity of each gene deletion was confirmed by PCR. *EcoRV*-linearized pHPR131 (expressing PTS2-DsRed) was stably integrated into the *trp1* locus of strains UTL-7A (yHPR251), UTL-7Apex7 Δ (yHPR252), UTL-7Apex13 Δ (yKAT92), and UTL-7Apex18 Δ pex21 Δ (yKat111). Plasmid pHPR132 (expressing DsRed) was similarly integrated into the *trp1* locus of UTL-7A (yHPR255). *PEX18* was genomically tagged with the TAP tag as described previously (38). In brief, primer pair RE301/302 was used to amplify the TAP tag cassette plus the *Kluyveromyces lactis TRP1* marker gene from plasmid pBS1479. The resulting PCR product was then used to transform strains UTL-7Apex7 Δ (yAS3) and UTL-7Apex21 Δ (yAS4). Correct integration of the cassette was verified by PCR.

Standard media for the cultivation of yeast and bacterial strains were prepared as described previously (10, 42). For the induction of peroxisomes, cells were grown in synthetic dextrose medium (SD) containing 0.3% glucose to mid-log phase, shifted into oleic acid medium (YNO), and incubated for 13 h. YNO contained 0.1% oleic acid, 0.05% Tween 40, 0.1% yeast extract, 0.67% yeast nitrogen base, and amino acids as required and was adjusted to pH 6.0 with KOH. For induction of the *CUP1* promoter, 25 mg of CuSO₄/liter was added to the YNO medium. Oleic acid plates were prepared as described previously and contained 0.1% (wt/vol) oleic acid, 0.5% (wt/vol) Tween 80, 0.1% yeast extract, 0.67% yeast nitrogen base, amino acids, and 0.5% potassium phosphate buffer (pH 6). Standard recombinant DNA techniques, including enzymatic modifica-

tion of DNA, transformation, and cultivation, were performed essentially as described previously (42). Yeast whole-cell extracts were prepared from 30 mg of cells according to a published protocol (55).

Plasmids and oligonucleotides. The plasmids and oligonucleotides used are listed in Tables 2 and 3, respectively. Unless otherwise stated, all PCRs were performed with genomic DNA of a wild-type strain as a template and products were subcloned into *EcoRV*-cut pBluescript SK(+) (Stratagene, La Jolla, Calif.). To construct the *PEX13-GFP* fusions, the complete *PEX13* open reading frame and a truncated form (representing amino acids [aa] 151 to 386) were amplified with primer pairs RE 432/435 and RE 550/435, respectively, cut with *Bam*HI and *Eco*RI, and cloned into the appropriately digested pUG35 (29), resulting in pMS9 and pKAT136.

Expression of full-length and truncated Pex13p from the *PEX13* promoter was achieved as follows. The *PEX13* promoter (base pairs -363 to +6) was amplified (RE 98/99) and cloned as a *Bam*HI/*Eco*RI fragment into pSK(+) (pKat46). The complete *PEX13* open reading frame, amplified with primer pair RE 421/423, and the truncated version (representing aa 56 to 386), amplified with primer pair RE 422/423, were digested with *Sph*I/*Hind*III and cloned into the *Sph*I/*Hind*III-digested pKat46, resulting in pKat111 and pKat112. The assembled constructs were excised from pKat111 and pKat112 with *Bam*HI/*Hind*III and ligated into the appropriately cut yeast expression vector pRS-FOXP-TERM (C. Clayton, Heidelberg, Germany), resulting in pKat113 and pKat122, respectively.

The PTS2-DsRed expression construct (pHPR131) was cloned by assembling the following fragments into *Xba*I/*Pst*I-cut YIplac204 (15): the *ADH2* promoter, excised from plasmid pADH2PIP2, as an *Xba*I/*Nde*I fragment (3); the amino-terminal 16 aa of Fox3p, obtained from pPTS2-GFP-prA (unpublished) by using primer pair RE 188/189, as an *Nde*I/*Nco*I fragment; and DsRed, amplified from pDsRed (Clontech, Palo Alto, Calif.) with primer pair RE 197/198 and subcloned into pBluescript SK(+) (pHPR126), as an *Nco*I/*Pst*I fragment. The DsRed expression vector lacking a PTS2 (pHPR132) was similarly assembled in *Eco*RI/*Pst*I-cut YIplac204 by using the DsRed-containing *Nco*I/*Pst*I fragment of pHPR126 and the *ADH2* promoter from pADH2-OAF1 as an *Eco*RI/*Nco*I fragment (39).

For the bacterial expression of a synthetic PTS2 protein (pRB107), PTS2-GFP-prA was amplified from pPTS2-GFP-prA with primer pair RE 188/415 and cloned as a *Bam*HI/*Eco*RI fragment into pGEX4T-2 (Amersham Pharmacia, Freiburg, Germany). To express Pex7p in *E. coli*, *PEX7* was amplified with primer pair RE 79/80 and cloned as a *Sac*I/*Pst*I fragment into the appropriately cut pQE31 (Qiagen, Hilden, Germany) or pMAL-c2X (New England Biolabs, Beverly, Mass.), yielding QE31-PEX7 and pKat128, respectively. To generate a His₆-Pex14p fusion construct (QE31-PEX14), *PEX14* was amplified with primer pair RE 87/88 and cloned into pQE31 as a *Bam*HI/*Pst*I fragment. The GST-

TABLE 2. Plasmids used

Plasmid	Description	Primer pair	Source or reference
pJR233	<i>MLS1</i> pr-GFP-SKL		24
YEpmcPex7	<i>CUP1</i> pr-myc-PEX7		36
pPC86-PEX5			1
pPC86-PEX14			1
pPC97-PEX7			1
pPC97-FOX3			36
pBS1479	<i>K. lactis</i> TRP1-marked TAP tag cassette		38
pHPR126	<i>DsRed</i> in SK(+)	RE197/198	This study
pHPR131	<i>ADH2</i> pr-PTS2- <i>DsRed</i>		This study
pHPR132	<i>ADH2</i> pr- <i>DsRed</i>		This study
pQE31-PEX7	His ₆ -PEX7	RE79/80	This study
pQE31-PEX14	His ₆ -PEX14	RE87/88	This study
pKat136	Pex13p ₁₅₁₋₃₈₆ -GFP	RE550/435	This study
pKat46	<i>PEX13</i> _{pr} in SK(+)	RE98/99	This study
pKat111	<i>PEX13</i> _{pr} - <i>PEX13</i> ₁₋₃₈₆ in SK(+)	RE421/423	This study
pKat112	<i>PEX13</i> _{pr} - <i>PEX13</i> ₅₆₋₃₈₆ in SK(+)	RE422/423	This study
pKat113	<i>PEX13</i> _{pr} - <i>PEX13</i> ₁₋₃₈₆ in pRSTERM		This study
pKat122	<i>PEX13</i> _{pr} - <i>PEX13</i> ₅₆₋₃₈₆ in pRSTERM		This study
pKat128	MBP-PEX7	RE79/80	This study
pKat118	GST-Pex13p ₁₋₁₀₀	RE298/379	This study
pKat16	pPC86-PEX13 (1-151)	RE25/32	This study
pKat17	pPC86-PEX13 (1-166)	RE25/33	This study
pKat18	pPC86-PEX13 (1-264)	RE25/34	This study
pKat19	pPC86-PEX13 (1-280)	RE25/35	This study
pKat20	pPC86-PEX13 (1-310)	RE25/36	This study
pKat21	pPC86-PEX13 (1-340)	RE25/37	This study
pKat22	pPC86-PEX13 (1-370)	RE25/52	This study
pKat23	pPC86-PEX13 (1-386)	RE25/38	This study
pKat24	pPC86-PEX13 (151-386)	RE26/38	This study
pKat25	pPC86-PEX13 (166-386)	RE27/38	This study
pKat26	pPC86-PEX13 (264-386)	RE28/38	This study
pKat27	pPC86-PEX13 (280-386)	RE29/38	This study
pKat28	pPC86-PEX13 (310-386)	RE30/38	This study
pKat29	pPC86-PEX13 (340-386)	RE31/38	This study
pKat30	pPC86-PEX13 (370-386)	RE53/38	This study
pKat31	pPC97-PEX13 (1-151)	RE25/32	This study
pKat86	pPC86-PEX13 (1-55)	RE25/378	This study
pKat87	pPC86-PEX13 (1-100)	RE25/379	This study
pKat88	pPC97-PEX13 (1-55)	RE25/378	This study
pKat89	pPC97-PEX13 (1-100)	RE25/379	This study
pKat90	pPC86-PEX13 (55-100)	RE380/379	This study
pKat91	pPC97-PEX13 (55-100)	RE380/379	This study
pKat59	pPC86-PEX21	RE173/174	This study
pKat66	pPC86-PEX18	RE169/170	This study
pKat60	pPC97-PEX21	RE173/174	This study
pKat68	pPC97-PEX18	RE169/170	This study
pKat146	pPC86-FOX3		This study
pMS9	Pex13p ₁₋₃₈₆ -GFP	RE432/435	This study
pRB107	GST-PTS2-GFP-prA	RE188/415	This study

Pex13p₁₋₁₀₀ construct (pKat118) was cloned by amplifying the Pex13p-encoding fragment with primer pair RE 298/379 and ligating the resulting *EcoRI/XbaI* fragment into pGEX4T-1.

Yeast two-hybrid assays. The two-hybrid assay was based on the method of Fields and Song (12). Selected *PEX* genes or truncations thereof were fused to the DNA-binding domain or transcription activation domain of Gal4p in the vectors pPC86 and pPC97 (6). To construct the Gal4p-AD-Pex13p fusions, *EcoRI/SpeI* fragments of the various *PEX13* fragments that had been amplified from genomic DNA were subcloned into the appropriately cut pPC86 (Table 2). To construct the Gal4p-BD-Pex13p fusions, the *PEX13* fragments were excised from the pPC86-PEX13 constructs with *SmaI/SpeI* and subcloned into *SmaI/SpeI*-digested pPC97 (Table 2). *PEX18* and *PEX21* were amplified from genomic DNA and subcloned into *Sall/SacI*-digested pPC86 and pPC97, respectively (Table 2). A *Sall/SacI* fragment of pPC97-FOX3 containing the *FOX3* open reading frame was subcloned into an appropriately cut pPC86, resulting in pPC86-FOX3 (pKat146). Cotransformation of two-hybrid plasmids into HF7c was performed according to the method of Schiestl and Gietz (44). Double

transformants were selected on SD synthetic medium without tryptophan and leucine. Histidine auxotrophy of transformed HF7c was determined by growth on selective plates lacking leucine, tryptophan, and histidine but containing 1 or 5 mM 3-aminotriazole.

Antibodies and Western blotting. To generate polyclonal antibodies against Pex7p, a His₆-tagged Pex7p was expressed in *E. coli* strain C41 (DE3) and purified under denaturing conditions according to the manufacturer's protocol (Qiagen). This protein was subsequently used to immunize rabbits at Eurogentec (Seraing, Belgium). The raised antiserum was specific, as it was able to detect Pex7p in whole-cell extracts from a wild type but not from the respective *pex7Δ* strain (Fig. 9C, lower panel). The other antibodies used have been described previously, namely, anti-Fox3p (9), anti-Cta1p (20), anti-Pex13p (16), anti-Pex14p (1), anti-maltose-binding protein (MBP; New England Biolabs), anti-protein A (Sigma, St. Louis, Mo.), and anti-myc from 9E10 cell lines (16). Immunoblotting was performed according to standard protocols. Horseradish peroxidase-coupled anti-rabbit or anti-mouse immunoglobulin G (IgG), in com-

TABLE 3. Oligonucleotides used

Designation	Sequence (5' to 3')
RE 25GTGAATTCGGATCCATATGTCATCCACAGCAGTACCA
RE 26GTGAATTCGGATCCATATGTTAATAGAAAGTTTGATAGGC
RE 27GTGAATTCGGATCCATATGCTGGAATCTACTTATATGGCC
RE 28GTGAATTCGGATCCATATGCCGCTACTATTTTTTTTGTATG
RE 29GTGAATTCGGATCCATATGAACAAAATTTATTAATAAACTACAG
RE 30GTGAATTCGGATCCATATGTTTGCAAGAGCGTTATATGAT
RE 31GTGAATTCGGATCCATATGAAAGACCCTCTTGGGAGGGAT
RE 32GCTCTAGAAGTAACTGAAAGGTGGCCTTCGT
RE 33GCTCTAGAAGTACAGCATTTGTGCAAATCCAGT
RE 34GCTCTAGAAGTACGGTTTCCATGAAATTTTCCT
RE 35GCTCTAGAAGTAGTGTAGTAGATATGGAAAACC
RE 36GCTCTAGAAGTAAATTTAACTTCGAAGGATC
RE 37GCTCTAGAAGTATTTTCTTACTCAAATTTGCCAT
RE 38GCTCTAGAAGTCTAGTGTGTACGCGTTTCATC
RE 52GCTCTAGAAGTATGATCTCAATATAGTTATA
RE 53GTGAATTCGGATCCATATGATAAAAAGACGGAAGAAAATTGAG
RE 79GCTCTAGAGACTACATATGCTCAGATATCACATGCAAGG
RE 80GGGGTACCCTCGAGCTGCAGTCAACCTAAGCCGTTCCATAC
RE 87GCTCTAGAGCTAGCAAGGATCCAATGAGTGACGTGGTTCAG
RE 88GGGGTACCGAATCCCTGCAGCTATGGGATGGAGTCTTC
RE 98CGGGTACCCTGCAGCTATTTATTCATAGTGTC
RE 99CGGAATTCGCATGCTGACATCGCAGGTATTGTTATAGT
RE 169GCGTCTGACTATGAATAGTAACCGATGC
RE 170GCGAGCTCTTAAGCAATTTCTGTCTTCAAC
RE 173GCGTCTGACTATGCCAGTGTCTGCCAT
RE 174GCGAGCTCTCAATCAAGTATGTCTTTGTG
RE 188CTGCAGTTAATTCGCGTCTACTTTCGG
RE 189CATATGCTCAAAGACTACAAAG
RE 197GCGGCCGACCATGGTGAGGTCTTCCAAGAATGTT
RE 198GGATCCGTCGCGGCCGCTAAAGGAAC
RE 221GTATAATCAGGTATGTAAGGG
RE 222CGACAATAAGTTCAGAAAAG
RE 298CGGAATTCATATGTCATCCACAGCAGTACCA
RE 301AACTGGGCTGGTCTTGAGTTCCATGATGTTGAAGACAGAATTGCTTCCATGGAAAAGAGAAG
RE 302TCTCTTGAAAGTGGTTAAAAGTGTATATATCTGAAATTCATGGTACGACTACTATAGGG
RE 378GCTCTAGAAGTCTAAGCAGACTCACTTGCCT
RE 379GCTCTAGAAGTCTATATAGAGTTTACTACTATACGG
RE 380GTGAATTCGGATCCATATGCCCCGAAGTTTTGCGCGG
RE 415GGATCCTCTCAAAGACTACAAAGTATC
RE 421GCATGCGGCGGCCGCTATCCACAGCAGTACCA
RE 422GCATGCGGCGGCCGCCCCGAAGTTTTGCGCGG
RE 423AAGCTTCTAGTGTGTACGCGTTTCAT
RE 432GAGGATCCTGCGATGTCATCCACA
RE 435TCTGAATTCGTGTGTACGCGTTTCATC
RE 550GAGGATCCTGCGATGTTAATAGAAAGTTTGATAGGC

bination with the ECL system (Amersham Pharmacia), was used to detect immunoreactive complexes.

Coimmunoprecipitation. Coimmunoprecipitation of *myc*-Pex7p was performed essentially as described previously (36), with the exception that breakage of the copper-induced cells was achieved by vortexing a suspension of 1 g of cells, 3 volumes of solution A (50 mM Tris-HCl [pH 7.5], 50 mM NaCl, 0.2% Triton X-100) that contained the Complete protease inhibitor cocktail (Roche, Mannheim, Germany), and 3 volumes of glass beads (diameter, 0.5 mm; Sigma) for 30 min at 4°C on an IKA-Vibrax VXR (Vibrax, Staufen, Germany). To precipitate Pex18p-TAP, soluble extracts obtained from 3 g of cells (suspended in 2.5 volumes of solution A) were incubated with IgG Sepharose (Amersham Pharmacia) followed by three washing steps with solution A. At that point, aliquots of each sample were removed for analysis of bound Pex18p-TAP. The remainder of the samples were adjusted to 10 mM dithiothreitol and 1 mM EDTA and incubated with 3 U of recombinant tobacco etch virus (TEV) protease (Life Technologies, Karlsruhe, Germany) for 2 h at room temperature with gentle agitation to release Pex18p from the matrix. The resulting eluates were analyzed for the presence of Pex7p and Fox3p.

In vitro binding assays. To analyze the Pex7p-Pex14p interaction, the soluble fraction of bacterially expressed His₆-Pex14p (QE31-PEX14) was incubated with nickel-nitrilotriacetic acid (Ni-NTA) matrix (Invitrogen, De Schelp, The Netherlands) for 2 h at 4°C. After being washed with column buffer (50 mM

NaH₂PO₄, 300 mM NaCl, 20 mM imidazole [pH 8]) in a column, equal amounts of the soluble fractions of bacterially expressed MBP-Pex7p (pKat128) or MBP alone (pMAL-c2X) were added to the column and incubated with the matrix for 1 h. After being washed with column buffer, the proteins were eluted with elution buffer (50 mM NaH₂PO₄, 300 mM NaCl, 500 mM imidazole [pH 8]). The eluted samples were analyzed by Coomassie brilliant blue staining or immunoblot analysis.

The *in vitro* binding of Pex7p to Pex13p was analyzed as follows. The soluble fractions of glutathione S-transferase (GST) and GST-Pex13p₁₋₁₀₀ that had been expressed in C41 (DE3) from plasmids pGEX4T-1 and pKat118, respectively, were bound for 2 h at 4°C on a glutathione Sepharose 4B matrix (Amersham Pharmacia). After washing the matrix with 1× phosphate-buffered saline (137 mM NaCl, 2.7 mM KCl, 4.3 mM Na₂HPO₄·7H₂O, 1.4 mM KH₂PO₄), the soluble fraction of bacterially expressed His₆-Pex7p (QE31-PEX7) was added and incubated with the matrix for 1 h at 4°C with gentle rotation. Alternatively, equal amounts of yeast lysates from *myc*-Pex7p-expressing UTL-7A or UTL-7A*pex18Δpex21Δ* cells were added. After washing the matrix with 1× phosphate-buffered saline, the proteins were eluted with 10 mM glutathione in 50 mM Tris-HCl (pH 8). The eluted samples were analyzed by Coomassie brilliant blue staining or immunoblot analysis.

The binding of PTS2 protein to Pex7p was tested by loading soluble extracts containing either GST-PTS2 protein (pRB107) or GST (pGEX4T-2) onto MBP-

Pex7p that had been purified by using the pMAL purification system from New England Biolabs. After being washed with 10 bed volumes of column buffer (20 mM Tris-HCl [pH 7.4], 200 mM NaCl, 1 mM EDTA), MBP-Pex7p was eluted with column buffer containing 10 mM maltose. The eluates were analyzed for the presence of GST-containing proteins by immunoblotting with an antibody directed against GST-Pex14p (1).

Miscellaneous. The analysis of live cells for DsRed and green fluorescent protein (GFP) fluorescence was performed according to the method of Westermann and Neupert (53). β -Galactosidase activities were assayed according to manufacturer's instructions and expressed as micromoles of CPRG (chlorophenol red- β -D-galactopyranoside) hydrolyzed per minute per cell (Clontech).

RESULTS

Pex7p interacts with the cytoplasmic N-terminal domain of Pex13p. We have shown previously that Pex13p functionally interacts with the PTS2 cargo receptor Pex7p (16). To further characterize the binding of Pex7p to Pex13p, a series of N- and C-terminal deletion constructs of Pex13p were tested in a yeast two-hybrid assay for interaction with Pex7p. The full-length Pex13p (aa 1 to 386) as well as the two C-terminal truncations that lacked part of the SH3 domain (aa 1 to 340) or the complete domain (aa 1 to 310) were able to interact with Pex7p (Fig. 1), thereby demonstrating that the SH3 domain of Pex13p is dispensable for the interaction of Pex13p with Pex7p. Interaction was also observed for even shorter Pex13p variants that lacked the cytosolically oriented C-terminal domain in total, with or without transmembrane domain 2 (TMD 2; aa 1 to 280 and 1 to 264, respectively). By further removing the internal loop (aa 1 to 166) and TMD 1 (aa 1 to 151) of Pex13p, binding to Pex7p remained proficient. On the other hand, removing these 151 N-terminal amino acids from Pex13p (aa 151 to 386) abolished its interaction with Pex7p. In accordance with this observation, all Pex13p fragments that harbored even larger N-terminal truncations also tested negative for Pex7p binding (Fig. 1). Thus, the N-terminal domain of Pex13p, which is exposed to the cytosol (16), is the only binding site for Pex7p. This novel protein interaction module was also tested for interaction with Pex5p and Pex14p, the other known binding partners of Pex13p. However, these proteins failed to bind aa 1 to 151 of Pex13p in a two-hybrid assay (Fig. 2A).

Pex18p, Pex21p, and Pex7p interact with the same region in Pex13p. Translocation of proteins with a PTS2 targeting signal specifically requires, in addition to Pex7p, the redundant, largely cytosolic Pex18p and Pex21p (34). In the absence of both proteins, PTS2-dependent import is abolished. Both proteins can interact with Pex7p and are thought to target cargo-loaded Pex7p to the peroxisomal membrane (34). We therefore tested whether Pex18p and Pex21p are able to contact the Pex7p interaction domain of Pex13p. When fused to the Gal4p DNA-binding domain, both proteins exhibited autoactivity in the two-hybrid assay (data not shown). However, the respective activation domain fusions of Pex18p and Pex21p caused growth of the tester strain only when coexpressed with the amino terminus of Pex13p (Fig. 2B). This result indicated that Pex18p and Pex21p can also interact with the N terminus of Pex13p. To narrow down the binding regions in the Pex13p N terminus for the PTS2-specific proteins, smaller fragments of Pex13p were tested in the two-hybrid assay (Fig. 3A). Amino acids 1 to 100 proved even more efficient than aa 1 to 151 in binding the three proteins, while the amino-terminal 55 aa of Pex13p were efficiently recognized only by Pex7p. The con-




FIG. 1. Identification of the Pex7p-binding region in Pex13p. (A) Known features of *PEX13*. The cytosolically exposed SH3 domain (checked box) of the *S. cerevisiae* peroxisomal membrane protein Pex13p (386 aa) is bound by Pex5p and Pex14p at distinct sites as indicated by the arrows. The two predicted TMDs are marked in dark gray. (B) Analysis of *PEX13* fragments for interaction with *PEX7* in the yeast two-hybrid assay. The indicated regions of *PEX13* were fused to the *GAL4* activation domain of plasmid pPC86 and cotransformed with a pPC97-derived plasmid expressing a *GAL4* DNA-binding domain-*PEX7* fusion into the wild-type strain HF7c. The resulting prototrophs on histidine dropout plates containing 5 mM aminotriazole were considered positive for interaction (+). Numbers flanking the bars indicate the Pex13p amino acid residues that delimit the fragments.

struct comprising aa 55 to 100 of Pex13p failed to give rise to a positive signal with any of the proteins analyzed (Fig. 3A).

Pex13p₁₋₁₀₀ and Pex13p₁₋₅₅ interacted more strongly with Pex7p than with Pex18p or Pex21p, as the Pex7p-containing strains grew faster on histidine dropout plates than those containing Pex18p and Pex21p (data not shown). This result was also verified by quantification of the two-hybrid data by means of a β -galactosidase assay in the strain background of PJ69-4A, which harbors a sensitive *GAL7* promoter-*lacZ* reporter gene. Activities were about threefold higher in the strain expressing Pex7p-Pex13p₁₋₁₀₀ ($2,509 \pm 123$ U) than in those expressing Pex7p-Pex13p₁₋₅₅ (811 ± 137 U), whereas in the strain harboring Pex7p and Pex13p₅₅₋₁₀₀, activities reached only background levels (<50 U). None of the Pex18p/Pex21p-containing strains turned out to yield activities above the detection limit of the assay (<50 U). These data were interpreted to mean that despite the N-terminal 55 aa of Pex13p being essential for the Pex7p interaction, a somewhat larger region of Pex13p, possi-




FIG. 2. Only the PTS2-specific peroxins interact with the N terminus of Pex13p. (A) Analysis of Pex5p and Pex14p for their interaction with Pex13p₁₋₁₅₁. The interaction of Pex7p with Pex13p₁₋₁₅₁ in a two-hybrid assay is indicated by the histidine prototrophy of two independent clones of the respective transformant. The corresponding *PEX5* and *PEX14* transformants failed to grow on the same plate. (B) Pex18p and Pex21p interaction with Pex13p₁₋₁₅₁. Coexpression of *PEX18* and *PEX21* fusions to the *GAL4* activation domain (AD) with the Pex13p₁₋₁₅₁ *GAL4* DNA-binding domain (BD) fusion resulted in the formation of histidine prototrophs.

bly up to its first 100 aa, constitutes the full interaction site for Pex7p. The association of Pex18p/Pex21p with this optimal site could be explained by the assumption of a Pex7p-mediated interaction of Pex18p/Pex21p with Pex13p.

Pex18p and Pex21p bind Pex13p via Pex7p. The resemblance in binding sites opened the possibility that Pex18p and Pex21p bound to Pex13p in a complex with Pex7p. This was addressed by testing the interaction of Pex7p with Pex13p₁₋₁₀₀ in a strain devoid of both Pex18p and Pex21p (Fig. 3B). The interaction was unaffected, suggesting that Pex7p forms contacts with Pex13p that are independent of Pex18p/Pex21p. In contrast, the presence of Pex7p was important for the Pex18p/Pex21p interaction with Pex13p₁₋₁₀₀ (Fig. 3B), thereby supporting the notion that Pex7p serves a bridging protein for the interaction of Pex18p/Pex21p with Pex13p.

We then analyzed the possibility that Pex18p and Pex21p are required for the binding of Pex7p to the other docking protein, Pex14p, but this binding event occurred in the *pex18Δ pex21Δ* mutant background (Fig. 3C). Pex18p/Pex21p interacted only




FIG. 3. Pex18p and Pex21p contact Pex13p and Pex14p only in the presence of Pex7p. (A) Confinement of the Pex13p-binding region for Pex7p, Pex18p, and Pex21p. The indicated fragments of Pex13p were tested in a two-hybrid assay for interaction with Pex7p, Pex18p, and Pex21p in the HF7c wild-type strain. (B) Dependence of the Pex13p₁₋₁₀₀ interactions with Pex7p, Pex18p, and Pex21p on endogenous *PEX7* and *PEX18/PEX21*. The assays with Pex13p₁₋₁₀₀ were repeated in the otherwise isogenic HF7c strains *pex7Δ* and *pex18Δ pex21Δ*. Histidine prototrophs were not observed for Pex18p/Pex21p in the absence of Pex7p. (C) Dependence of the Pex7p-Pex14p two-hybrid interaction on endogenous *PEX18/PEX21*. The respective Pex7p and Pex14p fusions were transformed into HF7c and the corresponding *pex18Δ pex21Δ* strain. In both strains, coexpression of Pex7p and Pex14p caused growth on histidine dropout plates.

very weakly with Pex14p, and this interaction depended on Pex7p (data not shown), suggesting that these redundant factors do not contact Pex14p directly, either.

Pex7p docking in the absence of Pex18p and Pex21p. To obtain independent evidence for our results, we set out coimmunoprecipitation experiments with a *myc*-tagged version of Pex7p. The same protein has previously been used to demonstrate Pex7p binding to Pex13p in the absence of Pex5p and Pex14p (16). The tagged protein was expressed in wild-type and *pex18Δ* and *pex18Δ pex21Δ* mutant backgrounds and also in the docking protein mutants *pex13Δ* and *pex14Δ*. Expression levels of *myc*-Pex7p were comparable in all strains (Fig. 4). When *myc*-Pex7p was immunoprecipitated from whole-cell extracts of the wild-type strain, Pex13p and Pex14p were also




FIG. 4. Docking of Pex7p in the absence of Pex18p and Pex21p. The wild-type strain UTL-7A as well as the indicated, otherwise isogenic set of *myc*-Pex7p-expressing strains was induced in oleic acid-containing medium and analyzed for the presence of Pex13p and Pex14p in the immunoprecipitate of *myc*-Pex7p (upper panel). The upper and lower panels show 10% of the immunoprecipitated samples and 0.5% of the cell lysates, respectively, that had been subjected to immunoblotting with antisera against the *myc* epitope, Pex13p, and Pex14p.

found in the precipitate (Fig. 4). As anticipated, in the absence of Pex14p, Pex13p was still found in a complex with Pex7p. Since this was also the case in the *pex18Δ pex21Δ* double mutant (Fig. 4), this result demonstrated that binding of Pex7p to the docking complex does not require the presence of Pex18p and Pex21p.

Pex7p binds to Pex14p in vitro. The ability of Pex7p to bind Pex14p in the absence of Pex18p/Pex21p was also analyzed with bacterially expressed proteins. A His₆-tagged version of Pex14p was purified on Ni-NTA affinity columns onto which soluble extracts were loaded that contained either MBP or an MBP-Pex7p fusion protein. After washing, His₆-Pex14p was eluted from the columns and the eluates were analyzed for the presence of coeluted MBP-Pex7p. Coomassie brilliant blue staining revealed that Pex7p coeluted with His₆-Pex14p (Fig. 5, upper panel, lane 3) whereas MBP did not (lane 4). To ensure that the indicated bands in the Coomassie blue-stained gels indeed represented the MBP fusion protein, the same samples were diluted 20-fold and subjected to immunoblot analysis with an antibody directed against MBP (Fig. 5, lower panel). MBP-Pex7p was clearly detectable (lane 3), whereas little MBP was detected (lane 4). Thus, heterologously expressed Pex7p specifically interacted with Pex14p, demonstrating that these two peroxins can bind each other in a direct manner.

Pex7p binds to Pex13p₁₋₁₀₀ in vitro. To analyze the Pex7p-Pex13p interaction in vitro, the first 100 aa of Pex13p were expressed as a GST fusion protein in *E. coli* and subsequently purified on GST affinity columns. Subsequently, yeast lysates of wild-type or *pex18Δ pex21Δ* mutant cells expressing *myc*-tagged Pex7p were added, and bound GST proteins were visualized by Coomassie blue staining (Fig. 6A, lower panel). *myc*-Pex7p was




FIG. 5. Direct binding of Pex7p to Pex14p. Bacterially expressed His₆-tagged Pex14p was purified on Ni-NTA affinity columns. Subsequently, soluble extracts containing equal amounts of MBP (lane 1) or MBP-Pex7p (lane 2) were loaded onto these columns. After being washed, the eluates of His₆-Pex14p were analyzed for retained MBP-Pex7p (lane 3) or MBP (lane 4) by Coomassie blue staining (upper panel). The same samples were diluted 20-fold and subjected to immunoblot analysis with anti-MBP antibodies (lower panel).

detected immunologically by using anti-*myc* antibodies (Fig. 6A, upper panel). The tagged Pex7p was specifically retained in the presence of Pex13p₁₋₁₀₀ even when expressed in cells devoid of Pex18p and Pex21p (lanes 2 and 3). When lysate from an untransformed wild-type strain was applied to the Pex13p₁₋₁₀₀-GST-loaded column, no signal was obtained (lane 1). Similarly, GST alone also failed to bind *myc*-Pex7p (lanes 4 and 5). Thus, it could be inferred that the amino-terminal 100 aa of Pex13p are indeed the binding site for Pex7p and that Pex18p/Pex21p are not required for Pex7p binding to Pex13p₁₋₁₀₀.

To exclude the possibility that another yeast protein contributed to the observed binding of Pex7p to Pex13p, both proteins were bacterially expressed and subjected to in vitro binding studies. When expressed as a His₆-tagged version, a large fraction of Pex7p was found in inclusion bodies. Nevertheless, we could obtain a small fraction of His₆-Pex7p that was soluble. This fraction was loaded onto columns containing purified GST-Pex13p₁₋₁₀₀ or GST (Fig. 6B). Anti-Pex7p antibodies were then used to detect bound Pex7p by immunoblot analysis. His₆-Pex7p was only retained in the presence of Pex13p₁₋₁₀₀




FIG. 6. Direct binding of Pex7p to Pex13p₁₋₁₀₀. (A) Yeast *myc*-Pex7p specifically interacts with GST-Pex13p₁₋₁₀₀. Equal amounts of GST or GST-Pex13p₁₋₁₀₀ were loaded onto affinity columns and incubated with total cell lysates of a wild-type strain or the *myc*-Pex7p-expressing wild-type or *pex18* Δ *pex21* Δ strains. After the columns were washed, bound proteins were eluted and analyzed by immunoblotting for the presence of *myc*-Pex7p (upper panel). The presence of GST and GST-Pex13p₁₋₁₀₀ in these eluates was determined by Coomassie blue staining (lower panel). (B) Binding of Pex7p to Pex13p in vitro. Equal amounts of GST or GST-Pex13p₁₋₁₀₀ were loaded onto affinity columns and incubated with soluble extracts of bacterially expressed His₆-tagged Pex7p (+) or a mock-transformed *E. coli* strain (-). The retention of His₆-Pex7p was analyzed by immunoblotting (upper panel). The lower panel shows the presence of GST and GST-Pex13p₁₋₁₀₀ in these eluates.

(Fig. 6B, lane 3), thereby demonstrating that Pex7p can directly bind to the N-terminal region of Pex13p.

The N terminus of Pex13p is required for PTS2-dependent import. To elucidate the physiological relevance of the Pex7p-binding region in Pex13p, full-length Pex13p as well as a truncated variant of Pex13p lacking the N-terminal 55 aa were expressed from the native *PEX13* promoter in a *pex13* Δ strain and analyzed for their capability to complement the mutant phenotype. Immunoblot analysis demonstrated that truncated Pex13p₅₆₋₃₈₆ was stably expressed, although to a lower degree than full-length Pex13p (Fig. 7A). However, in contrast to full-length Pex13p, Pex13p₅₆₋₃₈₆ was not able to restore the




FIG. 7. The Pex7p-binding region of Pex13p is important for its function. (A) Expression of a truncated Pex13p variant in *pex13* Δ cells. Whole-cell extracts of strain UTL-7*Apex13* Δ that had expressed plasmid-borne copies of either full-length Pex13p (Pex13p₁₋₃₈₆) or an N-terminally truncated version of Pex13p (Pex13p₅₆₋₃₈₆) were subjected to immunoblot analysis with antiserum against Pex13p. The asterisk denotes proteins that had cross-reacted with the antiserum. (B) Complementation analysis of *pex13* Δ with truncated Pex13p₅₆₋₃₈₆. The indicated strains were spotted as a series of 10-fold dilutions on an oleic acid plate and incubated for 5 days at 30°C.

growth defect of a *pex13* Δ mutant on fatty acids as a sole carbon source (Fig. 7B), indicating that the N-terminal region is important for the function of Pex13p.

To test whether truncated Pex13p was correctly targeted to peroxisomes, GFP was fused to the C terminus of full-length Pex13p (Pex13p₁₋₃₈₆) and to Pex13p that lacked the entire N-terminal cytosolic domain (Pex13p₁₅₁₋₃₈₆). The GFP fusions of both the truncated and the full-length Pex13p were expressed in a wild-type strain, and their intracellular distribution was analyzed by fluorescence microscopy. Both Pex13p fusion proteins showed a punctate staining pattern (Fig. 8A) that was reminiscent of clustered peroxisomes. This clustering did not affect peroxisomal function, as full-length Pex13p-GFP was able to fully complement a *pex13* Δ strain (data not shown). Moreover, the staining patterns of both the full-length and the truncated Pex13p coincided with that of the peroxisomal marker protein PTS2-DsRed (Fig. 8A and B). It was therefore concluded that Pex13p₁₋₃₈₆ and Pex13p₁₅₁₋₃₈₆ are correctly targeted to the peroxisomal membrane. We then investigated whether truncated Pex13p₅₆₋₃₈₆ is still functional in peroxisomal matrix protein import. This was achieved by monitor-




FIG. 8. Only PTS2-dependent import is blocked in the absence of the Pex13p-docking site for Pex7p. (A) Peroxisomal localization of Pex13p₁₅₁₋₃₈₆. The UTL-7A strains expressing PTS2-DsRed as well as Pex13p₁₋₃₈₆-GFP or Pex13p₁₅₁₋₃₈₆-GFP were examined for GFP (left panels) and DsRed (middle panels) fluorescence. Bright-field microscopy (BF) demonstrates the structural integrity of the cells. (B) Localization of the fluorescing PTS2-DsRed marker protein. The wild-type UTL-7A and the otherwise isogenic *pex7*Δ and *pex18*Δ *pex21*Δ strains expressing PTS2-DsRed were streaked on oleic acid plates. After 2 days, cells were examined for DsRed fluorescence. The expression of a PTS-less DsRed in a wild-type strain was also monitored. (C) Selective complementation of PTS1-dependent import by Pex13p₅₆₋₃₈₆. The *pex13*Δ strains coexpressing Pex13p₁₋₃₈₆ or Pex13p₅₆₋₃₈₆ in combination with GFP-PTS1 (pJR233) and PTS2-DsRed were streaked on oleic acid plates and examined after 2 days for staining patterns of the fluorescing PTS marker proteins.

ing the subcellular distribution of fluorescent marker proteins for PTS1 and PTS2 import. GFP-SKL was used for analyzing PTS1-dependent import, while PTS2-DsRed, composed of the 16 N-terminal amino acids of Fox3p and the red fluorescent protein DsRed, served as the PTS2 marker protein. This protein was imported into peroxisomes in a wild-type strain but not in a *pex7Δ* or *pex18Δ pex21Δ* strain, and DsRed without the PTS2 signal remained cytosolic in a wild-type strain (Fig. 8B). As expected, both GFP-SKL and PTS2-DsRed caused diffuse staining when expressed in the *pex13Δ* mutant, which is defective in both import pathways (Fig. 8C). Coexpression of full-length Pex13p restored matrix protein import, since both marker proteins led to a punctate fluorescence staining pattern. Most interestingly, upon coexpression of the truncated Pex13p₅₆₋₃₈₆, PTS2-DsRed fluorescence remained diffuse, whereas GFP-SKL gave rise to a punctate staining pattern (Fig. 8C). These results clearly demonstrate that in the absence of the N terminus of Pex13p, PTS1-dependent import can still proceed, while PTS2-dependent import is blocked, probably because the binding of Pex7p to this N-terminal domain is abolished.

Pex7p-dependent complex formation of Pex18p/Pex21p and Fox3p. To determine where in the translocation process Pex18p/Pex21p are required, we reinvestigated the two-hybrid interactions of the PTS2 protein Fox3p with the PTS2-specific peroxins. Interaction of Fox3p with Pex7p was observed in wild-type, *pex7Δ*, and *pex18Δ pex21Δ* strains (Fig. 9A, lower panel). On the other hand, the interaction between Fox3p and Pex18p or Pex21p (data not shown) was strictly dependent on Pex7p (Fig. 9A, upper panel), which was in accordance with previous findings (34). This result demonstrated that Pex18p/Pex21p are also not required for the Pex7p-PTS2 interaction. To exclude an involvement of other yeast proteins in the PTS2 recognition event, an *in vitro* binding assay using bacterially expressed MBP-Pex7p was set out. MBP-Pex7p was bound to an amylose resin onto which soluble extracts from *E. coli* cells were loaded that had expressed either a synthetic PTS2 protein fused to GST or GST alone. Subsequent analysis of the eluted MBP-Pex7p fractions revealed that the targeting signal-containing protein was specifically retained (Fig. 9B). Thus, Pex7p alone was sufficient to recognize the PTS2 targeting signal.

The apparent requirement for Pex7p in the formation of a complex between Pex18p and Fox3p was analyzed by coimmunoprecipitation. For that matter, Pex18p was chromosomally tagged with the TAP tag at its C terminus in *pex21Δ* and *pex7Δ* strains (Fig. 9C, lower panel). Expression of the fusion protein in a *pex21Δ* strain background revealed that Pex18p-TAP was fully functional, as this strain grew comparably to the wild-type strain on oleic acid plates (data not shown). This fusion protein could be specifically precipitated from whole-cell extracts via its TAP tag by using IgG Sepharose (Fig. 9C, upper panel). The Pex18p-TAP precipitates were released from the column by using TEV protease and analyzed for the presence of Pex7p, which could be detected in the precipitate of the *pex21Δ* strain but not in those of the *pex7Δ* or untransformed wild-type strains (Fig. 9C, upper panel). The same precipitates were also analyzed for the presence of Fox3p. While the PTS2 cargo protein was found in the *pex21Δ* sample, it was absent from the

precipitate derived from the *pex7Δ* strain (Fig. 9C, upper panel). It was therefore concluded that Fox3p can indeed associate with Pex18p *in vivo*, albeit only in the presence of Pex7p.

Pex7p- and Pex18p/Pex21p-dependent docking of Fox3p to Pex14p. We then analyzed whether Fox3p would be found in a complex with the docking proteins Pex14p and Pex13p when either Pex7p or Pex18p/Pex21p were absent. This was achieved by testing for an interaction of Fox3p with Pex14p and Pex13p in a two-hybrid assay. Remarkably, a strong interaction between Fox3p and Pex14p was observed in a wild-type strain, which was absent in a *pex7Δ* or *pex18Δ pex21Δ* mutant background (Fig. 10A). On the other hand, no interaction between Fox3p and Pex13p₁₋₁₀₀, a Pex13p fragment that strongly interacted with Pex7p, was detectable in a wild-type strain or a *pex18Δ pex21Δ* strain (Fig. 10B). Even in a *pex14Δ* strain, where Pex7p binding to Pex13p is likely to increase, no interaction between Fox3p and Pex13p was detected (Fig. 10B). These findings indicate that docking of the PTS2 cargo protein is critically dependent on both Pex18p/Pex21p and Pex7p. They further suggest that Pex14p represents the docking protein, whereas Pex13p is involved in the binding of the Pex7p-Pex18p/Pex21p complex without cargo protein.

Pex18p and Pex21p promote the formation of an import-competent PTS2 substrate complex. In the absence of a functional docking or translocation machinery, immunoprecipitates of *myc*-Pex7p contain a drastically increased amount of the PTS2 protein Fox3p (16, 37). This increased binding was interpreted to be due to the cytosolic accumulation of Fox3p in these mutants. Were Pex18p/Pex21p only to mediate the docking of Fox3p to Pex14p, then such an apparent accumulation of cargo-loaded receptor should also occur in a *pex18Δ pex21Δ* strain. As expected, the *myc*-Pex7p immunoprecipitates of a wild-type strain contained little Fox3p, whereas those of *pex13Δ* and *pex14Δ* strains showed an accumulation of Fox3p (Fig. 11A). However, in the *pex18Δ pex21Δ* strain, only very little Fox3p could be coprecipitated (Fig. 11A).

Since Fox3p also did not accumulate in a wild-type strain, we analyzed the possibility that in our experiment, ectopic expression of *myc*-Pex7p suppressed the defect of the *pex18Δ pex21Δ* strain. In that case, growth of the *pex18Δ pex21Δ* mutant on oleic acid plates should be restored when it is transformed with the *myc*-Pex7p expression plasmid. However, as can be seen in Fig. 11B, this strain (*pex18Δ pex21Δ* [*myc*-Pex7p]) remained deficient in utilizing oleic acid to a degree similar to that of the untransformed strain (*pex18Δ pex21Δ*).

Thus, Pex18p/Pex21p probably act in a step that precedes docking of Pex7p to the membrane. In that case, the observed increase of Fox3p bound to *myc*-Pex7p in a *pex14Δ* mutant would be abolished by additionally deleting *PEX18* and *PEX21*. The amount of coimmunoprecipitated Fox3p in a *pex14Δ pex18Δ pex21Δ* triple-mutant strain was therefore compared to those of the *pex14Δ*, *pex18Δ pex21Δ*, and wild-type strains. Strikingly, an accumulation of Fox3p was not observed in the triple-deletion strain (Fig. 11C). This result indicates that the accumulation of the Pex7p-Fox3p complex in a *pex14Δ* mutant is dependent on Pex18p/Pex21p and, as a consequence, that the function of Pex18p and Pex21p is already required in the import process prior to Pex14p and thus before the Fox3p-Pex7p-Pex18p/Pex21p complex docks at the peroxisomal membrane.






FIG. 9. Pex18p binds Fox3p only via Pex7p. (A) Fox3p was tested in a two-hybrid assay for interaction with Pex18p (upper panel) and Pex7p (lower panel) in the HF7c wild type as well as the *pex18Δpex21Δ* and *pex7Δ* strains. (B) Binding of Pex7p to a synthetic PTS2 protein in vitro. Equal amounts of MBP-Pex7p were loaded onto affinity columns and incubated with soluble extracts of bacterially expressed GST-PTS2-GFP-prA (GST-PTS2; right lane in each panel) or GST (left lane). The right panel shows a Coomassie blue stain of the loaded fractions. The retention of GST-containing proteins was analyzed by immunoblotting (upper-left panel). The lower panel shows the




FIG. 10. Fox3p interacts with Pex14p in a Pex7p- and Pex18p/Pex21p-dependent manner. The docking proteins Pex14p (A) and Pex13p (B) were tested for interaction with Fox3p in a two-hybrid assay in the indicated strain backgrounds.

DISCUSSION

Despite the identification of probably all components of the peroxisomal matrix protein import machinery in recent years (40, 47), the mechanism underlying the translocation process is still largely unknown. In this report, we have shown that *S. cerevisiae* Pex13p plays a direct role in PTS2-dependent peroxisomal protein import and begun to unravel the function of the PTS2-specific peroxins in the early steps of this process.

After their synthesis, PTS2 proteins such as thiolase are specifically recognized by Pex7p (36, 56). The recognition of PTS2 by Pex7p does not require *S. cerevisiae* Pex18p/Pex21p, whereas the latter proteins interact with Fox3p only in the presence of Pex7p (34). Our demonstration of yeast Pex7p binding a synthetic PTS2 protein in vitro goes beyond that of a previous report, where a Pex7p was used that had been immunoprecipitated from wild-type yeast extracts and thus might have contained other yeast proteins (36). Our first attempts to study in vitro the influence of Pex18p on the Pex7p-PTS2 interaction failed because bacterially expressed Pex18p did not bind Pex7p, even in the concomitant presence of PTS2 protein (data not shown). However, we could show by immunoprecipitation that such a ternary complex of Pex18p, Pex7p, and Fox3p indeed exists, and we substantiated the requirement for Pex7p in the formation of such a complex. Notwithstanding that, we found that the Fox3p-Pex7p-containing complex that accumulates in a *pex14Δ* strain vanished when *PEX18* and

presence of MBP-Pex7p in both eluates. (C) Coimmunoprecipitation of Fox3p with Pex18p. Strains *pex7Δ PEX18-TAP* (yAS3), *pex21Δ PEX18-TAP* (yAS4), and the untransformed UTL7-A wild type were induced in oleic acid-containing medium and analyzed for the presence of Pex7p and Fox3p in the precipitate of Pex18p-TAP by immunoblotting (upper panel). The upper and lower panels show 33% of the immunoprecipitated samples and 0.2% of the total cell lysates, respectively.




FIG. 11. The Pex7p-Fox3p-containing cytosolic complex does not accumulate in a *pex18* *pex21*Δ mutant. (A) Accumulation of a Fox3p-Pex7p complex in peroxin mutants. The indicated *myc*-Pex7p-expressing strains were analyzed for the presence of Fox3p in the immunoprecipitate of *myc*-Pex7p (10% of total) by immunoblotting (upper panel). The lower panel shows an immunoblot of the cell lysates (0.5% of total) that were used for precipitation. (B) The effect of overexpressing Pex7p in a *pex18* *pex21*Δ strain. Tenfold dilutions of the indicated strains were spotted on oleic acid-containing plates and incubated for 5 days at 30°C. (C) Epistatic analysis of *PEX14* and *PEX18/PEX21*. Precipitation of *myc*-Pex7p from the indicated oleic acid-induced cells and subsequent analysis of precipitated Fox3p were performed as described for panel A.

PEX21 were additionally deleted. *S. cerevisiae* thiolase is assembled into its active dimeric form in the cytosol even in the absence of its PTS2 sequence (17), arguing against an essential role for Pex18p/Pex21p in the assembly of enzymatically active Fox3p. Rather, these proteins are required to form an import-competent Fox3p complex.

A similar function in thiolase import was proposed for *Y. lipolytica* Pex20p. This protein binds thiolase autonomously, in a PTS2 targeting signal-independent fashion. In fact, Pex20p forms hetero-oligomers with Fox3p, an event that is probably mandatory for the generation of an import-competent complex (49). In analogy to Pex20p, Pex18p/Pex21p could physically contact Fox3p in the Pex7p-bound conformation. In that case,

the targeting signal would be bound by Pex7p, while Pex18p/Pex21p would contact both Pex7p, via their conserved motif (7), and Fox3p, which would then trigger higher-order oligomerization. Alternatively, Pex18p/Pex21p could assist in forming a Fox3p-Pex7p-containing complex by stabilizing the binding between Pex7p and Fox3p, although the existing two-hybrid data do not support this idea. Still, both cases would explain why Pex20p was able to partially complement a *pex18* *pex21*Δ mutant strain (8). Oligomerization as a requirement for import is not just an idiosyncrasy of thiolase. Such a postulate was already raised upon investigating the import of *Candida boidinii* alcohol oxidase, which seemed to depend on the oligomerization or aggregation at the cytoplasmic side of the

peroxisomal membrane (4). Since then, the ability of peroxisomes to import folded or even oligomeric proteins has been amply recorded (28, 52).

Our data also suggest that a complex is formed that contemporaneously contains membrane-associated Pex14p, Pex7p, Pex18p/Pex21p, and Fox3p (Fig. 10A). In the absence of either Pex7p or Pex18p/Pex21p, Fox3p was no longer able to interact with Pex14p. On the other hand, Fox3p failed to interact with the Pex7p-binding domain of Pex13p even in a wild-type strain (Fig. 10B). It is unlikely that the lack of interaction was caused by a masking of the interaction domain in Pex13p with the Gal4p moieties, as the same fusion construct was very efficient in recognizing Pex7p and Pex18p/Pex21p, which in turn were able to interact with the Fox3p fusion proteins. Thus, we believe that our data point to a scenario where Pex14p is the initial docking site in PTS2-dependent protein import. We also considered the possibility of Pex13p constituting the initial docking site for the empty Pex5p and Pex7p, which would only then bind their cargo proteins. However, this is hard to reconcile with the observed formation of a Pex7p-Pex18p/Pex21p-Fox3p complex in the *pex13Δ* mutant, where this complex must have formed in the cytosol. Future biochemical work will have to demonstrate whether cargo-loaded PTS2 receptor can indeed be found in a complex with Pex14p but not with Pex13p. In mammalian cells, the Pex7p-Pex5pL-PTS2 substrate complex also likely contacts Pex14p prior to Pex13p (30). It is worth noting that a similar conclusion was drawn for the PTS1 receptor Pex5p of *Pichia pastoris*, since substrate-loaded Pex5p bound Pex14p with higher affinity than Pex5p alone, whereas the opposite was true for Pex13p (50).

In contrast to Fox3p, Pex7p does not require the presence of Pex18p/Pex21p to contact Pex14p (Fig. 3C). Our *in vitro* binding analysis showed that this binding event can even occur in the absence of any other yeast protein. Pex7p also contacts Pex13p directly (Fig. 6), thereby ruling out the possibility of Pex18p/Pex21p being adapter proteins between Pex7p and the docking complex. Notably, *Y. lipolytica* Pex20p did bind *S. cerevisiae* Pex13p and Pex14p in the absence of Pex7p (8). Thus, in *Y. lipolytica* the function of a putative Pex7p might be restricted to targeting signal recognition, whereas Pex20p might be required to form an import-competent complex as well as to contact the docking machinery. Alternatively, Pex20p might contain the function of both Pex18p/Pex21p and Pex7p. In mammals, the Pex18p-like factor Pex5pL is able to interact with Pex14p via its diatomic pentapeptide repeats (13, 41, 54). It is therefore imaginable that Pex5pL mediates the docking of Pex7p to Pex14p. However, Shimizu and colleagues demonstrated direct binding of Pex7p to Pex14p (45). In this case, it is also possible that Pex5pL, in analogy to Pex18p, is involved in the formation of a PTS2 cargo import-competent complex.

In light of our results, the role of Pex13p in PTS2 protein import could be substantiated. Deletion of the N-terminal domain of Pex13p, which specifically interacted with Pex7p, affected PTS2-dependent import but left PTS1 protein import intact. The latter observation indicated that Pex14p was correctly targeted to the peroxisomal membrane. As a consequence, the role of Pex13p in PTS2 import is not restricted to the targeting of Pex14p. Rather, Pex13p fulfills an essential step in PTS2 import that is distinct from that governed by

Pex14p. This result also meant that although the import of both PTS1 and PTS2 requires Pex13p, the two pathways utilize different regions within Pex13p and therefore do not coincide at the stage of Pex13p in a more narrow sense. Mutations that specifically block one of the two routes have also been found previously for Pex8p and Pex2p, two peroxins that are possibly involved in later steps of the import cascade (22, 25).

The steps following docking are less clear. Peroxisomal import deviates considerably from the well-established import into mitochondria or the endoplasmic reticulum (43). An obvious protein translocation channel is lacking, and the ability to import folded or even oligomeric proteins must also be taken into account. Interestingly, Pex18p and Pex21p are constantly degraded due to ubiquitination, which is apparently restricted to the fraction of peroxisomally localized Pex18p/Pex21p (33). Although it remains to be determined whether the observed degradation of Pex18p/Pex21p is physiologically relevant for PTS2-dependent protein import, it could be that Pex18p/Pex21p are required not only for the formation of an import-competent PTS2 cargo protein complex but also for a step that follows docking.

ACKNOWLEDGMENTS

We are grateful to M. Nündel for excellent technical assistance. We thank X. Hong for plasmids His₆-PEX14 and His₆-PEX7, R. Bahadori for plasmid pRB107, A. Hartig for plasmid pJR233, B. Seraphin for plasmid pBS1479, H. Otto for anti-*myc* antibodies, and particularly W. H. Kunau for the provision of plasmids, strains, and antibodies.

This work was supported by the Deutsche Forschungsgemeinschaft, grants ER178/2-3 and SFB449, and by the Fonds der Deutschen Chem. Industrie. H.R. was supported by an EMBO long-term fellowship (ALTF255-2000).

REFERENCES

- Albertini, M., P. Rehling, R. Erdmann, W. Girzalsky, J. A. Kiel, M. Veenhuis, and W. H. Kunau. 1997. Pex14p, a peroxisomal membrane protein binding both receptors of the two PTS-dependent import pathways. *Cell* **89**:83–92.
- Barnett, P., G. Bottger, A. T. Klein, H. F. Tabak, and B. Distel. 2000. The peroxisomal membrane protein Pex13p shows a novel mode of SH3 interaction. *EMBO J.* **19**:6382–6391.
- Baumgartner, U., B. Hamilton, M. Piskacek, H. Ruis, and H. Rottensteiner. 1999. Functional analysis of the Zn(2)/Cys(6) transcription factors Oaf1p and Pip2p: different roles in fatty acid induction of beta-oxidation in *Saccharomyces cerevisiae*. *J. Biol. Chem.* **274**:22208–22216.
- Bellion, E., and J. M. Goodman. 1987. Proton ionophores prevent assembly of a peroxisomal protein. *Cell* **48**:165–173.
- Braverman, N., G. Dodt, S. J. Gould, and D. Valle. 1998. An isoform of Pex5p, the human PTS1 receptor, is required for the import of PTS2 proteins into peroxisomes. *Hum. Mol. Genet.* **7**:1195–1205.
- Chevray, P. M., and D. Nathans. 1992. Protein interaction cloning in yeast: identification of mammalian proteins that react with the leucine zipper of Jun. *Proc. Natl. Acad. Sci. USA* **89**:5789–5793.
- Dodt, G., D. Warren, E. Becker, P. Rehling, and S. J. Gould. 2001. Domain mapping of human PEX5 reveals functional and structural similarities to *Saccharomyces cerevisiae* Pex18p and Pex21p. *J. Biol. Chem.* **276**:41769–41781.
- Einwächter, H., S. Sowinski, W. H. Kunau, and W. Schliebs. 2001. *Yarrowia lipolytica* Pex20p, *Saccharomyces cerevisiae* Pex18p/Pex21p and mammalian Pex5p fulfil a common function in the early steps of the peroxisomal PTS2 import pathway. *EMBO Rep.* **2**:1035–1039.
- Erdmann, R., and W. H. Kunau. 1994. Purification and immunolocalization of the peroxisomal 3-oxoacyl-CoA thiolase from *Saccharomyces cerevisiae*. *Yeast* **10**:1173–1182.
- Erdmann, R., M. Veenhuis, D. Mertens, and W. H. Kunau. 1989. Isolation of peroxisome-deficient mutants of *Saccharomyces cerevisiae*. *Proc. Natl. Acad. Sci. USA* **86**:5419–5423.
- Feilotter, H. E., G. J. Hannon, C. J. Ruddell, and D. Beach. 1994. Construction of an improved host strain for two hybrid screening. *Nucleic Acids Res.* **22**:1502–1503.
- Fields, S., and O. Song. 1989. A novel genetic system to detect protein-protein interactions. *Nature* **340**:245–246.

13. Fransen, M., S. R. Terlecky, and S. Subramani. 1998. Identification of a human PTS1 receptor docking protein directly required for peroxisomal protein import. *Proc. Natl. Acad. Sci. USA* **95**:8087–8092.
14. Fujiki, Y. 2000. Peroxisome biogenesis and peroxisome biogenesis disorders. *FEBS Lett.* **476**:42–46.
15. Gietz, R. D., and A. Sugino. 1988. New yeast-*Escherichia coli* shuttle vectors constructed with in vitro mutagenized yeast genes lacking six-base pair restriction sites. *Gene* **74**:527–534.
16. Girzalsky, W., P. Rehling, K. Stein, J. Kipper, L. Blank, W. H. Kunau, and R. Erdmann. 1999. Involvement of Pex13p in Pex14p localization and peroxisomal targeting signal 2-dependent protein import into peroxisomes. *J. Cell Biol.* **144**:1151–1162.
17. Glover, J. R., D. W. Andrews, and R. A. Rachubinski. 1994. *Saccharomyces cerevisiae* peroxisomal thiolase is imported as a dimer. *Proc. Natl. Acad. Sci. USA* **91**:10541–10545.
18. Gould, S. J., G. A. Keller, and S. Subramani. 1988. Identification of peroxisomal targeting signals located at the carboxy terminus of four peroxisomal proteins. *J. Cell Biol.* **107**:897–905.
19. Gould, S. J., and D. Valle. 2000. Peroxisome biogenesis disorders: genetics and cell biology. *Trends Genet.* **16**:340–345.
20. Gurvitz, A., H. Rottensteiner, S. H. Kilpelainen, A. Hartig, J. K. Hiltunen, M. Binder, I. W. Dawes, and B. Hamilton. 1997. The *Saccharomyces cerevisiae* peroxisomal 2,4-dienoyl-CoA reductase is encoded by the oleate-inducible gene *SPS19*. *J. Biol. Chem.* **272**:22140–22147.
21. Holroyd, C., and R. Erdmann. 2001. Protein translocation machineries of peroxisomes. *FEBS Lett.* **501**:6–10.
22. Huang, Y., R. Ito, S. Miura, T. Hashimoto, and M. Ito. 2000. A missense mutation in the RING finger motif of PEX2 protein disturbs the import of peroxisome targeting signal 1 (PTS1)-containing protein but not the PTS2-containing protein. *Biochem. Biophys. Res. Commun.* **270**:717–721.
23. Huhse, B., P. Rehling, M. Albertini, L. Blank, K. Meller, and W. H. Kunau. 1998. Pex17p of *Saccharomyces cerevisiae* is a novel peroxin and component of the peroxisomal protein translocation machinery. *J. Cell Biol.* **140**:49–60.
24. Lametschwandtner, G., C. Brocard, M. Fransen, P. Van Veldhoven, J. Berger, and A. Hartig. 1998. The difference in recognition of terminal tripeptides as peroxisomal targeting signal 1 between yeast and human is due to different affinities of their receptor Pex5p to the cognate signal and to residues adjacent to it. *J. Biol. Chem.* **273**:33635–33643.
25. Liu, H., X. Tan, K. A. Russell, M. Veenhuis, and J. M. Cregg. 1995. *PER3*, a gene required for peroxisome biogenesis in *Pichia pastoris*, encodes a peroxisomal membrane protein involved in protein import. *J. Biol. Chem.* **270**:10940–10951.
26. Marzioch, M., R. Erdmann, M. Veenhuis, and W. H. Kunau. 1994. *PAS7* encodes a novel yeast member of the WD-40 protein family essential for import of 3-oxoacyl-CoA thiolase, a PTS2-containing protein, into peroxisomes. *EMBO J.* **13**:4908–4918.
27. Matsumura, T., H. Otera, and Y. Fujiki. 2000. Disruption of the interaction of the longer isoform of Pex5p, Pex5pL, with Pex7p abolishes peroxisome targeting signal type 2 protein import in mammals. Study with a novel Pex5-impaired Chinese hamster ovary cell mutant. *J. Biol. Chem.* **275**:21715–21721.
28. McNew, J. A., and J. M. Goodman. 1994. An oligomeric protein is imported into peroxisomes in vivo. *J. Cell Biol.* **127**:1245–1257.
29. Mumberg, D., R. Muller, and M. Funk. 1994. Regulatable promoters of *Saccharomyces cerevisiae*: comparison of transcriptional activity and their use for heterologous expression. *Nucleic Acids Res.* **22**:5767–5768.
30. Otera, H., T. Harano, M. Honsho, K. Ghaedi, S. Mukai, A. Tanaka, A. Kawai, N. Shimizu, and Y. Fujiki. 2000. The mammalian peroxin Pex5pL, the longer isoform of the mobile peroxisome targeting signal (PTS) type 1 transporter, translocates the Pex7p-PTS2 protein complex into peroxisomes via its initial docking site, Pex14p. *J. Biol. Chem.* **275**:21703–21714.
31. Otera, H., K. Okumoto, K. Tateishi, Y. Ikoma, E. Matsuda, M. Nishimura, T. Tsukamoto, T. Osumi, K. Ohashi, O. Higuchi, and Y. Fujiki. 1998. Peroxisome targeting signal type 1 (PTS1) receptor is involved in import of both PTS1 and PTS2: studies with PEX5-defective CHO cell mutants. *Mol. Cell Biol.* **18**:388–399.
32. Purdue, P. E., and P. B. Lazarow. 2001. Peroxisome biogenesis. *Annu. Rev. Cell Dev. Biol.* **17**:701–752.
33. Purdue, P. E., and P. B. Lazarow. 2001. Pex18p is constitutively degraded during peroxisome biogenesis. *J. Biol. Chem.* **276**:47684–47689.
34. Purdue, P. E., X. Yang, and P. B. Lazarow. 1998. Pex18p and Pex21p, a novel pair of related peroxins essential for peroxisomal targeting by the PTS2 pathway. *J. Cell Biol.* **143**:1859–1869.
35. Reguenga, C., M. E. Oliveira, A. M. Gouveia, C. Sa-Miranda, and J. E. Azevedo. 2001. Characterization of the mammalian peroxisomal import machinery: Pex2p, Pex5p, Pex12p, and Pex14p are subunits of the same protein assembly. *J. Biol. Chem.* **276**:29935–29942.
36. Rehling, P., M. Marzioch, F. Niesen, E. Wittke, M. Veenhuis, and W. H. Kunau. 1996. The import receptor for the peroxisomal targeting signal 2 (PTS2) in *Saccharomyces cerevisiae* is encoded by the *PAS7* gene. *EMBO J.* **15**:2901–2913.
37. Rehling, P., A. Skaletz-Rorowski, W. Girzalsky, T. Voorn-Brouwer, M. M. Franse, B. Distel, M. Veenhuis, W. H. Kunau, and R. Erdmann. 2000. Pex8p, an intraperoxisomal peroxin of *Saccharomyces cerevisiae* required for protein transport into peroxisomes binds the PTS1 receptor Pex5p. *J. Biol. Chem.* **275**:3593–3602.
38. Rigaut, G., A. Shevchenko, B. Rutz, M. Wilm, M. Mann, and B. Seraphin. 1999. A generic protein purification method for protein complex characterization and proteome exploration. *Nat. Biotechnol.* **17**:1030–1032.
39. Rottensteiner, H., A. J. Kal, B. Hamilton, H. Ruis, and H. F. Tabak. 1997. A heterodimer of the Zn2Cys6 transcription factors Pip2p and Oaf1p controls induction of genes encoding peroxisomal proteins in *Saccharomyces cerevisiae*. *Eur. J. Biochem.* **247**:776–783.
40. Sacksteder, K. A., and S. J. Gould. 2000. The genetics of peroxisome biogenesis. *Annu. Rev. Genet.* **34**:623–652.
41. Saidowsky, J., G. Dodt, K. Kirchberg, A. Wegner, W. Nastainczyk, W. H. Kunau, and W. Schliebs. 2001. The di-aromatic pentapeptide repeats of the human peroxisome import receptor PEX5 are separate high affinity binding sites for the peroxisomal membrane protein PEX14. *J. Biol. Chem.* **276**:34524–34529.
42. Sambrook, J., E. F. Fritsch, and T. Maniatis. 1989. *Molecular cloning: a laboratory manual*. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.
43. Schatz, G., and B. Dobberstein. 1996. Common principles of protein translocation across membranes. *Science* **271**:1519–1526.
44. Schiestl, R. H., and R. D. Gietz. 1989. High efficiency transformation of intact yeast cells using single stranded nucleic acids as a carrier. *Curr. Genet.* **16**:339–346.
45. Shimizu, N., R. Itoh, Y. Hirono, H. Otera, K. Ghaedi, K. Tateishi, S. Tamura, K. Okumoto, T. Harano, S. Mukai, and Y. Fujiki. 1999. The peroxin Pex14p. cDNA cloning by functional complementation on a Chinese hamster ovary cell mutant, characterization, and functional analysis. *J. Biol. Chem.* **274**:12593–12604.
46. Smith, J. J., and R. A. Rachubinski. 2001. A role for the peroxin Pex8p in Pex20p-dependent thiolase import into peroxisomes of the yeast *Yarrowia lipolytica*. *J. Biol. Chem.* **276**:1618–1625.
47. Subramani, S., A. Koller, and W. B. Snyder. 2000. Import of peroxisomal matrix and membrane proteins. *Annu. Rev. Biochem.* **69**:399–418.
48. Swinkels, B. W., S. J. Gould, A. G. Bodnar, R. A. Rachubinski, and S. Subramani. 1991. A novel, cleavable peroxisomal targeting signal at the amino-terminus of the rat 3-ketoacyl-CoA thiolase. *EMBO J.* **10**:3255–3262.
49. Titorenko, V. I., J. J. Smith, R. K. Szilard, and R. A. Rachubinski. 1998. Pex20p of the yeast *Yarrowia lipolytica* is required for the oligomerization of thiolase in the cytosol and for its targeting to the peroxisome. *J. Cell Biol.* **142**:403–420.
50. Urquhart, A. J., D. Kennedy, S. J. Gould, and D. I. Crane. 2000. Interaction of Pex5p, the type 1 peroxisome targeting signal receptor, with the peroxisomal membrane proteins Pex14p and Pex13p. *J. Biol. Chem.* **275**:4127–4136.
51. van den Bosch, H., R. B. Schutgens, R. J. Wanders, and J. M. Tager. 1992. Biochemistry of peroxisomes. *Annu. Rev. Biochem.* **61**:157–197.
52. Walton, P. A., P. E. Hill, and S. Subramani. 1995. Import of stably folded proteins into peroxisomes. *Mol. Biol. Cell* **6**:675–683.
53. Westermann, B., and W. Neupert. 2000. Mitochondria-targeted green fluorescent proteins: convenient tools for the study of organelle biogenesis in *Saccharomyces cerevisiae*. *Yeast* **16**:1421–1427.
54. Will, G. K., M. Soukupova, X. Hong, K. S. Erdmann, J. A. Kiel, G. Dodt, W. H. Kunau, and R. Erdmann. 1999. Identification and characterization of the human orthologue of yeast Pex14p. *Mol. Cell Biol.* **19**:2265–2277.
55. Yaffe, M. P., and G. Schatz. 1984. Two nuclear mutations that block mitochondrial protein import in yeast. *Proc. Natl. Acad. Sci. USA* **81**:4819–4823.
56. Zhang, J. W., and P. B. Lazarow. 1996. Pex1p (Pas7p) is an intraperoxisomal receptor for the NH2-terminal, type 2, peroxisomal targeting sequence of thiolase: Pex1p itself is targeted to peroxisomes by an NH2-terminal peptide. *J. Cell Biol.* **132**:325–334.