

pipsqueak Encodes a Factor Essential for Sequence-Specific Targeting of a Polycomb Group Protein Complex

Der-Hwa Huang,* Yuh-Long Chang, Chih-Chao Yang, I-Ching Pan,† and Balas King

Institute of Molecular Biology, Academia Sinica, Taipei, Taiwan 115, Republic of China

Received 11 February 2002/Returned for modification 25 March 2002/Accepted 6 June 2002

The Polycomb (Pc) group (Pc-G) of repressors is essential for transcriptional silencing of homeotic genes that determine the axial development of metazoan animals. It is generally believed that the multimeric complexes formed by these proteins nucleate certain chromatin structures to silence promoter activity upon binding to Pc-G response elements (PRE). Little is known, however, about the molecular mechanism involved in sequence-specific binding of these complexes. Here, we show that an immunoprecipitated Pc protein complex contains a DNA binding activity specific to the (GA)_n motif in a PRE from the *bithoraxoid* region. We found that this activity can be attributed primarily to the large protein isoform encoded by *pipsqueak* (*psq*) instead of to the well-characterized GAGA factor. The functional relevance of *psq* to the silencing mechanism is strongly supported by its synergistic interactions with a subset of Pc-G that cause misexpression of homeotic genes.

The Hox genes in the Antennapedia complex (ANTP-C) and the bithorax complex (BX-C) of *Drosophila melanogaster* are essential for segmental specification along the anteroposterior body axis (32, 41). Aberrant expression of these genes often causes homeotic transformation, i.e., misspecification of body parts. Complex regulatory mechanisms exist to ensure silencing of these genes outside specific domains throughout development (42). The silenced state is initially established by segmentation genes, including *hunchback* and *tailless*, that are expressed transiently in early embryos (46). Subsequent maintenance requires the presence of large numbers of the Polycomb (Pc) group (Pc-G) repressors.

Studies of *cis*-acting sequences capable of maintaining the silenced state of homeotic genes have revealed that, although extended sequences are necessary for full function, short sequences of several hundreds of bases have been proven to be critical. These sequences are commonly referred to as Pc-G response elements (PRE) (53). PRE appear to silence a closely linked marker gene in an epigenetically transmittable manner during cell division (10).

Several salient features have been noted about PRE. For example, PRE can silence a distant marker gene (54). PRE can also exhibit a pairing-sensitive silencing effect, resulting in much stronger silencing on the marker gene when PRE is present on the homologous chromosome (31). A high incidence of PRE insertion occurs at sites that contain preexisting PRE or PRE-like sequences (19). In general, PRE insertion creates a new chromosomal binding site for many Pc-G proteins (17). Further, PRE can confer transcription repression on *Ultrabithorax* (*Ubx*) in a Pc-dependent manner in cultured cells (13). Thus, PRE appear to act as the core sequences upon

which Pc-G proteins assemble into large functional silencing complexes. It has been speculated that PRE at different chromosomal sites, when spatially juxtaposed, might cooperate and become more effective (48).

How Pc-G can accomplish these tasks remains largely unclear. To date, less than half a dozen Pc-G have been thoroughly studied. Some Pc-G proteins contain domains that are capable of homophilic or heterophilic interaction (38, 45), potentially facilitating formation and/or interaction of multimeric protein complexes. Consistently, large protein complexes containing Pc-G proteins have been identified. For example, PC, Polyhomeotic (PH), and Posterior Sex Combs (PSC) are found in the Pc repression complex 1 of approximately 2 MDa (51). A smaller protein complex containing Enhancer of Zeste [E(Z)] and Extra Sex Combs (ESC) has also been reported (44, 59). Since some Pc-G proteins have not been shown to copurify with these complexes, additional complexes might be expected. Germ line clones of many Pc-G mutations display similar but distinct patterns of embryonic defects, suggesting partially overlapping functions (56). Chromatin immunoprecipitation has also revealed substantial variation in the composition of the Pc-G complexes at different sites (58). Surprisingly, some of these sites are found in actively expressed genes (58). Thus, multiple Pc-G complexes might function in different contexts during development.

A statistical estimation has suggested that there are about 40 Pc-G in the fly genome (29). Consistent with this view, many novel proteins have been found in Pc-G complexes (14, 44, 50, 51, 59). Recently, mutations of several new genes were found to enhance homeotic phenotypes when in combination with Pc-G mutations; however, they produced little effect by themselves (14, 33, 56). These types of genes might define functions distinct from those of previously characterized Pc-G. Indeed, the identification of MI-2 and histone deacetylase 1 (HDAC1) as crucial partners has revealed that nucleosome modification, via histone deacetylation, plays a crucial role in homeotic gene silencing (14, 33, 59).

* Corresponding author. Mailing address: N-420, Institute of Molecular Biology, Academia Sinica, Taipei, Taiwan 115, Republic of China. Phone: (886) 2-2789-9219. Fax: (886) 2-2782-6085. E-mail: mbdhuang@ccvax.sinica.edu.tw.

† Present address: Department of Genetics, Duke University, Durham, NC 27710.

A fundamental question yet to be addressed fully is how the Pc-G protein complexes recognize specific sequences in PRE. With the exception of *pleiohomeotic* (*pho*), which encodes the homologue of mammalian YY1 (8), no existing Pc-G has been shown to bind specific DNA sequences. The PHO binding site is a functional constituent of PRE (20); however, the inability of a LexA-PHO fusion protein to silence a linked reporter gene as other Pc-G fusion proteins suggests that PHO alone may not be sufficient to target functional Pc-G complexes (49). Recently, the $(GA)_n$ motif present in PRE has been suggested to be critical for homeotic gene silencing (24, 25). It has been further suggested that the GAGA factor (GAF), a well-characterized DNA binding protein for such a motif (55), is involved in the binding (24, 25). Contrary to the expected silencing effect, GAF has also been shown to act either as an antirepressor to alleviate the negative effects of histone H1 (36) or as a transactivator *in vitro*, in cultured cells, and in stress response (6, 21, 55). In addition, *Trithorax-like* (*Trl*), the GAF-encoding gene, has been formerly classified as a member of the trithorax group of genes (*trx-G*) that antagonize Pc-G (18). Therefore, the role of GAF remains unresolved.

An ~440-bp DNA fragment from the *bithoraxoid* (*bxd*) region of *Ubx* that can recapitulate both positive and negative effects of *trx* and *Pc*, respectively, has previously been identified (13). In this study, we used immunoaffinity chromatography to purify tagged Pc-G complexes and then we assayed their DNA binding activity. We found that the $(GA)_n$ motif in this fragment is indeed a primary binding site for the Pc-G complexes. We also provided several lines of evidence to show that the DNA binding protein is encoded by the *pipsqueak* (*psq*) gene, previously only known for its role in oogenesis and eye development (27, 52, 61).

MATERIALS AND METHODS

Fly works. Flies were raised at 25°C on standard food. To test genetic interactions, *psq* alleles were rebalanced with *SM6-TM6B,Tb* since the *CyO* balancer present in some of the original stocks appeared to show significant interaction with *Pc*⁴. Virgin females heterozygous for various *psq* alleles (0115, 2403, 8109, and $\Delta 18$ from C. Berg; F112, E34, and E39 from U. Weber) were crossed with males heterozygous for *Pc*⁴, *Psc*¹, *Scm*^{D1}, *esc*¹⁰, or *E(z)*⁶³, and the number of sex comb teeth in the second and third legs of their male progeny were counted as described previously (14). To identify larvae heterozygous for both *psq* and *Pc* mutations, female flies carrying *psq* alleles balanced with *SM6-TM6B,Tb* were crossed with *Pc*⁴/*TM6B,Tb* male flies. Larvae without the dominant *Tb* phenotype carried by the *TM6B* balancer were selected for subsequent disk staining.

Plasmids and probes. To generate the pMT/Pc-FH clone, the DNA sequence from the termination codon of *Pc* up to the *Bgl*II site of pAct/Pc-1 (13) was first replaced by the DNA sequence encoding a FLAG peptide and hexahistidine (TTGATATCAGATCTGATGGACTACAAGGACGATGACGATAAG AAC GCGTCCACCATCACCACCATCACTAGGATCT; details available upon request) to make pAct/Pc-FH. The *Sma*I/*Sal*I fragment of pAct/Pc-FH was then cloned into the *Xho*I/*Sal*I sites of pMT/Hy (37) after the repair of the *Xho*I site by Klenow enzyme. The *bxd*-a and *bxd*-b clones were generated from B-151 (13) by dividing the ~440-bp *bxd* fragment at the internal *Bgl*II site and inserting it into the *Pst*I/*Eco*RV sites of pBluescript after T4 polymerase treatment. *bxd*-1, -2, -3, and -4 were generated by PCR amplification and subsequent cloning into pBluescript (Stratagene). p(GA)_n and p(Z)_n were generated from B120/GAGA and B120/ZESTE plasmids (13, 39), respectively, by inserting into pBluescript the amplified DNA sequences corresponding to positions -321 to -12 of *Ubx*. For competition assays, DNA fragments were prepared following digestion by restriction enzymes: *Xba*I/*Xho*I for *bxd*-a and -b; *Sac*II/*Xba*I for *bxd*-1, -2, -3, and -4 and p(GA)_n and p(Z)_n; *Pvu*II for pBluescribe (Stratagene).

For the construction of pET/GAGA, a *Sma*I/*Kpn*I fragment corresponding to the full-length cDNA of GAGA-519 was excised from pBluescript/GAGA (a gift from T. Tsukiyama and C. Wu) and first cloned into a modified pPac5c-PL vector

(a gift from C. Thummel) that contains a DNA fragment encoding the FLAG peptide in the polylinker (details available upon request). The *Sna*BI/*Sac*I fragment was then inserted at the *Nde*I/*Hind*III sites of the pET15b vector (Novagene), after repair of the *Nde*I site. For the construction of pQE/PSQ-N, the *Spe*I/*Mlu*I fragment corresponding to the C-terminal part of *psq-1* was deleted from pHH14 (a gift from C. Berg) (27). For pQE/PSQ-C, the *Bam*HI/*Sma*I fragment corresponding to the N-terminal part of *psq-1* was deleted from pHH14.

Cell culture, protein, and antibody purification. The plasmid pMT/Pc-FH was transfected into S2 cells and selected for with hygromycin (200 μ g/ml) as described previously (37). Stable PC-FH cell lines were grown in spinner flasks and induced at a density of 2×10^6 to 4×10^6 cells/ml with 0.1 mM CuSO₄ for 67 h. The amount of Pc proteins induced at this concentration was less than 10% of the amount induced by 0.7 mM CuSO₄ (data not shown). As described earlier (14), the Pc protein complex was purified by immunoaffinity chromatography from the PC-FH cell line by making a 10 to 40% (NH₄)₂SO₄ differential precipitation of the nuclear extract, followed by FLAG peptide elution from a FLAG antibody column (M2; Kodak).

For expression of recombinant proteins, pET/GAGA was transformed into BL21(DE3) cells and pQE/PSQ-N and PSQ-C were transformed into TG-1 cells. Soluble proteins were obtained under the following induction conditions: 25°C for 90 min for GAGA and 25°C for 5 h for both PSQ-N and PSQ-C. Proteins were purified from a nickel-nitrilotriacetic acid column according to the vendor's instructions (Qiagen). Purified PSQ-N and PSQ-C were coupled to Affi-Gel 10 resin (Bio-Rad) in acetate buffer (0.2 M, pH 5.0) (16) and Clark and Lubs buffer (pH 8.5) (16), respectively.

AS-2 serum (a gift from C. Berg) (27) was used for affinity purification of PSQ antibody against specific parts of Psq proteins. However, the antibody purified from the PSQ-N column appeared to have strong cross-reactivity against many proteins on the Western blot, presumably due to the BTB and glutamine-rich domains. Serum extensively preabsorbed with PSQ-N resin was then passed through a PSQ-C column to purify the corresponding antibody by standard acid elution (23).

Electrophoretic mobility shift assay (EMSA). Binding reactions were carried out at 30°C for 20 min in a 10- μ l solution containing binding buffer (20 mM Tris-HCl [pH 7.8], 60 mM KCl, 0.1 mM EDTA, 1 mM dithiothreitol, 0.1% Tween 20, 10% glycerol), 5 μ g of bovine serum albumin, 0.5 to 1 ng of ³²P-labeled probe, and poly(dI-dC) (0.1 μ g for PC and 0.5 μ g for GAF). Samples were run at 4°C for ~3 h on a 3.5% polyacrylamide gel (37.5:1 acrylamide-to-bisacrylamide ratio; Bio-Rad) in 1 \times TAE (40 mM Tris-acetate [pH 8.0], 1 mM EDTA) after a preelectrophoresis for about half an hour. Competition experiments were carried out in the presence of ~30 ng of purified DNA fragments. For antibody supershift experiments, the PC protein complex was incubated with affinity-purified PC (2.5 mg/ml) antibody or goat immunoglobulin G (IgG) (2.5 mg/ml) for 1 h on ice prior to the addition of other components. The amount of poly(dI-dC) was increased to 0.25 μ g for these assays.

UV cross-linking. The oligo probe used for UV cross-linking studies was made by incorporating radioactive dCTP and cross-linkable 5-[N-(p-azidobenzoyl)-3-aminoallyl] dUTP (AB-dUTP) specifically at the binding site as described previously (2). The probe was made by first annealing a 47-mer (5'-CCTCCTCTTCCTGGAGAGGGGAGAGAGGCACGACTTAACGCATACAC-3') to a 19-mer (5'-GTGTATGCGTAAAGTCGTG-3'). Approximately 50 ng of hybrid DNA was partially filled by exonuclease-free Klenow enzyme (New England Biolabs) in the presence of ~0.67 μ M [α -³²P]dCTP (~3,000 Ci/mmol; Amersham), 1 μ M dCTP, 20 μ M AB-dUTP (a gift from P. Geiduschek), and 5 μ M dTTP, followed by the addition of 250 μ M dNTP to fill up the remaining gap in the probe DNA. The probe was purified by phenol-chloroform extraction and repeated ethanol precipitation in the presence of 0.5 M ammonium acetate and glycogen (200 μ g/ml) (Boehringer Mannheim).

DNA binding reactions with 10- μ l reaction mixtures were carried out as described above, except that bovine serum albumin was omitted. Open vials were UV irradiated for 3 min at 0.4 W in a UVC-515 UV multilinker (UV LUM). MgCl₂ and CaCl₂ were then added to a final concentration of 8 mM. Following a 30-min incubation at 30°C with 10 to 30 U of DNase I (Sigma), the DNA-protein complexes were further digested by 0.5 U of micrococcal nuclease (Sigma) for 20 min. Samples were then analyzed by sodium dodecyl sulfate (SDS)-polyacrylamide gel electrophoresis on a 9% gel according to standard procedure.

Immunostaining. For disc staining, imaginal discs from wandering third-instar larvae were fixed and stained with UBX antibody (1:2 dilution of FP.3.38 [62]). Labeling was detected by the ABC detection method (Vector) and diaminobenzidine staining. For polytene chromosome staining, salivary glands were dissected, fixed, and squashed according to the standard protocol (65), except that the formaldehyde concentration was reduced to 2.5%. For double immunoflu-




FIG. 1. Purification of tagged Pc-G complex. (A) Structure of the epitope-tagged *Pc* construct. The chromodomain is indicated by a solid box. The C-terminal FLAG epitope and hexahistidine tags are labeled FLAG and (H)₆, respectively. The extra amino acid residues introduced by the linker sequences are shown in single letter code. (B) Silver staining of eluted proteins. Aliquots (3.5 μ l) of proteins from the column input and different fractions were resolved on an SDS-7% polyacrylamide gel. The fraction numbers are indicated at the top of each lane and are consistent with those in Fig. 2D and 3B. The position of PC is indicated by the arrows. Note that several discrete protein bands can be resolved in this region on high-resolution gels. (C) Specific association of Pc-G proteins. The eluted fractions prepared from PC-FH or S2 cells were resolved on an SDS-7% polyacrylamide gel. Only a small number of nonspecific proteins were eluted from S2 cell extracts.

orescence staining, PSC monoclonal antibody 6E8 (1:10 purified antibody from Developmental Studies Hybridoma Bank) and affinity-purified rabbit PSQ antibody (1:500) were used. The secondary antibodies were conjugated with Cy5 and Rhodamine Red-X for the anti-mouse and anti-rabbit antibodies, respectively. Images were obtained with a Zeiss LSM310 confocal microscope.

RESULTS

(GA)_n as a critical motif in PRE. To facilitate the biochemical study of the Pc-G complex, we established a *Drosophila* S2 cell line PC-FH that can express *Pc* protein with both FLAG epitope and hexahistidine tags at its C terminus (Fig. 1A). These modifications do not appear to affect the activity of *Pc*, since similar constructs can repress the *Ubx* reporter gene in cultured cells and partially rescue *Pc* mutants in transgenic flies (data not shown). The tagged PC is under the control of a metallothionein promoter, which allows protein induction by the copper ion in a range up to 0.7 mM (9; data not shown). We chose a suboptimal concentration of CuSO₄ (i.e., 0.1 mM) for induction, since it appears to provide sufficient amounts of tagged protein complexes for purification. Nuclear extracts prepared from induced cells were fractionated by 10 to 40% (NH₄)₂SO₄ precipitation to enrich for large protein complexes. The extracts were then passed through a FLAG antibody column (i.e., M2 resin) and eluted with the FLAG peptide. We estimated that approximately 200-fold purification was obtained by affinity chromatography compared with the crude extracts (Fig. 1B). As shown in Fig. 1C, many proteins ap-

peared to be specifically coeluted with the *Pc* protein. Although the region corresponding to the size of *Pc* proteins appears to be heavily stained, the relative abundance of *Pc* proteins has been exaggerated by the presence of several proteins of similar size that can be better separated in high-resolution gels (data not shown). An earlier study has shown that a specific subset of Pc-G proteins, including PSC, PH, and HDAC1 were copurified, indicating the presence of multimeric Pc-G complexes in these fractions (14). Since this is the first *Drosophila* Pc-G complex shown to contain both histone modification activity (14) and DNA binding activity (see below) and since homeotic genes are its best characterized targets, we shall refer to this complex as CHRASCH (chromatin-associated silencing complex for homeotics) to distinguish it from commonly referred Pc-G complexes or complexes characterized by other workers (44, 51, 59).

Since a functional Pc-G complex must act specifically on its response element (i.e., PRE), we examined whether CHRASCH can bind such sequences *in vitro*. PRE from several homeotic genes have been mapped, including the one from the upstream *bxd* region of *Ubx* (12, 13, 15). It was previously shown that an ~440-bp fragment (B-151) from this region recapitulates transcriptional regulation by either *Pc* or *trx* in cultured cells (13). DNA fragments encompassing this region can also confer pairing-sensitive repression and are enriched for *Pc* proteins in chromatin immunoprecipitation experiments (12, 57). B-151 therefore contains physiologically relevant binding sites for the Pc-G complex. Using two subfragments from B-151 as probes for EMSAs, we found that CHRASCH binds strongly to the *bxd*-b fragment (Fig. 2B), resulting in several slow-migrating bands. As will be shown later, these bands presumably reflect the binding to a reiterated motif in the *bxd*-b fragment. The binding of CHRASCH to the *bxd*-b fragment appeared to be specific, since it could be completely competed out by the addition of *bxd*-b but not by *bxd*-a or nonspecific vector sequences (Fig. 2B and C). A much weaker but specific binding of CHRASCH to *bxd*-a was also detected. The observation that *bxd*-b could compete for binding to *bxd*-a (Fig. 2B, lane 4) but that *bxd*-a could not compete effectively for binding to *bxd*-b (Fig. 2B, lane 8) suggested that these fragments might contain similar binding sequences, albeit with a lower affinity in the *bxd*-a fragment. Due to the difficulty in studying a weak binding activity with certainty, our subsequent studies focused on *bxd*-b.

We used four partially overlapping fragments from B-151 in competition assays to further map the binding sites of CHRASCH. As shown in Fig. 2C, only the *bxd*-3 and *bxd*-4 fragments compete effectively for the binding to CHRASCH. In addition, *bxd*-4 appeared to compete better than *bxd*-3. Therefore, we deduced that the right half of B-151 must contain the primary binding sites for CHRASCH. Interestingly, transgenes containing small deletions in this region, but not in the left half of B-151, failed to silence the reporter gene effectively and could no longer respond to mutations in several Pc-G (60), indicating that this region is indeed relevant for Pc-G-mediated silencing. Three major sequence motifs can be identified in B-151. The first motif (C/T)GAG(C/T)G is the consensus binding site of the *Zeste* protein (47). Both the left and right halves of B-151 contain one *Zeste* binding site. The second motif ATGGC represents the binding site of a newly characterized member of




FIG. 2. CHRASCH binds the $(GA)_n$ motif. (A) Map of DNA fragments in B-151 used for binding and competition assays. The locations of the $(GA)_n$ motif (solid rectangular box), YY1 (solid circles), and ZESTE (open circles) binding sites are shown. The coordinates of restriction sites and various fragments are indicated. (B) Preferential binding of CHRASCH to the bxd-b fragment. Binding of labeled bxd-a (a) or bxd-b (b) probe was carried out with or without (-) an excess amount of unlabeled bxd-a or bxd-b or a DNA fragment from the polylinker of a pBluescribe vector

Pc-G, *pho*, which encodes the *Drosophila* homologue of YY1 (8, 43). *bxd-3* and *bxd-4* each contain one copy of an almost identical YY1 site. The third motif is a $(GA)_n$ repeat, which represents the consensus binding site of GAF encoded by *Trithorax-like* (*Trl*) (21). *Trl* was originally identified as a member of *trx-G* (18); however, some recent studies suggest that it may also share some characteristics with Pc-G (22, 24). While both *bxd-3* and *bxd-4* contain 2 separate clusters of this motif, one cluster in *bxd-4* is much further extended (GAGAGAGG GAGAG versus GAGAG). Since *bxd-4* has been shown to be more effective in competition assays, it is likely that the $(GA)_n$ motif is most critical for CHRASCH binding. This possibility is further supported by the observation that CHRASCH binding to *bxd-b* was completely competed out by a fragment containing multiple $(GA)_n$ repeats (Fig. 2C, lane 10) but not by the one containing multiple Zeste repeats (Fig. 2C, lane 11) or an oligonucleotide containing a YY1 binding site (data not shown). Therefore, we conclude that the binding sequences of CHRASCH consist primarily of the $(GA)_n$ motif.

The association of the $(GA)_n$ binding activity with CHRASCH was further confirmed by the following observations. When the DNA binding activity was examined in peptide-eluted fractions, we found that fractions 6 and 8 had the strongest binding activities (Fig. 2D). These fractions also contained the highest amounts of PC and other associated proteins (compare Fig. 1B and 2D). Thus, the binding protein coeluted with CHRASCH in the immunoaffinity chromatography. In addition, we found that the DNA-protein complexes formed on *bxd-b* could be slightly supershifted by a preincubation with an affinity-purified PC antibody but not with a nonspecific IgG (Fig. 2E, lanes 4 and 5). The small supershift might be expected for a large complex in a gel composed of 3.5% polyacrylamide. Taken together, these results indicate that the $(GA)_n$ binding protein is physically associated with CHRASCH.

Psq proteins are responsible for $(GA)_n$ binding. Since chromatin immunoprecipitation experiments have shown that GAF is enriched in the region encompassing B-151 and its vicinity (57), we examined whether the binding protein of CHRASCH is related to GAF. To our surprise, our results were not consistent with this notion. As expected, a purified recombinant GAF bound specifically to the *bxd-b* fragment. In addition, the binding activity of GAF was drastically stimulated by zinc ions over a wide range, resulting in a further retardation of the DNA-protein complexes in the EMSA (Fig. 3A). By contrast, the binding activity of CHRASCH was not significantly affected at intermediate concentrations of zinc ions and became completely inactivated at a high concentration (i.e., 0.5 mM). The differential effects of zinc ions on DNA binding properties argue that different DNA binding proteins are involved in the




FIG. 3. Lack of GAF in CHRASCH. (A) Effects of zinc ion on binding activities of GAF and CHRASCH. Different concentrations of $ZnSO_4$ were included in binding reaction mixtures, followed by electrophoresis on 3.5% polyacrylamide gels. The concentrations used are as follows: none (lanes 2 and 10), 10 μM (lanes 3 and 11), 20 μM (lanes 4 and 12), 50 μM (lanes 5 and 13), 100 μM (lanes 6 and 14), 200 μM (lanes 7 and 15), and 500 μM (lanes 8 and 16). Specific bands are indicated by dots. In contrast to stronger binding for GAF at higher $ZnSO_4$ concentrations, CHRASCH is refractory to intermediate concentrations and then becomes completely inactive. (B) Proteins from both column input (1 μl) (lanes 1 and 3) and fraction 8 (5 μl) (lanes 2 and 4) were resolved on an SDS-10% polyacrylamide gel, transferred, and assayed by GAF antiserum before (lanes 1 and 2) or after (lanes 3 and 4) a 1-h preincubation with purified recombinant GAF (~16 $\mu g/ml$). Two major clusters of GAF isoforms, GAGA-519 and GAGA-581, are indicated. Preincubation of recombinant GAF strongly reduces the reactivity of the antiserum to GAF in the input but not to the unknown protein in the purified fraction, indicating that this protein is not related to GAF.

binding. Furthermore, we used a GAF-specific antiserum to examine our protein preparations. GAF antiserum detected multiple bands corresponding to two major classes of isoforms, GAF-519 and -581 (4), from the column input. A weak band migrating slightly slower than one GAF-519 isoform, however, was detected in the eluted fraction (Fig. 3B, compare lanes 1 and 2). To ascertain its identity, we asked whether the reactivity of GAF antiserum could be blocked by GAF. Preincubation

(ns). Specific binding to *bxd-a* and *bxd-b* probes is indicated by the arrowhead and arrow, respectively. (C) Binding of CHRASCH to the $(GA)_n$ motif. Binding assays were carried out with the *bxd-b* probe. The competitors were *bxd-a* (a), *bxd-b* (b), *bxd-1* to -4 (1 to 4), linker DNA (ns), or fragments containing multiple $(GA)_n$ (G) or ZESTE (Z) binding sites. (D) Elution profile of the DNA binding activity. Aliquots (2 μl) of eluted fractions (same fractions as shown in Fig. 1A) were assayed with the *bxd-b* probe. (E) Antibody supershift assay. Affinity-purified PC antibody (2.5 mg/ml), IgG (2.5 mg/ml), or buffer were preincubated with CHRASCH before the addition of a reaction mixture containing the *bxd-b* probe. The volume of antibody solution is indicated. The supershifted complex is indicated by the arrowhead. Note that fast-migrating bands appeared after 1 h of preincubation (lanes 2 and 6), suggesting instability of the complex. The EMSA shown here was done with 3.5% polyacrylamide gels. B, bound DNA-protein complex; F, free probe.

of GAF antiserum with purified recombinant GAF could cause almost complete loss of its reactivity to GAF in the input fraction but had no effect on the reactivity to the band detected in the eluted fraction (Fig. 3B, lane 4). Therefore, this band most likely arises from a nonspecific cross-reactivity.

To further characterize the DNA binding protein of CHRASCH, we used a cross-linking method to specifically label this protein (2). An oligonucleotide probe was designed that allowed specific incorporation of both radioactive dCTP and a photoactivated cross-linking dTTP analogue (i.e., AB-dUTP) into the nucleotide sequences that correspond to the extended (GA)_n motif in *bxd-4* (Fig. 4A). Following DNA binding and UV irradiation, the cross-linked nucleoprotein complex was digested extensively with both DNase I and micrococcal nuclease to remove excessive DNA sequences. The indirectly labeled proteins were then resolved on an SDS-9% polyacrylamide gel. As shown in Fig. 4B, a major band of ~85 kDa was identified for the purified recombinant GAF, whereas a different pattern was observed for CHRASCH, which consists of a doublet of ~130 kDa and a doublet of ~70 kDa. Taken together, our results provide strong evidence that the DNA binding activity of CHRASCH is not contributed by GAF but by a novel factor(s).

A new (GA)_n-binding protein has been identified incidentally in *Apis mellifera* by screening an expression library with (GA)_n repeats (40). The *Drosophila* homologue of this protein was found to be encoded by *pipsqueak* (*psq*), identified originally by its grandchildless phenotype and subsequently by its effect on eye development (27, 52, 61). By differential transcriptional and translational initiation, *psq* produces two major mRNAs containing open reading frames for 1,065 (PSQ-A) and 630 to 646 amino acids (PSQ-B). These two isoforms share a common C-terminal PSQ domain capable of binding to the (GA)_n motif (40). The size similarity between one of the cross-linked proteins and PSQ-A prompted us to test whether Psq proteins are copurified with CHRASCH. Using an antibody that reacts with both PSQ-A and B (27), we found that PSQ-A and much less PSQ-B are clearly detectable in our sample (Fig. 4C). For a preparation of FLAG-tagged TATA-box binding protein (TBP), however, we found a large amount of PSQ-A in the input but not in the eluted fraction (Fig. 4C). These results demonstrate that PSQ-A and CHRASCH have indeed been copurified. To further examine the association between PC and PSQ in vivo, we performed coimmunoprecipitation with embryonic nuclear extracts. As shown in Fig. 4D, PC was precipitated by the PSQ antibody but not by the control serum. Although it is not clear whether the smaller ~70-kDa proteins detected in the cross-linking experiments represent degradation products of PSQ or other unrelated proteins, our results strongly suggest that PSQ-A may play a major role in DNA binding.

Selective role of *psq* in vivo. To determine the role of *psq* in vivo, we used a dosage-sensitive assay to test the genetic interactions between *Pc* and *psq* mutations (35). Two classes of *psq* mutations have been previously identified, each has been implicated in a distinct developmental function. For example, the *0115*, *2403*, and *8109* alleles (referred to as class I mutations) that result from P element insertion mutations in the ~40-kb intron (see Fig. 7A) appear to primarily affect oogenesis, giving rise to the grandchildless phenotype (27, 52), whereas the




FIG. 4. Psq proteins are directly involved in (GA)_n binding. (A) Probe used for UV cross-linking of (GA)_n binding protein. The DNA sequences of the upper and the lower strands are shown. Two stretches of the (GA)_n motif are underlined, which are almost identical to the extended (GA)_n motif in the *bxd-4* fragment, except that their exact order is reversed. The sequences flanking both sides of the (GA)_n motif in the probe bear no similarity to those in the *bxd-4* fragment. Note that the incorporation of AB-dUTP and radioactive dCTP are restricted to the (GA)_n motif only. (B) UV cross-linking of the binding protein in CHRASCH. Aliquots of recombinant GAF and CHRASCH that gave equivalent DNA binding activities, as judged by EMSA (data not shown), were used for UV cross-linking. After extensive digestion with various amounts of DNase I (10, 20, and 30 U) and 0.5 U of micrococcal nuclease, indirectly labeled proteins were resolved on an SDS-9% polyacrylamide gel. One major protein species of ~85 kDa was found for purified GAF. Two sets of bands were found for CHRASCH. One set of bands was substantially larger than the 106.5-kDa mass marker, while the other one was ~70 kDa. (C) Copurification of Psq proteins. The input and peak fractions from extracts containing tagged PC or TBP were resolved on a 7% polyacrylamide gel. After transfer to a nitrocellulose filter, the proteins were probed with an affinity-purified antibody specific for the C-terminal common half of PSQ. The left panel represents the immunoblot for CHRASCH, while the right panel represents that for TBP as a control. Note that PSQ-A and PSQ-B migrate as ~150- and ~95-kDa proteins, respectively. Note that the size of PSQ-B is larger than the one (~80 kDa) reported by Horowitz and Berg (27). (D) Association of PSQ and PC in vivo. Embryonic nuclear extracts were immunoprecipitated by an affinity-purified PSQ antibody (IP) or nonimmune serum (control). The extracts (input) and immunoprecipitates were probed with an affinity-purified PC antibody. The positions for PC and Ig are indicated.

F112, *E34*, and *E39* alleles (referred to as class II mutations) that are clustered around the first exon of *psq-B* (or *psq-2* by Horowitz and Berg [27]) appear to affect eye development (61). Although none of these *psq* mutations alone have been known to cause homeotic phenotypes in adults, it does not necessarily preclude their role in homeotic gene silencing, since an increasing number of *Pc-G* interacting genes have been found to cause little, if any, homeotic phenotype by themselves (14, 33, 56). Thus, we crossed females heterozygous for various *psq* alleles with male *Pc*⁴ heterozygotes and examined their effects on homeotic leg transformation, i.e., production of

TABLE 1. Genetic interaction between *psq* and *Pc*

Expt ^a	<i>psq</i> allele	Avg no. of ectopic sex comb teeth/leg for genotype ^b :	
		<i>Bal Pc</i>	<i>psq Pc</i>
A	<i>0115</i>	4.08 (118)	10.13 (167)
	<i>2403</i>	3.44 (148)	8.58 (184)
	<i>8109</i>	3.31 (140)	10.02 (131)
	<i>F112</i>	3.67 (168)	5.17 (216)
	<i>E34</i>	3.86 (175)	4.41 (216)
	<i>E39</i>	3.44 (126)	3.81 (108)
B	<i>0115</i>	3.77 (102)	7.51 (135)
	$\Delta 18$	2.52 (84)	8.90 (132)
	<i>8109</i>	3.00 (66)	8.78 (136)

^a Virgin females of various *psq* mutant alleles were crossed with *Pc*⁴ mutant males. Two separate experiments were done to determine the allelic strength in interactions with the *Pc*⁴ mutation.

^b The average number of ectopic sex comb teeth per leg was calculated by dividing the total number of ectopic sex comb teeth by the number of second and third legs (values in parentheses) for each genotype. The number of sex comb teeth on the first leg of wild-type males usually varies between 10 and 12. A reciprocal cross between *Pc*⁴ females and *psq*²⁴⁰³ males gave a lower number (1.31) of ectopic sex comb teeth in *Pc*⁴ progeny but a similar number (8.30 versus 8.58) in double-mutant progeny, indicating a strong maternal effect of *psq*.

ectopic sex comb teeth on the second and third legs of F₁ males. As shown in Table 1, the average number of ectopic sex comb teeth is strikingly enhanced for all alleles of class I mutations when doubly heterozygous with the *Pc*⁴ mutation (ranging from 9 to 10 teeth per leg), and the second and third legs were almost completely transformed into the first leg. In contrast, class II alleles showed relatively weak but significant




FIG. 5. Synergistic effect of *psq* and *Pc* mutations on ectopic UBX expression in imaginal discs. First (A and C)- and second (B and D)-leg discs and wing discs (E and F) from wild type (A, B, and E) or double heterozygotes (C, D, and F) of *psq*²⁴⁰³ and *Pc*⁴ were stained with a monoclonal UBX antibody. UBX is normally undetectable in the first-leg disk and detectable at a low level in second-leg and wing discs of wild-type larvae. A slight increase of UBX is observed in the wing discs of larvae heterozygous for *Pc*⁴. Strong UBX is observed in the discs of double mutants. The *psq* mutation alone has no detectable effect on UBX expression. Essentially the same results were observed for *psq*⁰¹¹⁵.




FIG. 6. Colocalization of PSC and PSQ on ANTP-C and BX-C. Polytene chromosomes from third-instar larvae were simultaneously stained with an affinity-purified rabbit PSQ antibody and a mouse monoclonal PSC antibody. The merged images are shown with DNA in green (A and E), PSQ staining in red (A, B, E, and F), and PSC staining in blue (A, B, E, and F). The images for single staining of PSQ (C and G) or PSC (D and H) are also shown. The sites for ANTP-C (A to D) and BX-C (E to H) are indicated by arrows.

effects on leg transformation, giving rise to 3 to 5 ectopic teeth per leg.

Since three major protein complexes containing different combinations of Pc-G proteins have been described previously (14, 24, 44, 51, 59), it is interesting to determine whether the function of *psq* is generally or specifically required for these complexes. Pc-G mutations that are representative of these protein complexes were tested. Our preliminary results showed that there was a remarkable increase in the number of ectopic sex comb teeth in the progeny carrying both *Psc*¹ and *psq*²⁴⁰³ mutations compared to those carrying the *Psc*¹ mutation alone. The *Scm*^{D1} mutation displays an intermediate level of interaction, giving rise to approximately two- to threefold increases, whereas no significant enhancement was found for the *esc*¹⁰ or *E(z)*⁶³ mutations. These results suggest that *psq* is crucial for proper function of a subset of Pc-G proteins that are constituents of CHRASCH.

We further examined the effect of *psq* mutations on *Ubx* expression in imaginal discs. UBX proteins are normally expressed at high levels in the haltere- and third-leg discs but at low levels in the peripodial membranes of the wing discs. No significant change was observed in larvae heterozygous for either the *Pc*⁴ or *psq* mutation (data not shown). However, high levels of UBX proteins were observed in the medial sections of the wing discs from larvae doubly heterozygous for *Pc*⁴ and *psq* (Fig. 5F). In addition, substantial amounts of UBX proteins could be detected in the first- and second-leg discs (Fig. 5C and D).

Given the copurification and genetic interactions between *psq* and Pc-G, one might expect PSQ to be detected on polytene chromosomes at sites corresponding to ANTP-C (at 84AB) and BX-C (at 89E), where strong Pc-G signals

are normally observed. Using antibodies against PSC and PSQ for double immunofluorescent staining of polytene chromosomes under standard fixation conditions (i.e., 3.7% formaldehyde), we could not find consistent PSQ staining at ANTP-C and BX-C sites. It was possible, however, that the PSQ staining at these sites might have been masked by other proteins of the complex that are apparently layered upon it. Thus, we tested conditions to alleviate any potential masking effect. Using 2.5% formaldehyde for fixation, we found that PSQ signals can be detected at ANTP-C and BX-C sites (Fig. 6), with the PSC staining being consistently weaker than what was observed under standard conditions. The staining of PSC and PSQ at these sites was not always coextensive. For example, the PSQ signal represents only a part of the broader region of PSC staining at the ANTP-C site (Fig. 6A to D). The significance of this observation remains unclear.

We also examined the staining patterns on polytene chromosomes from transgenic flies containing insertions of small PRE fragments (e.g., PRE-D) (20). While these fragments showed ectopic PSC staining as expected, most of these fragments appeared to insert at sites that correspond to endogenous PSQ sites on wild-type chromosomes (data not shown). Thus, we could not use this approach to map the exact locations of PSQ binding sites within the PRE.

PSQ-A is essential for silencing. Class I *psq* mutations have previously been shown to cause a more-severe reduction in the level of PSQ-A than in that of PSQ-B (27). It raised an interesting possibility that different classes of *psq* alleles might have differential effects on these two major protein species and that such effects might be correlated with their contribution to the silencing function. The relative abundance of these proteins from adult ovaries of several homozygous mutants was determined. As shown in Fig. 7B, there are indeed some fluctuations in the relative abundance of these two proteins, however, the fluctuations between different alleles within one class or between different classes do not seem to be consistent with the possibility that a specific Psq protein may be more critical for gene silencing. For example, while both *psq*⁰¹¹⁵ and *psq*²⁴⁰³ mutants contain much less PSQ than *psq*⁸¹⁰⁹, *psq*²⁴⁰³ shows consistently weaker interaction with *Pc* than *psq*⁰¹¹⁵ and *psq*⁸¹⁰⁹. In addition, although *psq*⁸¹⁰⁹ and *psq*^{E34} contain similar amounts of PSQ, *psq*⁸¹⁰⁹ shows substantially stronger interaction with *Pc* than *psq*^{E34}. Thus, it seems unlikely that the homeotic phenotype can be attributed to the mere presence of different PSQ in a simple manner.

In addition to their effects at the level of gene expression, however, class I mutations can also result in the synthesis of a novel BTB-domain-containing fusion product [i.e., *psq*-*l(3)S12*] due to aberrant splicing (26). It has further been shown that overexpression of this fusion protein in wild-type flies is sufficient to solicit phenotypes similar to those of class I mutations (27). It is believed that this fusion protein may interfere with the function of PSQ-A through the shared BTB protein interaction domain (1, 27) since this domain is present in PSQ-A protein but not in PSQ-B protein. Taking this possibility into consideration, the stronger interaction of class I mutations with the *Pc* mutation described above might result from a combination of reduced expression and an interference with PSQ-A protein, leading to a specific loss of PSQ-A function. To substantiate this possibility, we tested whether the loss




FIG. 7. Relative abundance of Psq proteins in ovaries of *psq* mutants. (A) A map of the *psq* locus is redrawn on the basis of the information from Horowitz and Berg (27) and Weber et al. (61). The genomic structure of *psq* and the positions of class I mutations including three PZ insertion mutants (0115, 2403, and 8109) and a deletion mutant ($\Delta 18$) are shown above the map. The structures of two major cDNAs, *psq*-1 and *psq*-2, are shown below, with their splicing junctions indicated. The non-coding sequences (blank boxes) and the coding regions (solid boxes), including the N-terminal BTB (stippled boxes) and C-terminal PSQ domains (grey boxes), are also shown. Three class II mutations, F112, E34, and E39, are located around the first exon of *psq*-2. The approximate sites of insertion (F112) or deletion (E34 and E39) are indicated. *psq*-1/*l(3)S12* represents a fusion product resulting from an aberrant splicing between *psq*-1 and *l(3)S12* adventitiously present in the PZ transposons. The positions of two recombinant proteins, PSQ-N and PSQ-C, used for antibody purification are also indicated. The DNA fragments are drawn to scale except for the ~ 40 -kb intron. (B) Protein analysis of *psq* mutants. Ovaries were dissected from homozygous *psq* or transheterozygous mutant adults, and proteins were analyzed by SDS-polyacrylamide gel electrophoresis. *psq* proteins were detected with affinity-purified antibody against PSQ-C. Extracts derived from wild-type (WT) or mutants are indicated above the respective lanes. Homozygous *psq*^{E39} animals died during pupation. The positions for PSQ-A, PSQ-B, and an internal control (α -tubulin) are marked by arrows. The positions of mass markers (in kilodaltons) are marked. Note that trace amounts of PSQ-A could be detected in *psq* ^{$\Delta 18$} /*psq*⁰¹¹⁵ mutants when the film was exposed for a longer amount of time. (C) Relative abundance of *psq* proteins. The relative amounts of PSQ-A and PSQ-B in these mutants are derived by first calibrating against the internal control (i.e., α -tubulin) in each sample. Each value was then calculated by using wild-type proteins as references.

of PSQ-A alone is sufficient to cause a strong interaction with the *Pc* mutation. *psq* ^{$\Delta 18$} is a homozygous lethal mutation that deletes the 5' exons of *psq*-1 and the intergenic region between *psq* and a divergently transcribed gene, *lola*. As expected, while

the level of PSQ-B remains normal, very little PSQ-A is detected in ovaries from *psq*⁰¹¹⁵/*psq*^{Δ18} transheterozygotes (Fig. 7B), indicating that PSQ-A is specifically affected by the *psq*^{Δ18} mutation. When heterozygous *psq*^{Δ18} females were crossed with heterozygous *Pc*⁴ males, a very strong leg transformation was observed in their male progeny (Table 1, experiment B). Indeed, both the penetrance and expressivity caused by the *psq*^{Δ18} mutation are comparable to, if not stronger than, the two strongest *psq* alleles. Therefore, our results strongly support an essential role for PSQ-A in homeotic gene silencing.

DISCUSSION

The molecular mechanism responsible for sequence-specific targeting of Pc-G proteins has been elusive to date, partly because of the lack of strict consensus sequences of PRE and partly because of the plethora of proteins involved. In addition, given the identification of at least three physically separable Pc-G protein complexes (14, 24, 44, 51, 59) and the finding that different combinations of Pc-G proteins are detected on different target genes (58), it is likely that several strategies are employed by these complexes at different developmental stages, in different cell types, or at different target genes. Nevertheless, in this report, we provide several lines of evidence to show that a novel DNA binding factor encoded by *psq* is a constituent of CHRASCH, a previously characterized major Pc-G protein complex (14). Since CHRASCH also contains a histone modification factor, HDAC1, we suggest that this complex may represent a fully functional entity that can nucleate certain chromatin structures at and around specific sequences (i.e., PRE) of homeotic genes.

Biochemical purification of Pc-G protein complexes has been limited by their apparent instability (50; data not shown). Thus, a balance between biochemical purity and functional integrity might be considered. Different approaches are required subsequently to substantiate the physiological relevance of copurified proteins. To meet these criteria, we adopted the strategy of purifying Pc-G protein complexes to sufficient homogeneity mainly by immunoaffinity chromatography under moderate conditions, then examining the biochemical functions potentially relevant to these complexes, followed by identifying the functional constituents of the complex and corresponding genes, and finally validating their roles with genetic studies.

The *bxd* region has been extensively examined for PRE. Although different fragments ranging from ~400 bp to ~1 kb have been studied (12, 13, 24, 25, 60), they share a common region represented almost entirely by the B-151 fragment analyzed in this study. Among the three binding motifs of this fragment, we found that the (GA)_n motif represents the most prominent binding site for CHRASCH. In recent studies, the role of this motif in silencing has been demonstrated in transgenic flies (24). Thus, we believe that this motif plays a critical role in anchoring one of the major Pc-G complexes (i.e., CHRASCH). Our results, however, are not mutually exclusive to the possibility that other motifs may be required for different functional aspects of PRE.

One of the most critical issues concerning the specific targeting of the Pc-G complex appears to reside in the identity of the DNA binding factor. Our results support the conclusion

that PSQ-A plays a primary role in such a function for the following reasons. First, PSQ-A, but not GAF, is copurified with CHRASCH. Second, UV cross-linking studies strongly indicate that PSQ-A binds directly to the (GA)_n motif. Additional proteins, however, were also evident in these studies. At present, we cannot distinguish between the possibilities that these proteins represent degradation products of PSQ-A, other novel binding proteins, or spurious cross-linking to sterically adjacent proteins in the complex. Nevertheless, it is clear that PSQ-A is involved in the binding of the (GA)_n motif in vitro. Third, PSQ is colocalized with Pc-G protein at both ANTP-C and BX-C sites on polytene chromosomes. Fourth, there is a remarkably strong genetic interaction between Pc-G and *psq* that gives rise to leg transformation and ectopic *Ubx* expression. Finally, we show that the lack of PSQ-A in one mutant (i.e., *psq*^{Δ18}) is sufficient to account for genetic interaction with *Pc*.

Recent studies have indicated that GAF (25) or a combination of novel forms of GAF and PSQ (24) is responsible for the binding of the Pc-G complex to the (GA)_n motif. In one study, embryonic nuclear extracts were used to form the DNA-protein complex, followed by immunodetection with GAF antibody. Since multiple (GA)_n motifs are present in the probes, it is difficult to exclude the possibility that GAF and Pc-G complexes might bind these motifs independently. Similar problems also arise from subsequent studies in which fusion proteins of LexA and Pc-G have been used to bind probes containing LexA binding sites (49), since the minimal GAF binding site, the GAG trinucleotide (63), is also present in the LexA probe. Although more purified fractions were used for DNA binding analysis in the other study (24), a combination of Bio-Rex 70 and Q-Sepharose may not provide sufficient resolving power to exclude the possibility that a large number of unrelated proteins are copurified. In addition, the final fractions appear to be enriched for a GAF of ~54 kDa and to exclusively contain a PSQ of ~70 kDa. Both proteins appear substantially smaller than the smallest forms detected in the original extracts (~67 kDa for GAF and ~95 kDa for PSQ). Since both GAF and PSQ antisera have nonspecific cross-reactivities (Fig. 3B for GAF) (see reference 27 for PSQ) (data not shown), the identities of these proteins remain obscure. Nonetheless, despite these uncertainties, it is possible that GAF may play a role in certain aspects of the silencing mechanism as suggested by genetic studies (22, 24).

Other sequence-specific DNA binding factors have also been implicated for Pc-G targeting by genetic and/or biochemical studies (8, 50). PHO is the only one that has been formally categorized as a Pc-G. Its binding sites are present in many PRE (43). In addition, mutations of the PHO binding site compromise the ability of PRE to silence reporter gene in larval tissues (20). However, PHO does not appear to be directly associated with many Pc-G proteins (49). Thus, despite its important role in homeotic gene silencing, it is not clear whether PHO is directly involved in the targeting of Pc-G complexes.

Another potential candidate involved in the binding of the Pc-G complex is the Zeste protein for its copurification with Pc repression complex 1 (50). It has been speculated that Zeste proteins act as the scaffold via self-multimerization to bring together regulatory sequences situated on the same chromo-

some or different chromosomes (3). Its binding site has also been found in several PRE. In contrast to the proposed role for silencing, however, previous molecular and genetic studies have shown that the Zeste protein is most likely an activator. For example, it stimulates transcription of the *Ubx* promoter in vitro (5). Expression of a *Ubx*-LacZ transgene is completely abolished by a *zeste* mutation (39). For its transactivating effect, *zeste* has been considered a *trx-G* (34). Consistent with this notion, direct physical interaction has recently been demonstrated between the Zeste protein and two *trx-G* proteins, MOIRE and OSA, of the BRAHMA nucleosome remodeling complex (30). Genetically, *zeste* has also been defined as a transactivator involved in transvection of several genes, including *Ubx* (47). In addition, several Pc-G have been identified as suppressors of *zeste* (7, 64). These observations cast some doubts on the physiological relevance of the Zeste protein in homeotic gene silencing. It is important to note that the two best characterized PRE (i.e., *bx*d and *Fab7*) also respond to *trx-G* (11, 13, 60). Thus, the mere existence of binding sites in PRE may not necessarily provide an unambiguous indication of their functions. While the manuscript was under review, however, a recent study has shown that *zeste* mutations result in an extended expression of a *Ubx* transgene containing a replacement of the proximal promoter with a combination of multiple Zeste and NTF-1 binding sites (28), suggesting a role for *zeste* in *Ubx* silencing. However, since extended expression was also observed for a *Ubx* transgene containing multiple NTF-1 binding sites at the proximal promoter region (28), the exact role of *zeste* may need to be more thoroughly examined.

In conclusion, we believe that our results provide direct evidence that a specific PSQ isoform is critically involved in the targeting of a major Pc-G protein complex CHRASCH to the (GA)_n motifs that are commonly found in PRE. Earlier studies have demonstrated that a functional HDAC1 is associated with CHRASCH and is required for the silencing in vivo (14). We suggest a simple model for homeotic gene silencing that involves the assembly of multimeric complexes by known Pc-G proteins and other novel proteins yet to be identified, direct binding to specific sequences of PRE, and subsequent modification of N-terminal tails of core histones to establish a silencing code for stable maintenance of an inactive state.

It is also relevant to note that the functions of Pc-G silencing complexes may not be fully revealed by previous genetic or biochemical approaches because of the lack of suitable mutations, easily tractable phenotypes, or sufficient stability of the protein complexes. In the case of *psq*, a grandchildless class of mutations, sufficient amounts of PSQ remain detectable in most homozygous mutant adults (Fig. 7), yet embryos produced by these adults become severely defective before the manifestation of homeotic genes (27). In addition, the presence of more PSQ sites than Pc-G sites on polytene chromosomes suggests a much wider spectrum of target genes for PSQ. These effects altogether could conceivably obscure the homeotic effect caused by *psq* mutations, unless a more-sensitized genetic background (e.g., *Pc* mutations) is provided. The roles of MI-2 and HDAC1 in homeotic gene silencing also become apparent with similar approaches (14, 33). We speculate that some novel functions of the silencing complex may be defined by more-systematic studies.

ACKNOWLEDGMENTS

We thank the Bloomington Stock Center for fly stocks and the Developmental Studies Hybridoma Bank at the University of Iowa (under the auspices of the National Institute of Child Health and Human Development) for the PSC antibody developed by P. Adler. We are grateful to C. Berg for class I *psq* alleles and PSQ antisera, to P. Geiduschek for AB-dUTP, to M. Koelle and D. Hogness for pMK33, to T. Tsukiyama and C. Wu for GAF cDNA and antisera, and to U. Weber and M. Mlodzik for class II *psq* alleles. We also thank the following people for help at different stages of the experiments: S.-C. Lin, I.-L. Chen, Y.-H. Peng, and P.-Y. Ho. We appreciate critical comments from Ken Deen on the manuscript.

This work was supported by grants from the Academia Sinica and the National Science Council (NSC 86-2316-B-001-009, 87-2311-B-001-118, 88-2311-B-001-126, and 89-2311-B-001-081).

REFERENCES

- Bardwell, V. J., and R. Treisman. 1994. The POZ domain: a conserved protein-protein interaction motif. *Genes Dev.* 8:1664–1677.
- Bartholomew, B., R. L. Tinker, G. Kassavetis, and E. P. Geiduschek. 1995. Photochemical cross-linking assay for DNA tracking by replication proteins. *Methods Enzymol.* 262:476–494.
- Benson, M., and V. Pirrotta. 1988. The Drosophila zeste protein binds cooperatively to sites in many gene regulatory regions: implications for transvection and gene regulation. *EMBO J.* 7:3907–3915.
- Benyajati, C., L. Mueller, N. Xu, M. Pappano, J. Gao, M. Mosammamaparast, D. Conklin, D. Granok, C. Craig, and S. Elgin. 1997. Multiple isoforms of GAGA factor, a critical component of chromatin structure. *Nucleic Acids Res.* 25:3345–3353.
- Biggin, M. D., S. Bickel, M. Benson, V. Pirrotta, and R. Tjian. 1988. Zeste encodes a sequence-specific transcription factor that activates the Ultrabithorax promoter *in vitro*. *Cell* 53:713–722.
- Biggin, M. D., and R. Tjian. 1988. Transcription factors that activate the Ultrabithorax promoter in developmentally staged extracts. *Cell* 53:699–711.
- Bornemann, D., E. Miller, and J. Simon. 1998. Expression and properties of wild-type and mutant forms of the Drosophila sex comb on midleg (SCM) repressor protein. *Genetics* 150:675–686.
- Brown, J. L., D. Mucci, M. Whiteley, M.-L. Dirksen, and J. A. Kassis. 1998. The Drosophila Polycomb group gene pleiohomeotic encodes a DNA binding protein with homology to the transcription factor YY1. *Mol. Cell* 1:1057–1064.
- Bunch, T. A., Y. Grinblat, and L. S. B. Goldstein. 1988. Characterization and use of the Drosophila metallothionein promoter in cultured Drosophila melanogaster cells. *Nucleic Acids Res.* 16:1043–1061.
- Cavalli, G., and R. Paro. 1998. The Drosophila Fab-7 chromosomal element conveys epigenetic inheritance during mitosis and meiosis. *Cell* 93:505–518.
- Cavalli, G., and R. Paro. 1999. Epigenetic inheritance of active chromatin after removal of the main transactivator. *Science* 286:955–958.
- Chan, C.-S., L. Rastelli, and V. Pirrotta. 1994. A Polycomb response element in the *Ubx* gene that determines an epigenetically inherited state of repression. *EMBO J.* 13:2553–2564.
- Chang, Y.-L., B. O. King, M. O'Connor, A. Mazo, and D.-H. Huang. 1995. Functional reconstruction of *trans* regulation of the *Ultrabithorax* promoter by the products of two antagonistic genes, *trithorax* and *Polycomb*. *Mol. Cell Biol.* 15:6601–6612.
- Chang, Y.-L., Y.-H. Peng, I.-C. Pan, D.-S. Sun, B. King, and D.-H. Huang. 2001. Essential role of Drosophila Hda1 in homeotic gene silencing. *Proc. Natl. Acad. Sci. USA* 98:9730–9735.
- Chiang, A., M. O'Connor, R. Paro, J. Simon, and W. Bender. 1995. Discrete Polycomb-binding sites in each parasegmental domain of the bithorax complex. *Development* 121:1681–1689.
- Dawson, R. M. C., D. C. Elliott, W. H. Elliott, and K. M. Jones. 1986. Data for biochemical research, 3rd ed. Clarendon Press, Oxford, United Kingdom.
- DeCamillis, M., N. Cheng, D. Pierre, and H. W. Brock. 1992. The polyhomeotic gene of Drosophila encodes a chromatin protein that shares polytene chromosome-binding sites with Polycomb. *Genes Dev.* 6:223–232.
- Farkas, G., J. Gausz, M. Galloni, G. Reuter, H. Gyurkovics, and F. Karch. 1994. The *Trithorax*-like gene encodes the Drosophila GAGA factor. *Nature* 371:806–808.
- Fauvarque, M.-O., and J.-M. Dura. 1993. Polyhomeotic regulatory sequences induce developmental regulator-dependent variegation and targeted P-element insertion in Drosophila. *Genes Dev.* 7:1508–1520.
- Fritsch, C., J. L. Brown, J. K. Kassis, and J. Mueller. 1999. The DNA-binding Polycomb group protein Pleiohomeotic mediates silencing of a Drosophila homeotic gene. *Development* 126:3905–3913.
- Granok, H., B. A. Leibovitch, C. D. Shaffer, and S. C. R. Elgin. 1995. Ga-ga over GAGA factor. *Curr. Biol.* 5:238–241.
- Hagstrom, K., M. Muller, and P. Schedl. 1997. A Polycomb and GAGA

- dependent silencer adjoin the Fab-7 boundary in the *Drosophila* bithorax complex. *Genetics* **146**:1365–1380.
23. Harlow, E., and D. Lane. 1988. *Antibodies: a laboratory manual*. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.
 24. Hodgson, J. W., B. Argiropoulos, and H. W. Brock. 2001. Site-specific recognition of a 70-base-pair element containing d(GA)n repeats mediates bithoraxoid Polycomb group response element-dependent silencing. *Mol. Cell. Biol.* **21**:4528–4543.
 25. Horard, B., C. Tatout, S. Poux, and V. Pirrotta. 2000. Structure of a Polycomb response element and *in vitro* binding of Polycomb group complexes containing GAGA factor. *Mol. Cell. Biol.* **20**:3187–3197.
 26. Horowitz, H., and C. A. Berg. 1995. Aberrant splicing and transcription termination caused by P element insertion into the intron of a *Drosophila* gene. *Genetics* **139**:327–335.
 27. Horowitz, H., and C. A. Berg. 1996. The *Drosophila* pipsqueak gene encodes a nuclear BTB-domain-containing protein required early in oogenesis. *Development* **122**:1859–1887.
 28. Hur, M.-W., J. D. Laney, S.-H. Jeon, J. Ali, and M. D. Biggin. 2002. Zeste maintains repression of *Ubx* transgenes: support for a new model of Polycomb repression. *Development* **129**:1339–1343.
 29. Jurgens, G. 1985. A group of genes controlling the spatial expression of the bithorax complex in *Drosophila*. *Nature* **316**:153–155.
 30. Kal, A. J., T. Mahmoudi, N. B. Zak, and C. P. Verrijzer. 2000. The *Drosophila* Brahma complex is an essential coactivator for the trithorax group protein Zeste. *Genes Dev.* **14**:1058–1071.
 31. Kassiss, J. A. 1994. Unusual properties of regulatory DNA from the *Drosophila* engrailed gene: three “pairing-sensitive” sites within a 1.6 kb region. *Genetics* **136**:1025–1038.
 32. Kaufman, T. C., R. A. Lewis, and B. T. Wakimoto. 1980. Cytogenetic analysis of chromosome 3 in *Drosophila melanogaster*: the homeotic gene complex in polytene chromosome interval 84A-B. *Genetics* **94**:115–133.
 33. Kehle, J., D. Beuchle, S. Treuheit, B. Christen, J. A. Kennison, M. Bienz, and J. Mueller. 1998. dMi-2, a hunchback-interacting protein that functions in Polycomb repression. *Science* **282**:1897–1900.
 34. Kennison, J. A. 1995. The Polycomb and trithorax group proteins of *Drosophila*: trans-regulators of homeotic gene function. *Annu. Rev. Genet.* **29**:289–303.
 35. Kennison, J. A., and J. W. Tamkun. 1988. Dosage-dependent modifiers of Polycomb and Antennapedia mutations in *Drosophila*. *Proc. Natl. Acad. Sci. USA* **85**:8136–8140.
 36. Kerrigan, L. A., G. E. Croston, L. M. Lira, and J. T. Kadonaga. 1991. Sequence-specific transcriptional anti-repression of the *Drosophila* Kruppel gene by the GAGA factor. *J. Biol. Chem.* **266**:574–582.
 37. Koelle, M. R., W. S. Talbot, W. A. Segraves, M. T. Bender, P. Cherbas, and D. S. Hogness. 1991. The *Drosophila* EcR gene encodes an ecdysone receptor, a new member of the steroid receptor superfamily. *Cell* **67**:59–77.
 38. Kyba, M., and H. W. Brock. 1998. The *Drosophila* Polycomb group protein Psc contacts ph and Pc through specific conserved domains. *Mol. Cell. Biol.* **18**:2712–2720.
 39. Laney, J. D., and M. D. Biggin. 1997. Zeste-mediated activation by an enhancer is independent of cooperative DNA binding *in vivo*. *Proc. Natl. Acad. Sci. USA* **15**:3602–3604.
 40. Lehmann, M., T. Siegmund, K. Lintermann, and G. Korge. 1998. The Pipsqueak protein of *Drosophila melanogaster* binds to GAGA sequences through a novel DNA-binding domain. *J. Biol. Chem.* **273**:28504–28509.
 41. Lewis, E. B. 1978. A gene complex controlling segmentation in *Drosophila*. *Nature* **276**:565–570.
 42. McGinnis, W., and R. Krumlauf. 1992. Homeobox genes and axial patterning. *Cell* **68**:283–302.
 43. Mihaly, J., R. K. Mishra, and F. Karch. 1998. A conserved sequence motif in Polycomb-response elements. *Mol. Cell* **1**:1065–1066.
 44. Ng, J., C. M. Hart, K. Morgan, and J. A. Simon. 2000. A *Drosophila* ESC-E(Z) protein complex is distinct from other Polycomb group complexes and contains covalently modified ESC. *Mol. Cell. Biol.* **20**:3069–3078.
 45. Peterson, A., M. Kyba, D. Borneman, K. Morgan, H. Brock, and J. A. Simon. 1997. A domain shared by the *Polycomb* group proteins Scm and ph mediates heterotypic and homotypic interactions. *Mol. Cell. Biol.* **17**:6683–6692.
 46. Pirrotta, V. 1997. Chromatin silencing mechanisms in *Drosophila* maintain patterns of gene expression. *Trends Genet.* **13**:314–318.
 47. Pirrotta, V. 1991. The genetics and molecular biology of zeste in *Drosophila melanogaster*. *Adv. Genet.* **29**:301–348.
 48. Pirrotta, V. 1998. Polycombing the genome: Pc-G, trx-G and chromatin silencing. *Cell* **93**:333–336.
 49. Poux, S., D. McCabe, and V. Pirrotta. 2001. Recruitment of components of Polycomb group chromatin complexes in *Drosophila*. *Development* **128**:75–85.
 50. Saurin, A. J., Z. Shao, H. Erdjument-Bromage, P. Tempst, and R. E. Kingston. 2001. A *Drosophila* Polycomb group complex includes zeste and dTFII proteins. *Nature* **412**:655–660.
 51. Shao, Z., F. Raible, R. Mollaaghababa, J. R. Guyon, C.-T. Wu, W. Bender, and R. E. Kingston. 1999. Stability of chromatin structure by PRC1, a Polycomb complex. *Cell* **98**:37–46.
 52. Siegel, V., T. A. Jongens, L. Y. Jan, and Y. N. Jan. 1993. Pipsqueak, an early acting member of the posterior group of genes, affects vasa level and germ cell-somatic cell interaction in the developing egg chamber. *Development* **119**:1187–1202.
 53. Simon, J., A. Chiang, W. Bender, M. J. Shimell, and M. O'Connor. 1993. Elements of the *Drosophila* Bithorax complex that mediate repression by Polycomb group products. *Dev. Biol.* **158**:131–144.
 54. Simon, J., M. Peifer, W. Bender, and M. O'Connor. 1990. Regulatory elements of the bithorax complex that control expression along the anterior-posterior axis. *EMBO J.* **9**:3945–3956.
 55. Soeller, W. C., W. E. Oh, and T. B. Kornberg. 1993. Isolation of cDNAs encoding the *Drosophila* GAGA transcription factor. *Mol. Cell. Biol.* **13**:7961–7970.
 56. Soto, M. C., T. B. Chou, and W. Bender. 1995. Comparison of germ-line mosaics of genes in the Polycomb group of *Drosophila melanogaster*. *Genetics* **140**:231–243.
 57. Strutt, H., G. Cavalli, and R. Paro. 1997. Co-localization of Polycomb protein and GAGA factor on regulatory elements responsible for the maintenance of homeotic gene expression. *EMBO J.* **16**:3621–3632.
 58. Strutt, H., and R. Paro. 1997. The Polycomb group protein complex of *Drosophila melanogaster* has different compositions at different target genes. *Mol. Cell. Biol.* **17**:6773–6783.
 59. Tie, F., T. Furuyama, J. Prasad-Sinha, E. P. Jane, and P. J. Harte. 2001. The *Drosophila* Polycomb group proteins ESC and E(Z) are present in a complex containing the histone-binding protein p55 and the histone deacetylase RPD3. *Development* **128**:275–286.
 60. Tillib, S., S. Petruk, Y. Sedkov, A. Kuzin, M. Fujioka, T. Goto, and A. Mazo. 1999. Trithorax- and Polycomb-group response elements within an *Ultra-bithorax* transcription maintenance unit consist of closely situated but separable sequences. *Mol. Cell. Biol.* **19**:5189–5202.
 61. Weber, U., V. Siegel, and M. Mlodzik. 1995. Pipsqueak encodes a novel nuclear protein required downstream of seven-up for the development of photoreceptors R3 and R4. *EMBO J.* **14**:6247–6257.
 62. White, R. A. and M. Wilcox. 1984. Protein products of the Bithorax complex in *Drosophila*. *Cell* **39**:163–171.
 63. Wilkins, R. C., and J. T. Lis. 1998. GAGA factor binding to DNA via a single trinucleotide sequence element. *Nucleic Acids Res.* **1**:2672–2678.
 64. Wu, C., R. S. Jones, P. F. Lasko, and W. M. Gelbart. 1989. Homeosis and the interaction of zeste and white in *Drosophila*. *Mol. Gen. Genet.* **218**:559–564.
 65. Zink, B., and R. Paro. 1989. *In vivo* binding of a trans-regulator of homeotic genes in *Drosophila melanogaster*. *Nature* **337**:468–471.