

Ytm1, Nop7, and Erb1 Form a Complex Necessary for Maturation of Yeast 66S Preribosomes

Tiffany D. Miles, Jelena Jakovljevic, Edward W. Horsey,[†] Piyanun Harnpicharnchai,[‡]
 Lan Tang, and John L. Woolford, Jr.*

Department of Biological Sciences, Carnegie Mellon University, Pittsburgh, Pennsylvania 15213

Received 2 March 2005/Returned for modification 29 March 2005/Accepted 20 September 2005

The essential, conserved yeast nucleolar protein Ytm1 is one of 17 proteins in ribosome assembly intermediates that contain WD40 protein-protein interaction motifs. Such proteins may play key roles in organizing other molecules necessary for ribosome biogenesis. Ytm1 is present in four consecutive 66S preribosomes containing 27SA₂, 27SA₃, 27SB, and 25.5S plus 7S pre-rRNAs plus ribosome assembly factors and ribosomal proteins. Ytm1 binds directly to Erb1 and is present in a heterotrimeric subcomplex together with Erb1 and Nop7, both within preribosomes and independently of preribosomes. However, Nop7 and Erb1 assemble into preribosomes prior to Ytm1. Mutations in the WD40 motifs of Ytm1 disrupt binding to Erb1, destabilize the heterotrimer, and delay pre-rRNA processing and nuclear export of preribosomes. Nevertheless, 66S preribosomes lacking Ytm1 remain otherwise intact.

Biogenesis of eukaryotic ribosomes is a highly regulated and dynamic process that begins in the nucleolus with transcription of a precursor rRNA (pre-rRNA) that is rapidly packaged into the 90S ribonucleoprotein particle containing ribosomal proteins, nonribosomal proteins, and snoRNA-containing ribonucleoprotein particles (snoRNPs). The 90S pre-RNPs are converted into 43S and 66S ribosome assembly intermediates, which ultimately give rise to mature 40S and 60S ribosomal subunits (Fig. 1).

Molecular genetic approaches in yeast identified more than 70 *trans*-acting factors required for ribosome assembly (12, 14, 46). Subsequent advances in proteomics enabled purification of pre-rRNPs from yeast and identification of an additional 80 assembly factors present in preribosomes, as well as most of those proteins previously discovered using genetic screens (3, 7, 11, 17, 20, 21, 24, 26, 37, 38, 41, 49–51). Metazoan homologues of most of the yeast ribosome assembly factors were discovered by proteomic analysis of purified nucleoli (2, 52).

Among the assembly factors found in yeast preribosomes are 17 proteins containing WD40 motifs (14). These motifs function as protein-protein interaction domains (53). Therefore, such WD40-containing proteins may nucleate assembly of preribosomes by interacting sequentially or simultaneously with other assembly factors or ribosomal proteins. Previously, we identified the WD40 protein Ytm1 as a constituent of purified 66S pre-rRNPs and showed that depletion of Ytm1 results in a deficiency of 60S ribosomal subunits (21).

In this study, we have further investigated the role of Ytm1 in ribosome biogenesis. Ytm1 is a constituent of multiple con-

secutive 66S preribosomes containing 27SA₂, 27SA₃, 27SB, 25.5S, and 7S pre-rRNAs plus a collection of ribosomal and nonribosomal proteins. Ytm1 is present in a heterotrimer with two other assembly factors, Nop7 and Erb1, both within 66S pre-rRNPs and as a subcomplex independent of preribosomes. Mutations in Ytm1 disrupt interactions between Ytm1 and Erb1, destabilize the heterotrimer, and significantly reduce association of these three proteins with 66S preribosomes. These 66S pre-rRNPs otherwise remain intact in the *ytm1-1* mutant, but processing of 27SA₃ pre-rRNAs is delayed and release of 66S preribosomes from the nucleolus is partially blocked. Thus, Ytm1 is necessary to nucleate the assembly of a heterotrimer that is important for intermediate-to-late steps in maturation of 66S preribosomes.

MATERIALS AND METHODS

Strains, plasmids, and media. Yeast strains used in this work (Table 1) were grown in YEPD medium (2% dextrose, 2% peptone, and 1% yeast extract) or YEPGal medium (2% galactose, 2% peptone, and 1% yeast extract) at 30°C and harvested at 5×10^7 cells/ml unless otherwise indicated. The *ytm1-1* mutant strain JWY7128 was generated by mutagenizing plasmid pRS317 containing wild-type *YTM1* and *LYS2* with hydroxylamine and transforming it into yeast strain SM412 cells which have *GAL-YTM1* at the *YTM1* locus. The *ytm1-1* mutant plasmid that conferred temperature sensitivity to strains grown on selective medium containing 2% glucose was rescued, shuttled through *Escherichia coli*, and used to replace the chromosomal *YTM1* gene in strain JWY7132 to generate *ytm1-1* strain JWY7128. The *GAL-YTM1* strain JWY6992 was described previously (21).

Three-hemagglutinin epitope (HA3)- and tandem affinity purification (TAP)-tagged strains were generated as described in references 21 and 26. Integration of the HA3 tag or the TAP cassette in-frame with the last codon of each open reading frame was confirmed by genomic PCR and Western immunoblotting. Yeast strain JWY6790 expressing enhanced green fluorescent protein (eGFP)-tagged rPL25 was generated by transforming JWY7128 (*ytm1-1*) with a plasmid bearing rPL25eGFP (28). Transformants were screened by fluorescence microscopy for expression of rPL25eGFP. Yeast strains JWY7808 (*GST-YTM1*), JWY7810 (*GST-ERB1*) and JWY7807 (*GST-NOP7*) expressing glutathione S-transferase (GST) fusion proteins were obtained from Mike Snyder (Yale University).

Sucrose gradient analysis. Ribosomes, preribosomes, and polyribosomes in yeast strains JWY3400 (*YTM1*), JWY7124 (*YTM1-TAP*), and JWY7128 (*ytm1-1*) were analyzed as described previously (26). JWY7128 was grown at 25°C and harvested or else shifted from 25 to 37°C for 3 h and harvested. Ytm1-HA3 or

* Corresponding author: Department of Biological Sciences, Carnegie Mellon University, 616 Mellon Institute, 4400 Fifth Ave., Pittsburgh, PA 15213. Phone: (412) 268-3193. Fax: (412) 268-7129. E-mail: jw17@andrew.cmu.edu.

[†] Present address: Department of Chemical Engineering, Carnegie Mellon University, 700 Technology Drive, PTC Room 4305, Pittsburgh, PA 15219.

[‡] Present address: National Center for Genetic Engineering and Biotechnology, Thailand Science Park, 113 Paholyothin Road, Klong 1, Klong Luang, Pathumthani 12120, Thailand.




FIG. 1. Pre-rRNA processing and pre-rRNP maturation pathway in *Saccharomyces cerevisiae*. (A) The 35S pre-rRNA contains sequences for mature 18S, 5.8S, and 25S rRNAs (represented as thick horizontal lines) along with additional internal and external spacer sequences (represented as thin horizontal lines). The 35S pre-rRNA is transcribed by RNA polymerase I and rapidly modified and processed to produce 33S pre-rRNA. Cleavage of 33S pre-rRNA at site A_0 generates 32S pre-rRNA. The 20S and 27SA₂ pre-rRNA processing intermediates are generated through internal cleavage of 32S pre-rRNA at the A_2 site. Subsequent processing and cleavage of 20S and 27SA₂ pre-rRNAs result in the production of the mature 18S, 25S, and 5.8 rRNAs, respectively. 5S rRNA is transcribed separately by RNA polymerase III. (B) Pre-rRNA processing occurs in preribosomal particles. The 35S primary transcript is found within the 90S pre-rRNP (dark gray circle). Cleavage at site A_2 initiates subunit-specific maturation, generating the 43S and 66S pre-rRNPs (light gray circles). The 43S preribosome is exported to the cytoplasm, where final steps in 20S maturation take place. Multiple 66S preribosomes exist that contain each of the 27S or 25.5S plus 7S pre-rRNA processing intermediates. The mature 40S subunit contains 18S rRNA, whereas the 60S subunit contains 25S, 5.8S, and 5S rRNA (white circles).

Ytm1-TAP were identified in gradient fractions by Western blotting using mouse monoclonal antibody 12CA5 or rabbit anti-mouse immunoglobulin G (Pierce), respectively.

Analysis of rRNA. Steady-state levels of rRNAs were analyzed by Northern blotting (26) or primer extension assays (58). To carry out primer extension, radiolabeled oligonucleotide primers complementary to 35S, 27S, or 25.5S pre-rRNA were first annealed for 90 min at 46°C to total RNA or affinity-purified rRNA. Primer extension reaction mixtures containing the annealed oligonucleotide primer/RNA hybrid, 10 mM deoxynucleoside triphosphates (dNTPs) (Amersham Biosciences), 12.5 U avian myeloblastosis virus reverse transcriptase (Promega), and 20 U RNasin (Promega) were incubated for 40 min at 46°C. To hydrolyze the RNA, 6 μ l of 1 M NaOH and 1 μ l of 0.5 M EDTA were added to each primer extension reaction mixture for 30 min to 1 h at 55°C. Next, 6 μ l of 1 M HCl was added to each reaction mixture and DNA was precipitated with 4 μ g glycogen, 30 μ l 7.5 M NH₄OAc, and 250 μ l of 100% ethanol. DNA was suspended in DNA dye (95% formamide, 0.05% xylene cyanol, 0.05% bromophenol blue in 20 mM EDTA, pH 8.0) and subjected to electrophoresis on a 6% polyacrylamide-urea sequencing gel. Gels were dried and directly exposed to film for autoradiography. Oligonucleotide sequences are available upon request. Pulse-chase assays of pre-rRNA processing were carried out as previously described (26).

Ribosome export assays. Release of preribosomes from nucleoli and export from the nucleoplasm to the cytoplasm were assayed as previously described

(28), except that strains were grown overnight in C-Trp medium, washed and suspended in YEPD medium, and grown at 25°C or shifted to 37°C for 5 h.

Affinity purification and mass spectrometry. Cell extracts were prepared, tandem affinity purification was carried out, and identification of copurifying proteins and RNAs was performed as previously described (26, 47).

Ribosome assembly subcomplexes were separated from 66S preribosomes and 60S ribosomal subunits by centrifugation on 7% to 47% sucrose gradients (26) or by differential centrifugation as performed previously (32) with the following modifications: whole-cell extracts were centrifuged for 2 h at 180,000 $\times g$ at 4°C, followed by a second centrifugation of supernatants for 30 to 45 min at 180,000 $\times g$ at 4°C. Subcomplexes were affinity purified from gradient fractions or from the 180,000 $\times g$ supernatant using TAP-tagged Nop7.

Assembly subcomplexes were isolated directly from whole-cell extracts by adding to the lysis buffer and calmodulin binding buffer a phosphatase inhibitor cocktail (20 mM pyrophosphate, 10 mM sodium azide, 20 mM sodium fluoride, 1 mM sodium orthovanadate, and 100 mM β -glycerophosphate) that disrupts pre-rRNPs.

Generation of anti-Ytm1 antibodies and Western immunoblotting. Rabbit antibodies generated against the synthetic peptide ITREDKSVQKGVNDK (Alpha Diagnostics, Inc.) were used to detect Ytm1. Antibodies were concentrated by ammonium sulfate precipitation, dialyzed, and affinity purified using full-length filter-bound Ytm1 protein previously subjected to electrophoresis through

TABLE 1. Strains used in this study

Strain	Genotype	Source or reference
JWY3400	<i>MATa ura3-52 lys2-801 trp1-1 leu2-1 his3-Δ200 pep4::HIS3 prb1-Δ1.6R can YTM1</i>	E. Jones
JWY7128	<i>MATα ura3-52 lys2-801 ade2-101 trp1-Δ63 his3-Δ200 leu2-Δ1 ytm1-1 URA3</i>	S. Matsumoto
JWY7132	<i>MATα ura3-52 lys2-801 ade2-101 trp1-Δ63 his3-Δ200 leu2-Δ1 YTM1</i>	S. Matsumoto
JWY6149	<i>MATa his3-Δ1 leu2-Δ0 lys2-Δ0 MET15 ura3-Δ0 YTM1</i>	26
JWY6992	<i>MATα met15-Δ0 ura3-Δ0 his3-Δ1 leu2-Δ0 YTM1::kanMX4 pGAL-YTM1 URA3</i>	21
JWY6779	<i>MATa ura3-52 trp1-Δ101 lys2-801 his3-Δ200 leu2-Δ1 YTM1 p1877 (RPL25eGFP URA3)</i>	This study
JWY6790	<i>MATα ura3-52 lys2-801 ade2-101 trp1-Δ63 his3-Δ200 leu2-Δ1 GAL1 ytm1-1 p1878 (RPL25-eGFP TRP1)</i>	This study
JWY6147	<i>MATa ura3-52 trp1-Δ101 lys2-801 his3-Δ200 leu2-Δ1 YTM1</i>	E. Jones
JWY6770	<i>MATa ura3-52 trp1-Δ101 lys2-801 his3-Δ200 leu2-Δ1 YTM1-HA3 HIS3</i>	This study
JWY7124	<i>MATa ura3-52 lys2-801 trp1-1 leu2-1 his3-Δ200 pep4::HIS3 prb1-Δ1.6R can YTM1-TAP TRP1</i>	This study
JWY7129	<i>MATα ura3-52 lys2-801 trp1 leu2-1 his3-Δ200 ade2-101 pep4::HIS3 prb1-Δ1.6R can 1 nop4::TRP1 [pRS315 LEU2 nop4-3] YTM1-TAP URA3</i>	This study
JWY6729	<i>MATα ura3-52 lys2-801 trp1 leu2-1 his3-Δ200 ade2-101 pep4::HIS3 prb1-Δ1.6R can1 nop4::TRP1 [pRS315 LEU2 nop4-3] NOP7-TAP TRP1</i>	21
JWY7131	<i>MATa his7 ade2 trp1 lys2 tyr1 gal1 gal2 rrp1-1 YTM1-TAP TRP1</i>	This study
JWY6970	<i>MATa his7 ade2 trp1 lys2 tyr1 gal1 gal2 rrp1-1 NOP7-TAP TRP1</i>	This study
JWY6938	<i>MATa ura3-52 lys2-801 trp1-1 leu2-1 his3-Δ200 pep4::HIS3 prb1-Δ1.6R can YTM1 NOP7-TAP TRP1</i>	21
JWY7019	<i>MATα ura3-52 lys2-801 ade2-101 trp1-Δ63 his3-Δ200 leu2-Δ1 ytm1-1 NOP7-TAP TRP1</i>	This study
JWY7810	<i>MATa pep4-3 his4-500 ura3-52 leu2-3,112 pEG(KG) (GST-ERB1 LEU2 URA3)</i>	M. Snyder
JWY7808	<i>MATa pep4-3 his4-500 ura3-52 leu2-3,112 pEG(KG) (GST-YTM1 LEU2 URA3)</i>	M. Snyder
JWY7807	<i>MATa pep4-3 his4-500 ura3-52 leu2-3,112 pEG(KG) (GST-NOP7 LEU2 URA3)</i>	M. Snyder
JWY6300	<i>MATα trp1-901 leu2-3,112 ura3-52 his3-Δ200 gal4Δ gal80Δ LYS2-GAL1-HIS3 ade2-101 GAL2-ADE2 met2::GAL7-LacZ</i>	PJ69-4α (29)
JWY4340	<i>MATa trp1-901 leu2-3,112 ura3-52 his3-Δ200 gal4 gal80 LYS2::GAL1-HIS3 GAL2-ADE2 met2::GAL7-lacZ</i>	PJ69-4a (29)
JWY7138	<i>MATa ura3-52 lys2-801 trp1-1 leu2-1 his3-Δ200 pep4::HIS3 prb1-Δ1.6R can BRX1-TAP TRP1 YTM1</i>	This study
JWY7140	<i>MATα ura3-52 lys2-801 ade2-101 trp1Δ63 his-3Δ200 leu2-Δ1 BRX1-TAP TRP1 ytm1-1</i>	This study

a 10% polyacrylamide gel and electroblotted to nitrocellulose (Optitran; Schleicher and Schuell). Immunoblotting was carried out using standard protocols (26).

GST pull-down assays. GST fusion proteins were harvested from yeast by glass bead lysis of frozen cell pellets suspended in 1.6 ml sorbitol buffer (300 mM sorbitol, 5 mM MgCl₂, 100 mM NaCl, 10 mM Tris-HCl, pH 7.5, 1 mM phenylmethylsulfonyl fluoride, 1 μg/ml aprotinin, 1 μg/ml pepstatin, 1 μg/ml leupeptin). One milliliter of protein extract was incubated with 50 μl glutathione-agarose beads overnight at 4°C. Beads were washed three times with 1 ml high-salt wash buffer (300 mM sorbitol, 5 mM MgCl₂, 1 M NaCl, 10 mM Tris-HCl, pH 7.5), three times with 1 ml HKT buffer (10 mM HEPES, 100 mM KCl, 0.5% Triton X, 1% IGEPAL, 5% bovine serum albumin) and once with 1 ml sorbitol buffer lacking proteinase inhibitors. ³⁵S-labeled Nop7, Erb1, and Ytm1 were synthesized in vitro using the TNT T7 Quick for PCR DNA kit (Promega Corporation) and oligonucleotides T7_NOP7_UP and NOP7_TRUC_2HY_GAP_DN, T7_ERB1_UP and ERB1_DN, or T7_YTM1_UP and YTM1-2HYB-GAP_REPR-DN. Labeled proteins (5 μl or 10% of the labeling reaction) were incubated with the glutathione bead-bound GST fusion proteins for 2 h at 4°C. GST beads were washed once with 0.5 ml of HKT buffer containing 1 M NaCl and three times with 1 ml HKT buffer. Protein complexes were eluted from GST beads by boiling in 35 μl of 2× sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) sample buffer and subjected to electrophoresis on 10% polyacrylamide gels. Gels were fixed in a 50% methanol-15% acetic acid solution for 15 min, washed with water, treated with 1 M sodium salicylate for 1 h, dried, and subjected to autoradiography.

Yeast two-hybrid assays. The *YTM1* and *ERB1* open reading frames were amplified from the genome using primer pairs YTM1-2HYB-GAP_REPR-UP and YTM1-2HYB-GAP_REPR-DN or ERB1-2HYB-GAP_REPR-UP and ERB1-2HYB-GAP_REPR-DN, respectively. PCR products were cloned into the DNA binding domain vector pOBD-2 by gap repair and transformed into the two-hybrid host strain PJ69-4α (JWY6300). Yeast cells containing *NOP7*, *ERB1*, and *YTM1* cloned into the DNA activation domain vector (pOAD) were obtained as a pool in the two-hybrid host strain PJ69-4a (JWY4340) from the Yeast Resource Center (University of Washington). The yeast two-hybrid assay was performed as previously described (5). Diploids containing activation domain and binding domain plasmids were screened on media lacking adenine or containing 2 to 50 mM 3-aminotriazole to assay the *ADE3* and *HIS3* reporter genes, respectively.

RESULTS

The *ytm1-1* mutant contains mutations in two of the WD40 repeat motifs of Ytm1. Ytm1 is an essential 51-kDa yeast nucleolar protein that is highly conserved from fungi to humans. Seven WD repeats are clustered at the C terminus of the protein (Fig. 2A). WD repeat-containing proteins are predicted to share a common circularized beta-propeller structure (Fig. 2B) based on the crystal structure of the Gβ subunit of heterotrimeric G proteins (16, 33, 54, 59). A paramount feature of WD repeat proteins is their ability to participate in multiple interactions either simultaneously or sequentially. The WD repeat motif is not known to have any enzymatic activity. Therefore, WD repeat proteins acting in disparate pathways may share a common function of establishing and regulating interactions within multiprotein complexes.

To investigate the importance of WD40 motifs in Ytm1, we used the *ytm1-1* temperature-sensitive mutant that contains two mutations, G398D and S442N, in WD40 repeats 6 and 7 of Ytm1 (Fig. 2A). This mutant strain was generated by random mutagenesis of cloned *YTM1* and replacement of chromosomal *YTM1* with the mutant allele. The counterpart of Ytm1 residue G398 in the Gβ protein contacts phosphatidylinositol (54). Thus, residues at this position may be important for mediating interactions with associated proteins. Within the conserved WD40 repeat consensus sequence is the structural tetrad, which contributes to local and global stability through inter- and intrablock hydrogen bonding (40, 53, 54). The G398D substitution occurs in the position adjacent to the highly conserved aspartate residue in the structural tetrad. Similarly, the S442N substitution occurs in a residue that is next to the serine/threonine




FIG. 2. Ytm1 is a WD40 repeat-containing protein. (A) Predicted amino acid sequence of *S. cerevisiae* Ytm1. WD40 repeats are overlined. Amino acid residues altered in the *ytm1-1* mutant are indicated by asterisks. (B) Ras Mol 2.6 was used to generate the top and side view of a model for amino acids 103 to 450 of Ytm1, based on the crystal structure of the WD repeat protein G β .

residue of the structural tetrad. Thus, the ability of Ytm1 to bind to ligands may be compromised in the *ytm1-1* mutant, which may result in defects in ribosome assembly.

The *ytm1-1* mutant was unable to grow at 37°C and grew slower than the wild-type control strain at all temperatures below 37°C (data not shown). The G398D and S442N mutations might inactivate Ytm1 at 37°C, perhaps by distorting the structure of Ytm1 or by disrupting interactions with ligands of Ytm1. The *ytm1-1* mutant protein is relatively stable: amounts of Ytm1-1 protein did not change drastically when compared

to those of wild-type Ytm1 in strains grown at 25°C or shifted from 25°C to 37°C for 5 h (data not shown).

The *ytm1-1* mutant is deficient in 60S ribosomal subunits. To determine the effect of inactivation of Ytm1 on ribosome biogenesis, we assayed levels of ribosomal subunits, monoribosomes, and polyribosomes in *ytm1-1* cells grown at 25°C or shifted from 25°C to 37°C for 3 h. Upon shifting the *ytm1-1* mutant cells to 37°C, amounts of 60S ribosomal subunits and 80S monosomes were greatly reduced and half-mer polyribosomes were apparent. At 25°C, *ytm1-1* mutant cells contained fewer free 60S subunits and 80S monoribosomes and accumulated half-mer polyribosomes, compared to wild-type *YTM1* cells (Fig. 3). This suggests that the function of Ytm1-1 is compromised even at the permissive temperature and is consistent with the slow-growth phenotype at 25°C. These findings are consistent with previous results obtained when Ytm1 is metabolically depleted (21).

Pre-rRNA processing is slowed when Ytm1 is inactivated or depleted. The kinetics of pre-rRNA processing were analyzed by pulse-labeling *YTM1* wild-type and *ytm1-1* mutant cells, grown at 25°C, and shifted to 37°C for 5 h, as well as the *GAL-YTM1* cells shifted from galactose- to glucose-containing medium, in which Ytm1 is depleted. Processing of 35S pre-rRNA to mature 25S and 18S rRNAs occurred rapidly in the *YTM1* strain. The 27S pre-rRNA processing intermediate was completely converted to 25S rRNA by the 60-min chase point. In the *ytm1-1* mutant, however, 27S pre-rRNA was still present at the 60-min chase point; 27SA pre-rRNAs were converted to 27SB pre-rRNA less efficiently than in wild-type cells (Fig. 4B). Consequently, 3.6-fold less mature 25S rRNA was produced relative to 18S rRNA in the *ytm1-1* mutant than in the wild-type cells (Fig. 4B, compare lanes 5 and 10). Both 35S pre-rRNA and 23S pre-rRNA accumulated in the *ytm1-1* mutant. The effects on 35S and 23S are thought to be indirect and often are observed when 60S ribosomal subunit assembly is perturbed (4, 19). Effects on pre-rRNA processing observed in Ytm1-depleted cells are identical to those found in the *ytm1-1* mutant (data not shown).

We analyzed the steady-state levels of rRNA intermediates and mature rRNAs by primer extension and Northern blotting




FIG. 3. The *ytm1-1* mutant is deficient in 60S ribosomal subunits. Free 40S and 60S ribosomal subunits, monoribosomes, and polyribosomes were assayed in yeast strains JWY3400 (*YTM1*) (left) or JWY7128 (*ytm1-1*) (center) grown at 25°C or JWY7128 grown at 25°C and shifted to 37°C for 3 h (right). Whole-cell extracts prepared from each strain were fractionated on 7 to 47% sucrose gradients. A_{260} peaks representing 40S and 60S ribosomal subunits and 80S monosomes are labeled. Half-mer polyribosomes are indicated by vertical arrows.

(Fig. 4C and D). Ytm1 was depleted from *GAL-YTM1* cells by shifting galactose-grown cells to glucose for 0 h, 10 h, 12 h, 15 h, or 18 h. The *ytm1-1* cells were grown at 25°C and shifted to 37°C for 0 h, 3 h, or 6 h. Primer extension using an oligonucleotide that detects 27S pre-rRNAs (Fig. 4A) indicated that under nonpermissive conditions amounts of 27SA₂ pre-rRNA decrease in *ytm1-1* cells but increase in *GAL-YTM1* cells (Fig. 4C). 27SA₃ pre-rRNA strongly accumulated upon shifting *GAL-YTM1* and *ytm1-1* cells to the nonpermissive conditions (Fig. 4C). Amounts of 27SB pre-rRNAs were also affected: 27SB_S pre-rRNA was drastically diminished relative to 27SB_L pre-rRNA, making these species nearly equal in Ytm1-depleted or Ytm1-inactivated cells (Fig. 4C).

Phosphorimage analysis of Northern blots also indicated that 7S pre-rRNA and 5S rRNA decrease slightly upon shifting *GAL-YTM1* and *ytm1-1* cells to nonpermissive conditions (Fig. 4D). Despite the observed effects on processing of early and intermediate pre-rRNAs present in 66S pre-rRNPs, amounts of 25S and 5.8S rRNA were largely unaffected. Because preexisting 25S and 5.8S rRNAs are present in vast quantities, changes in steady-state amounts of 25S and 5.8S rRNAs may be masked and therefore difficult to observe by standard assays.

Taken together, these results indicate that in *ytm1* mutants pre-rRNA processing delays begin early, at the step when 27SA₂ pre-rRNA is converted to 27SA₃ pre-rRNA. Subsequent steps in pre-rRNA processing are similarly slowed down, but no step in pre-rRNA processing is completely blocked. Mutation of Ytm1 through depletion or inactivation results in nearly identical phenotypes, suggesting that Ytm1-1 protein is largely inactive.

Ytm1 is necessary for release of 66S preribosomes from the nucleolus. To further investigate the timing and role of Ytm1 in ribosome biogenesis, we assayed the ability of 60S preribosomes to exit the nucleolus and nucleus in the *ytm1-1* mutant, using the ribosome export assay (28), in which eGFP-tagged rpL25 functions as a reporter. In *ytm1-1* cells grown at the permissive temperature, 66S preribosomes were released to the cytoplasm (Fig. 5C). In wild-type cells grown at 25°C or shifted from 25°C to 37°C, rpL25eGFP signal was cytoplasmic (data not shown). When the *ytm1-1* mutant strain was grown at 25°C and shifted to 37°C for 5 h, rpL25eGFP was strongly retained in the nucleolus in most cells (Fig. 5D, arrows), although in some cells signal was distributed throughout the nucleoplasm. Thus, Ytm1 is important for nucleolar release of 66S preribosomes and perhaps for subsequent nuclear export.

Ytm1 is a component of 66S preribosomes. Previously Ytm1 was identified in 66S pre-rRNPs purified using TAP-tagged ribosome assembly factors (3, 21, 26, 41). Consistent with this result, HA3-tagged Ytm1 peaks in sucrose gradient fractions 15 to 17 containing 66S preribosomes (21) (Fig. 6). A small amount of Ytm1 can be detected sedimenting at the top of the gradient in lighter fractions and larger amounts at the bottom of the gradient in heavier fractions. The significance of the sedimentation in heavier fractions is unclear.


To examine in more detail ribosome assembly intermediates containing Ytm1, we identified the pre-rRNAs and proteins associated with affinity-purified Ytm1-TAP. TAP-tagged Ytm1 is fully functional: the tagged strain grows at wild-type rates and has a wild-type polysome profile, and Ytm1-TAP sediments on sucrose gradients with a peak at 66S (data not shown).

To determine in which preribosomes Ytm1 is present, we assayed which pre-rRNAs copurify with TAP-tagged Ytm1. The amounts of 27SA₂, 27SA₃, 27SB, 25.5S, and 7S pre-rRNAs recovered relative to each other were similar to those found in whole cells. Smaller relative amounts of 5.8S rRNA and no 35S or 20S pre-rRNA or 18S rRNA copurified with Ytm1-TAP (Fig. 7). Enrichment of Ytm1 with these RNA molecules is consistent with our finding that Ytm1 is important for assembly of 60S ribosomal subunits and for processing of 27S pre-rRNA (Fig. 3 and 4).


Fifty-three different proteins that copurify with TAP-tagged Ytm1 were identified by SDS-PAGE and mass spectrometry (Fig. 8). Among these are ribosomal proteins from both the large and small ribosomal subunits. Since these ribosomal proteins frequently contaminate TAPs (3, 11, 21), their significance cannot be assessed. Seventeen nonribosomal proteins specifically required for biogenesis of 60S ribosomal subunits are present in the Ytm1-containing particles (Fig. 8 and Table 2). Copurification of these pre-rRNAs and proteins with TAP-tagged Ytm1 indicates that Ytm1 first enters the ribosome assembly pathway by associating with 66S preribosomes and remains stably associated with 66S pre-rRNPs until late stages of maturation in the nucleoplasm.

Ribosome assembly factors Ytm1, Nop7, and Erb1 are present together in a subcomplex. Several different experiments demonstrate that Ytm1, Erb1, and Nop7 form a heterotrimeric subcomplex that is present within 66S preribosomes and also exists independently of pre-rRNPs. (i) Affinity purification from whole-cell extracts using TAP-tagged Ytm1, Nop7, or Erb1 yielded 50 to 60 proteins present in 66S pre-rRNPs (Fig. 8) (21; data not shown). In each case, greater amounts of Ytm1, Nop7, and Erb1 (Fig. 8, bands 33, 37, and 40, respectively) were recovered than those of any of the other proteins. This suggests that a mixture of 66S pre-rRNPs and a heterotrimer of Ytm1, Nop7, and Erb1 copurify with each of these TAP-tagged proteins (Fig. 8 and see Fig. 11). (ii) Affinity purification using TAP-tagged Nop7 or Ytm1 from *rrp1-1* or *nop4-3* mutants in which 66S pre-rRNPs are unstable (21, 26) yielded mostly Ytm1, Nop7, and Erb1, and greatly diminished amounts of molecules comprising 66S pre-rRNPs (21) (Fig. 9A). Thus, under these conditions, many fewer 66S pre-rRNPs were recovered, but the heterotrimeric subcomplexes remained intact. (iii) The Ytm1/Nop7/Erb1 heterotrimer could be separated from 66S pre-rRNPs by sucrose gradient fractionation or differential centrifugation of whole-cell extracts. Affinity purification from such enriched fractions using TAP-tagged Ytm1, Nop7, or Erb1 yielded primarily Ytm1, Nop7, and Erb1 (8, 32) (Fig. 9B, lane 1, and C, lane 2). (iv) Treatment of whole-cell extracts with a cocktail of phosphatase inhibitors caused pre-rRNPs to be disrupted (P. Harnpicharnchai, unpublished), while the Nop7/Ytm1/Erb1 subcomplex remained intact. Nop7-TAP or Ytm1-TAP under these conditions resulted in the recovery of only the Ytm1/Nop7/Erb1 heterotrimer (Fig. 9D, lane 2) (5; data not shown). (v) Nop7-TAP from sucrose gradient fractions containing 66S preribosomes yielded most protein components of the 66S pre-rRNPs (Fig. 9E, lane 1). However, when gradient fractions containing 66S pre-rRNPs were treated with the phosphatase inhibitor cocktail prior to affinity purification with Nop7-TAP, mostly the heterotrimer was recovered (Fig. 9E, lane 2). The last result indicates that Ytm1,


A




B



C



D






FIG. 5. Inactivation of Ytm1 in the *ytm1-1* mutant causes 66S preribosomes to accumulate in the nucleolus. The *ytm1-1* mutant strain JWY6790 expressing eGFP-tagged rpL25 was grown in C-Trp medium at 25°C, washed and suspended in YEPD, and grown at 25°C (A and C) or shifted to 37°C for 5 h (B and D). Nuclei stained with 4',6'-diamidino-2-phenylindole (DAPI) are shown in panels A and B (typically, nucleoli do not stain with DAPI). The signal from RpL25eGFP is shown in panels C and D. Arrows indicate nucleolar accumulation of rpL25eGFP (D) and corresponding DAPI staining (B).

Erb1, and Nop7 form a stable complex within preribosomes sedimenting at 66S and can be released from the 66S pre-rRNPs by treatment with the phosphatase inhibitor cocktail. The molecular basis of the effect(s) of the phosphatase inhibitors is unknown.

Our current data suggest that the components of the Ytm1/Nop7/Erb1 heterotrimer join preribosomes separately since significant or small amounts of 35S pre-rRNA coprecipitate with Nop7 and Erb1, respectively, although no 35S pre-rRNA copurifies with Ytm1 (Fig. 7C). Thus, Nop7 likely joins nascent preribosomes first, followed by Erb1. Ytm1 later associates with 66S preribosomes containing 27SA₂ pre-rRNA (Fig. 7B).

Ytm1 and Nop7 directly interact with Erb1. To assay pairwise interactions between components of the heterotrimer and to determine whether the interactions are direct, we carried out GST pull-down assays. Ytm1 bound specifically to GST-Erb1, and Erb1 bound to GST-Ytm1, while Nop7 displayed strong binding to GST-Erb1 but not to GST-Ytm1 (Fig. 10A) (data not shown). Consistent with these observations, Pes1 and Bop1, the mammalian homologues of Nop7 and Erb1, bind to each other in vitro and interact in two-hybrid assays in vivo

(34). The interactions between Erb1 and Ytm1 were corroborated by two-hybrid assays in vivo. Cells expressing *AD-YTM1* and *BD-ERB1* or *AD-ERB1* and *BD-YTM1* displayed strong expression of the *GAL-HIS3* reporter gene (growth on 50 mM 3-aminotriazole) (data not shown). Thus, Ytm1 and Nop7 each bind directly to Erb1 but not to one another (Fig. 10C). Erb1, like Ytm1, contains WD40 repeats. Nop7 also contains a known protein-protein interaction motif, the BRCT domain (1). Further analysis is necessary to test whether these or other domains dictate the strong interactions among these three proteins.

Association of heterotrimer with 66S preribosomes is significantly weakened in the *ytm1-1* mutant. Since the two *ytm1-1* mutations are in residues that may be important for interactions with ligands, we determined the effects of these mutations on the integrity of the heterotrimer and 66S preribosomes. We used TAP-tagged Nop7 or Brx1 to purify 66S preribosomes from *YTM1* and *ytm1-1* strains, since both of these proteins are present in all seven different 66S pre-rRNPs (21, 60) (Fig. 1B). SDS-PAGE, silver staining, and Western blotting revealed that most of the proteins present in wild-type 66S pre-rRNPs are also present in equivalent amounts in the particles isolated

FIG. 4. Processing of pre-rRNAs is altered in the *ytm1-1* mutant. (A) Oligonucleotide probes or primers used to detect rRNAs and pre-rRNAs. (B) Yeast strains JWY3400 (*YTM1*) and JWY7128 (*ytm1-1*) were grown in YEPD medium at 25°C and shifted to 37°C for 5 h. Cells were pulse-labeled with [5,6-³H]uracil for 5 min and chased with an excess of unlabeled uracil for 2, 5, 10, and 60 min. Equal cpm of RNA isolated from cells at each time point were subjected to electrophoresis on agarose-formaldehyde gels to separate each pre-rRNA or rRNA and detected by autoradiography. (C) Primer extension was performed to determine steady-state levels of 27SA₂, 27SA₃, 27SB_L plus 7S_L, and 27SB_S plus 7S_S pre-rRNAs. RNA was extracted from whole-cell extracts from strains JWY3400 (*YTM1*) and JWY7128 (*ytm1-1*) grown in YEPD medium at 25°C or shifted from 25°C to 37°C for 3 h or 6 h or from strain JWY6149 (*YTM1*) or JWY6992 (*GAL-YTM1*) grown in galactose-containing medium and shifted to glucose-containing medium for 0, 10, 12, 15, or 18 h. (D) Northern blotting was used to determine steady-state levels of 25S, 18S, 5.8S, and 5S rRNA and 7S pre-rRNA. High-molecular-weight RNAs were subjected to electrophoresis on agarose-formaldehyde gels, whereas acrylamide-urea gels were used to separate low-molecular-weight RNAs. U3 snoRNA was used as a loading control. RNA was quantified by phosphorimaging and normalized to U3 snoRNA.




FIG. 6. Ytm1-HA3 cosediments on sucrose gradients with 66S preribosomes. Whole-cell extracts were prepared from yeast strain JWY6770 (*YTM1-HA3*) and fractionated on 7 to 47% sucrose velocity gradients. Fractions containing 40S and 60S ribosomal subunits and 80S monosomes are labeled. Proteins were trichloroacetic acid precipitated from gradient fractions and subjected to Western immunoblot analysis to detect Ytm1-HA3.

from the *ytm1-1* mutant, indicating that the 66S preribosomes are largely intact in the *ytm1-1* mutant (Fig. 11). However, Ytm1-1 mutant protein was greatly diminished or absent from the pool of purified 66S preribosomes (Fig. 11A, lanes 2 and 4, and B, lanes 2 and 4). Nop7 and Erb1 were still present, but in reduced amounts (Fig. 11A, lanes 2 and 4). Thus some of the mutant preribosomes contain Nop7 and/or Erb1 but not Ytm1, and others lack all three proteins. The relative amounts of each pre-rRNA copurifying with TAP-tagged Nop7 or Brx1 in 66S pre-rRNPs parallel those in whole-cell extracts (data not shown).

The Ytm1/Nop7/Erb1 heterotrimer is destabilized in the *ytm1-1* mutant. The effects of the *ytm1-1* mutations on 66S pre-rRNPs and pre-rRNA processing could result from alterations of the heterotrimer containing Ytm1. Therefore, we purified the heterotrimer from the *ytm1-1* mutant and examined its integrity, using three assays: sucrose gradient centrifugation, differential centrifugation, and treatment of whole-cell extracts with phosphatase inhibitors. In each case, only small amounts, if any, of Erb1 copurified with Nop7 and no Ytm1 could be detected (Fig. 9B, lane 2; C, lane 4, and D, lane 4). GST-pulldown assays confirmed that Ytm1-1 does not bind to Erb1 *in vitro* at 37°C (Fig. 10B). These results suggest that at the nonpermissive temperature, Ytm1-1 fails to interact with Erb1 and the Nop7-Erb1 association is significantly weakened, leading to destabilization of the heterotrimer and perturbations of preribosome maturation.

DISCUSSION

Although there has been much progress to identify yeast ribosome assembly intermediates and their protein and RNA constituents (reviewed in references 12 and 14), nothing is known about the architecture of these pre-rRNPs. For example, it is unclear which proteins are nearest neighbors within assembling ribosomes and to what extent neighboring molecules function together. Since Ytm1 contains multiple WD40 protein-protein interaction motifs, it is an excellent candidate for a molecule present in a multiprotein subcomplex comprising a neighborhood in assembling ribosomes.

Here we have shown that Ytm1 is a constituent of four consecutive 66S preribosomes and is necessary for steps in their maturation to 60S ribosomal subunits. Ytm1 associates with ribosome assembly factors Erb1 and Nop7 to form a stable subcomplex that is present within 66S preribosomes and that also exists separately from pre-rRNPs. Erb1 and Nop7 assemble into preribosomes prior to Ytm1. Mutations in WD40 motifs 6 and 7 of Ytm1 destabilize the heterotrimer and weaken association of each of the three proteins with 66S pre-rRNPs. Consequently, processing of pre-rRNAs and release of preribosomes from the nucleolus are delayed, resulting in production of fewer 60S




FIG. 7. Ytm1 associates with pre-rRNAs in 66S preribosomes. (A) Whole-cell extracts were prepared from the *YTM1-TAP* strain JWY7124 and from untagged strain JWY3400 grown at 30°C in YEPD medium to 6×10^7 cells/ml. RNA was extracted from whole cells and from tandem affinity-purified samples, subjected to electrophoresis on agarose-formaldehyde or acrylamide-urea gels, blotted to nitrocellulose, and assayed by Northern blotting with specific oligonucleotide probes complementary to pre-rRNAs and mature rRNAs. Five percent of total RNA and 100% of tandem affinity-purified RNA were assayed. (B) Primer extension analysis was used to assay 35S, 27SA₂, 27SA₃, and 25.5S pre-rRNAs, as well as the B_S and B_L 5' ends of 27S and 7S pre-rRNAs, using ³²P-labeled oligonucleotides. Products of primer extension were resolved on sequencing gels, dried, and exposed to X-ray film for detection by autoradiography. (C) 35S pre-rRNA copurifies with TAP-tagged Erb1 and Nop7 but not Ytm1. RNA in whole-cell extracts and copurifying RNAs were assayed by primer extension as described above.




FIG. 8. Nonribosomal proteins necessary for biogenesis of 60S ribosomal subunits, as well as ribosomal proteins, copurify with TAP-tagged Ytm1. Whole-cell extract was prepared from the *YTM1-TAP* strain JWY7124 grown at 30°C in YEPD medium to 6×10^7 cells/ml and subjected to tandem affinity purification. Proteins were trichloroacetic acid precipitated from column eluates and subjected to electrophoresis on 4 to 20% polyacrylamide gels. Proteins were stained with colloidal Coomassie blue, manually excised from the gel, digested with trypsin, and identified by matrix-assisted laser desorption ionization-time of flight mass spectrometry (Table 2).

ribosomal subunits. This is one of the first examples demonstrating the importance of protein-protein interactions and subcomplex integrity for assembly of eukaryotic ribosomes.

Ytm1 associates with 66S preribosomes. The copurification of Ytm1 with the same relative proportions of 27SA₂, 27SA₃, 27SB, 25.5S, and 7S pre-rRNAs as found in whole-cell extracts (Fig. 7) indicates that Ytm1 is present in each of the four consecutive 66S pre-rRNPs containing these pre-rRNAs, through out most or all of their lifetimes. No 20S pre-rRNA or 18S rRNA copurifies with Ytm1, consistent with Ytm1 participating in biogenesis of 60S ribosomal subunits but not 40S subunits. Although Ytm1-TAP particles contain moderate amounts of 5.8S rRNA, we failed to detect significant levels of 25S rRNA, consistent with a previous report indicating that conversion of 7S pre-rRNA to 5.8S rRNA occurs more rapidly than production of 25S rRNA from 25.5S pre-rRNA (18).

We also identified 53 proteins that copurify with TAP-tagged Ytm1 (Fig. 8, Table 2). Of the proteins isolated, 17 are ribosome assembly factors previously shown to be components of 66S preribosomes (Table 2). None of the proteins found specifically in late nucleoplasmic or cytoplasmic 66S pre-rRNPs (41) are present in preribosomes purified using Ytm1-TAP. This is consistent with our RNA and subcellular localization data indicating that Ytm1 dissociates from pre-rRNPs in the nucleoplasm prior to the latest nucleoplasmic and cytoplasmic stages of ribosome maturation. Thus 66S preribosomes purified using Ytm1-TAP may represent core complexes of molecules important for intermediate steps in 60S subunit ribosome biogenesis.

Ytm1 is present in a heterotrimeric complex with Nop7 and Erb1. Our results and those of others indicate that Ytm1, Erb1, and Nop7 form a stable heterotrimeric complex (8, 21, 32) (Fig. 8, 9, and 10). The sedimentation of Nop7, Ytm1, and Erb1 on sucrose gradients indicates that in wild-type cells most of this microparticle exists within 66S preribosomes (21; our unpublished results) (Fig. 6 and 9E). Both Ytm1 and Nop7

TABLE 2. Nonribosomal proteins that copurify with Ytm1-TAP

Gel ID no. ^a	Protein	ORF	Characteristic				Pre-rRNA processing ^c	Localization ^b	Reference
			Metazoan homolog	Essential	Sediments at 66S	Decreased 60S Subunits			
33	Ytm1	YOR272w	+	+	+	+	27SA ₃ ↑ 25.5S ↓ 7S ↓	No/Nu	This work (21)
10	Nip7	YPL211w	+	+	+	+	ND ^d	No/Nu/C	61
24	Nop16	YER002c	-	-	+	+	ND	No/Nu	21
26	Brx1	YOL077c	+	+	ND	+	27SA ↑ 27SB ↓	No	31
28	Rlp7	YNL002c	+	+	ND	+	27SA ₃ ↑ 27SB ↓ 7S ↓	No	9, 15
28	Rpf2	YKR081c	+	+	+	+	27SB ↑	No	39, 60
29	Cic1/Nsa3	YHR052c	-	+	+	ND	27S ↓ 7S ↓	No	13, 21, 41
31	Nsa1	YGL111w	-	+	ND	+	ND	ND	21
32, 33	Has1	YMR290c	+	+	ND	+/-	35S ↑ 27SB ↑	No	10
34	Ebp2	YKL172w	+	+	ND	+	35S ↑ 27SA ₃ ↑	No	27, 57
35, 36	Nog1	YPL093w	+	+	+	+	27S ↓ 7S ↓	No	30
37	Nop7	YGR103w	+	+	+	+	27S → 25S slow	No	1
37	Noc3	YLR002c	+	+	+	+	35S ↑ 27SB ↑	No/Nu	37
38	Noc2	YOR206w	+	+	+	+	35S ↑ 27SB ↑	No/Nu	37
38	Nop2	YGL111w	+	+	+	+	35S ↑ 27S ↑	No	6, 25
39	Drs1	YLL008w	-	+	ND	+	27SB ↑	No	48
40	Erb1	YMR049c	+	+	+	+	27S ↑	No	44

^a Protein band 30 was identified as Tef2 and protein band 36 also contained Ssa1. These proteins are common contaminants of affinity-purified complexes (17, 24).

^b No, nucleolar; Nu, nuclear; C, cytoplasmic.

^c ↑ and ↓, increased and decreased amounts of indicated RNAs, respectively.

^d ND, not determined.




FIG. 9. Ytm1, Erb1, and Nop7 form a heterotrimeric subcomplex both within 66S preribosomes and independently of these particles. (A) Ytm1, Erb1, and Nop7 are enriched (relative to other proteins found in 66S pre-rRNPs) among proteins copurifying with Ytm1-TAP or Nop7-TAP from *rrp1-1* or *nop4-3* mutants in which 66S preribosomes are unstable. Heterotrimer was purified (B) from sucrose gradient fractions, (C) by differential centrifugation, or (D) from whole-cell extracts treated with a cocktail of phosphatase inhibitors that disrupt 66S pre-rRNPs. Wild-type cells or mutant cells were grown at 25°C and shifted to 37°C for 5 h. Tandem affinity purification using Nop7-TAP or Ytm1-TAP was carried out from (A) whole-cell extracts from a 50-ml culture, (B) gradient fractions 5 to 7 (prepared from a 900-ml culture), (C) whole-cell extracts from 50 ml of cells (lanes 1 and 3) or 180,000 × g spin supernatants prepared from 500 ml of cells (lanes 2 and 4), or (D) untreated (–) or phosphatase inhibitor cocktail-treated extracts (+). Purified proteins were resolved by SDS-PAGE. Note that the heterotrimer is destabilized in the *ytm1-1* mutant (B, lane 2; C, lane 4; D, lane 4). (E) Ytm1, Erb1, and Nop7 form a stable subcomplex within 66S preribosomes. Whole-cell extracts from *YTM1* cells were subjected to centrifugation on 7 to 47% gradients. Fractions containing 66S preribosomes were pooled and subjected to tandem affinity purification in the presence (+) or absence (–) of phosphatase inhibitors, and proteins were resolved by SDS-PAGE. Bands indicated by asterisks in B and E are common contaminants that we observe upon TAP from any fractions of sucrose gradients (top, middle, or bottom) using any TAP-tagged protein.

bind tightly to Erb1; however, we find no evidence for direct interactions between Ytm1 and Nop7 (Fig. 10). The ability to purify the heterotrimer directly from 66S preribosomes using the phosphatase inhibitor cocktail that disrupts 66S pre-rRNPs (Fig. 9E) suggests that interactions among these three proteins are stronger than those with other molecules in the 66S pre-rRNP. Nevertheless, it is reasonable to assume that this subcomplex influences the assembly or function of a larger protein or RNP neighborhood within assembling ribosomes necessary for their efficient maturation (see below). Thus, it will be important to identify other molecules that are adjacent to Ytm1, Erb1, and Nop7 within preribosomes.

The heterotrimeric complex also exists independently of 66S preribosomes. It can be purified from fractions near the top of

gradients or from high-speed supernatants (Fig. 9B and C). Our present data suggest that Nop7 and Erb1 may assemble into preribosomes before Ytm1. Consistent with this finding is the presence of Nop7 and Erb1, but not Ytm1, in 66S pre-rRNPs isolated from the *ytm1-1* mutant. Nop7 and Erb1 may not require Ytm1 to assemble into or remain in preribosomes. Upon completion of their functions, Nop7, Erb1, and Ytm1 might dissociate from preribosomes together as a heterotrimer, prior to recycling into other nascent preribosomes. The substantial amount of heterotrimer that we purify from cells might also result from disassembly in vivo of unstable abortive assembly intermediates or upon dissociation from preribosomes in vitro during fractionation or purification.




FIG. 10. Ytm1 and Nop7 directly interact with Erb1. (A) Synthetic radiolabeled proteins (*) were incubated with GST fusion proteins (lanes 2, 5, and 8). As negative controls, synthetic peptides were incubated with GST beads only (lanes 1, 4, and 7) or GST fusion proteins were incubated with the unrelated, radiolabeled 40S ribosome assembly factor Krr1 or ribosomal protein L11 (lanes 3, 6, and 9). (B) Radiolabeled wild-type Ytm1 protein was preincubated at 37°C for 15 min (lane 2). Mutant Ytm1-1 protein was preincubated at 37°C for 15 min (lane 3), 30 min (lane 4), or 60 min (lane 5). Following preincubation, wild-type Ytm1 or mutant Ytm1-1 radiolabeled protein was incubated with GST-Erb1. Mutant Ytm1-1 protein was incubated with GST beads only (lane 1) as a negative control. Complexes were eluted from glutathione beads, subjected to electrophoresis on 10% polyacrylamide gels, and detected by autoradiography. (C) Model for interactions between Ytm1, Erb1, and Nop7. Gray lines indicate interactions detected using GST pulldown assays, whereas the black line indicates interactions detected by two-hybrid assays.

A conserved network of protein interactions. Interactions among Ytm1, Erb1, and Nop7 required for ribosome biogenesis are likely to be conserved. Interactions of Pes1, the mouse homologue of Nop7, with Bop1, the mouse homologue of Erb1, are required for assembly of Pes1 into preribosomes (34). Like their yeast counterparts, Pes1 and Bop1 are required for similar steps in processing of pre-rRNAs to mature 25S and 5.8S rRNA (1, 35, 44, 55, 56).

Mutations in the WD40 motifs of Ytm1 destabilize the heterotrimer and weaken its association with preribosomes. The G398D and S442N mutations in WD40 repeats 6 and 7 of *YTM1* prevent binding of Ytm1-1 to Erb1 in vitro and significantly weaken association between Nop7 and Erb1 (Fig. 9 and 10). Destabilization of the heterotrimer also weakens association of Ytm1, Erb1, and Nop7 with 66S pre-rRNPs, resulting in recovery of greatly diminished amounts of Ytm1 and slightly decreased amounts of Erb1, relative to Nop7, in the pool of purified 66S preribosomes (Fig. 11). The mixture of 66S pre-rRNPs purified from the *ytm1-1* mutant using Brx1-TAP contains no detectable Ytm1-1 and less Nop7 and Erb1 compared to preribosomes isolated from wild-type cells (Fig. 11). (Note that no free heterotrimer could copurify with Brx1-TAP.) The greater decrease of Ytm1 compared to Erb1 in Nop7-TAP-purified *ytm1-1* mutant particles suggests that the Ytm1-Erb1 interaction may be necessary to recruit or maintain Ytm1 in 66S pre-rRNPs. This interaction may also influence assembly of Nop7 and Erb1 into preribosomes or may be required to stabilize their association with preribosomes, since amounts of

Nop7 are reduced slightly in the 66S preribosomes purified from *ytm1-1* mutant cells expressing Brx1-TAP.

Ytm1 is required for pre-rRNA processing and trafficking of 66S preribosomes. Our data show that pre-rRNA processing is slowed in the *ytm1* mutants. The conversion of 27SA₃ pre-rRNA to 27SB pre-rRNA is delayed, resulting in changes in amounts of 27SB_S and 27SB_L pre-rRNAs. Subsequent steps in pre-rRNA processing are also slowed, resulting in reduced amounts of 7S pre-rRNA and 5S rRNA (Fig. 4). Processing of 27SA₃ to 27SB pre-rRNA involves rapid exonucleolytic trimming of the 5' end of the 27SA₃ pre-rRNA by Rat1 and Xrn1 (23). Ytm1 might function directly in all of these steps. Alternatively, the decreased rate of processing of 27SA₃ pre-rRNA in the *ytm1* mutants might indirectly affect subsequent processing of 27SB pre-rRNA, for example, by perturbing the architecture of pre-rRNPs. Depletion of Nop7 and Erb1 has effects on pre-rRNA processing similar but not completely identical to those of *ytm1* mutants: increased levels of 27SA₃ pre-rRNA and decreased amounts of 27SB and 7S pre-rRNAs (42, 44). Thus, the pre-rRNA processing phenotypes of the *ytm1* mutants may reflect a combination of effects on the presence and/or functions of all three proteins in the Ytm1/Erb1/Nop7 heterotrimer.

Most, but not all steps of processing of precursors to 25S and 5.8S rRNA are thought to occur prior to nucleolar release and most likely require or are accompanied by many changes in the topology of these pre-rRNPs. Although the mechanism whereby preribosomes exit the nucleolus remains a mystery, it may require




FIG. 11. 66S preribosomes are largely intact but lack Ytm1 in the *ytm1-1* mutant. Wild-type *YTM1* cells and mutant *ytm1-1* cells expressing Nop7-TAP or Brx1-TAP were grown in YEPD medium at 25°C and shifted to 37°C for 5 h. (A) Silver staining or (B) Western immunoblot analysis was performed on proteins trichloroacetic acid precipitated from samples affinity purified from extracts from these strains.

some changes in the composition and structure of preribosomes, including those resulting from pre-rRNA processing. Nuclear export of preribosomes requires binding of export factors to preribosomes, perhaps timed by the availability of ligands on the surface of preribosomes (22). Thus pre-rRNA processing, nucleolar release, and nuclear export may be intertwined to prevent premature exit of incompletely assembled ribosomes from the nucleolus and nucleus. Therefore, rather than binding directly to export factors, Ytm1, as well as Erb1 and Nop7, is likely to participate indirectly in nucleolar release and nuclear export of preribosomes, by functioning together to establish or maintain structures necessary for these processes. Indeed, as observed for the *ytm1-1* mutant, preribosomes also accumulate in the nucleus upon depletion of Nop7 (42). The *ytm1-1* mutations, however, cause 66S preribosomes to be retained in the nucleolus (Fig. 5).

Functions of the heterotrimer in ribosome biogenesis. Taken together, our results suggest the following model for effects of

ytm1-1 mutations on ribosome assembly. In wild-type cells, early in ribosome assembly during or after synthesis of 35S pre-rRNA, Nop7 and Erb1 join nascent preribosomes. Slightly later, many 60S ribosome assembly proteins, including Ytm1, associate with the 27SA₂ pre-rRNA to form the earliest detectable 66S pre-rRNP. In the *ytm1-1* mutant, Ytm1-1 may not efficiently assemble into this pre-rRNP or may not remain stably associated with preribosomes, also resulting in destabilization of association of Nop7 and Erb1 with pre-60S ribosomes. Although it is also possible that Ytm1, Nop7, or Erb1 dissociates from the mutant particles during purification rather than in vivo, these mutant particles are likely otherwise intact. However, their overall architecture might be perturbed by the absence or weakened association of Ytm1, Erb1, and Nop7. When preribosomes lack these putative scaffolding proteins, they might be unable to establish or maintain structures necessary for efficient pre-rRNA processing, ribosome assembly, release of nascent ribosomes from the nucleolus, or export of preribosomes from the nucleus to the cytoplasm.

Moonlighting functions of the heterotrimeric proteins. Clearly Ytm1 is necessary for efficient ribosome production. Depletion or inactivation of Ytm1 decreases the rate of 60S subunit biogenesis below levels necessary to sustain viability. However, Ytm1 may have a second “moonlighting” function, such as mitosis or chromosome transmission (36, 43; Matsumoto et al., personal communication). Interestingly, Nop7 and its metazoan homologue Pescadillo, and Bop1, the mammalian homologue of Erb1, also have independent functions. Nop7 and Pescadillo are implicated in DNA replication or cell proliferation (8, 35). Mutations in *BOP1* induce p53-dependent cell cycle arrest (45). Thus, the heterotrimeric complex of Ytm1, Erb1, and Nop7 might be a depot for directing multiple functions of these proteins.

ACKNOWLEDGMENTS

We thank Seiji Matsumoto for generously providing *ytm1-1* and *YTM1* strains and for communicating unpublished results. We thank Stan Fields and Tony Hazbun (Yeast Resource Center, University of Washington) for plasmids containing *YTM1*, *ERB1*, or *NOP7* cloned into pOAD. We also thank Michael McAlear, Jonathan Warner, David Goldfarb, Janine Maddock, Jennifer Fuentes, and Bernard Trumpower for antibodies against ribosomal proteins or ribosome assembly factors. We are grateful to Susan Dowd and Mark Bier (Center for Molecular Analysis, Carnegie Mellon University) for assistance with mass spectrometry. We thank Jon Minden, Peter Berget, Jeff Brodsky, Elizabeth Jones, and members of our laboratory for fruitful discussions and for comments on the manuscript. We thank Brooke McCartney for use of her microscope.

This work was supported by National Institutes of Health grants RO1 GM28301 to J.L.W., F31 GM65067 to T.D.R., and F31 GM19937 to E.W.H. P.H. was supported by the government of Thailand.

REFERENCES

- Adams, C. C., J. Jakovljevic, J. Roman, P. Harnpicharnchai, and J. L. Woolford, Jr. 2002. *Saccharomyces cerevisiae* nucleolar protein Nop7 is necessary for biogenesis of 60S ribosomal subunits. *RNA* 8:150–165.
- Andersen, J. S., C. E. Lyon, A. H. Fox, A. K. L. Leung, Y. W. Lam, H. Steen, M. Mann, and A. I. Lamond. 2002. Directed proteomic analysis of the human nucleolus. *Curr. Biol.* 12:1–11.
- Baßler, J., P. Grandi, O. Gadal, T. Lessmann, E. Petfalski, D. Tollervey, J. Lechner, and E. Hurt. 2001. Identification of a 60S pre-ribosomal particle that is closely linked to nuclear export. *Mol. Cell* 8:517–529.
- Basu, U., K. Si, J. R. Warner, and U. Maitra. 2001. The *Saccharomyces cerevisiae* *TIF6* gene encoding translation initiation factor 6 is required for 60S ribosomal subunit biogenesis. *Mol. Cell. Biol.* 21:1453–1462.
- Cagney, G., P. Uetz, and S. Fields. 2001. Two-hybrid analysis of the *Saccharomyces cerevisiae* 26S proteasome. *Physiol. Genomics* 7:27–34.

6. de Beus, E., J. S. Brockenbrough, B. Hong, and J. P. Aris. 1994. Yeast *NOP2* encodes an essential nucleolar protein with homology to a human proliferation marker. *J. Cell Biol.* **127**:1799–1813.
7. Dragon, F., J. E. Gallagher, P. A. Compagnone-Post, B. M. Mitchell, K. A. Porwancher, K. A. Wehner, S. Wormsley, R. E. Settlage, J. Shabanowitz, Y. Osheim, A. L. Beyer, D. F. Hunt, and S. J. Baserga. 2002. A large U3 snoRNP complex which is required for 18S rRNA biogenesis. *Nature* **417**: 967–970.
8. Du, Y. C., and B. Stillman. 2002. Yph1p, an ORC-interacting protein: potential links between cell proliferation control, DNA replication, and ribosome biogenesis. *Cell* **109**:835–848.
9. Dunbar, D. A., F. Dragon, S. J. Lee, and S. J. Baserga. 2000. A nucleolar protein related to ribosomal protein L7 is required for an early step in large ribosomal subunit biogenesis. *Proc. Natl. Acad. Sci. USA* **97**:13027–13032.
10. Emery, B., J. de la Cruz, S. Rocak, O. Deloche, and P. Linder. 2004. Has1p, a member of the DEAD-box family, is required for 40S ribosomal subunit biogenesis in *Saccharomyces cerevisiae*. *Mol. Microbiol.* **52**:141–158.
11. Fatica, A., A. D. Cronshaw, M. Dlakis, and D. Tollervey. 2002. Ssl1p prevents premature processing of an early pre-60S ribosomal particle. *Mol. Cell* **9**:341–351.
12. Fatica, A., and D. Tollervey. 2002. Making ribosomes. *Curr. Opin. Cell Biol.* **14**:313–318.
13. Fatica, A., M. Oeffinger, D. Tollervey, and I. Bozzoni. 2003. Cic1p/Nsa3p is required for synthesis and nuclear export of 60S ribosomal subunits. *RNA* **9**:1431–1436.
14. Fromont-Racine, M., B. Senger, C. Saveanu, and F. Fasiolo. 2003. Ribosome assembly in eukaryotes. *Gene* **313**:17–42.
15. Gadal, O., D. Strauss, E. Petfalski, P. E. Gleizes, N. Gas, D. Tollervey, and E. Hurt. 2002. Rlp7p is associated with 60S preribosomes, restricted to the granular component of the nucleolus, and required for pre-rRNA processing. *J. Cell Biol.* **157**:941–951.
16. Gaudet, R., A. Bohm, and P. B. Sigler. 1996. Crystal structure at 2.4 Å resolution of the complex of transducin $\beta\gamma$ and its regulator, phosducin. *Cell* **87**:577–588.
17. Gavin, A. C., M. Bosche, R. Krause, P. Grandi, M. Marzoch, A. Bauer, J. Schultz, J. M. Rick, A. M. Michon, C. M. Cruciat, M. Remor, C. Hofert, M. Schelder, M. Brajenovic, H. Ruffner, A. Merino, K. Klein, M. Hudak, D. Dickson, T. Rudi, V. Gnau, A. Bauch, S. Bastuck, B. Huhse, C. Leutwein, M. A. Heurtier, R. R. Copley, A. Edelmann, E. Querfurth, V. Rybin, G. Drewes, M. Raida, T. Bouwmeester, P. Bork, B. Seraphin, B. Kuster, G. Neubauer, and G. Superti-Furga. 2002. Functional organization of the yeast proteome by systematic analysis of protein complexes. *Nature* **415**:141–147.
18. Geerlings, T. H., J. C. Vos, and H. A. Raué. 2000. The final step in the formation of 25S rRNA in *Saccharomyces cerevisiae* is performed by 5'→3' exonucleases. *RNA* **6**:1698–1703.
19. Gelperin, D., L. Horton, J. Beckman, J. Hensold, and S. K. Lemmon. 2001. Bms1p, a novel GTP-binding protein, and the related Tsr1p are required for distinct steps of 40S ribosome biogenesis in yeast. *RNA* **9**:1268–1283.
20. Grandi, P., V. Rybin, J. Bassler, E. Petfalski, D. Strauss, M. Marzoch, T. Schafer, B. Kuster, H. Tschochner, D. Tollervey, A. C. Gavin, and E. Hurt. 2002. 90S pre-ribosomes include the 35S pre-rRNA, the U3 snoRNP, and 40S subunit processing factors but predominantly lack 60S synthesis factors. *Mol. Cell* **10**:105–115.
21. Harnpicharnchai, P., J. Jakovljevic, E. Horsey, T. Miles, J. Roman, M. Rout, D. Meagher, B. Imai, Y. Guo, C. J. Brame, J. Shabanowitz, D. F. Hunt, and J. L. Woolford, Jr. 2001. Composition and functional characterization of yeast 66S ribosome assembly intermediates. *Mol. Cell* **8**:505–515.
22. Hedges, J., M. West, and A. R. Johnson. 2005. Release of the export adapter, Nmd3p, from the 60S ribosomal subunit requires Rpl10p and the cytoplasmic GTPase Lsg1p. *EMBO J.* **24**:567–579.
23. Henry, Y., H. Wood, J. P. Morrissey, E. Petfalski, S. Kearsy, and D. Tollervey. 1994. The 5' end of yeast 5.8S rRNA is generated by exonucleases from an upstream cleavage site. *EMBO J.* **10**:2452–2463.
24. Ho, Y., A. Gruhler, A. Heilbut, G. D. Bader, L. Moore, S. L. Adams, A. Millar, P. Taylor, K. Bennett, K. Boutlier, L. Yang, C. Wolting, I. Donaldson, S. Schandorff, J. Shewnarane, M. Vo, J. Taggart, M. Goudreau, B. Muskaf, C. Alfano, D. Dewar, Z. Lin, K. Michalikova, A. R. Willems, H. Sassi, P. A. Nielsen, K. J. Rasmussen, J. R. Andersen, L. E. Johansen, L. H. Hansen, H. Jespersen, A. Podtelejnikov, E. Nielsen, S. Crawford, V. Poulsen, B. D. Sorensen, J. Matthiesen, R. C. Hendrickson, F. Gleeson, T. Pawson, M. F. Moran, D. Durocher, M. Mann, C. W. Hogue, D. Figey, and M. Tyers. 2002. Systematic identification of protein complexes in *Saccharomyces cerevisiae* by mass spectrometry. *Nature* **415**:180–183.
25. Hong, B., J. S. Brockenbrough, P. Wu, and J. P. Aris. 1997. Nop2p is required for pre-rRNA processing and 60S ribosome subunit synthesis in yeast. *Mol. Cell Biol.* **17**:378–388.
26. Horsey, E. W., J. Jakovljevic, T. D. Miles, P. Harnpicharnchai, and J. L. Woolford, Jr. 2004. Role of the yeast Rpl1 protein in the dynamics of pre-ribosome maturation. *RNA* **10**:813–827.
27. Huber, M. D., J. H. Dworet, K. Shire, L. Frappier, and M. A. McAlear. 2000. The budding yeast homolog of the human EBNA1-binding protein 2 (Ebp2p) is an essential nucleolar protein required for pre-rRNA processing. *J. Biol. Chem.* **275**:28764–28773.
28. Hurt, E., S. Hannus, B. Schmeizl, D. Lau, D. Tollervey, and G. Simos. 1999. A novel *in vivo* assay reveals inhibition of ribosomal nuclear export in Ran-cycle and nucleoporin mutants. *J. Cell Biol.* **144**:389–401.
29. James, P., J. Halladay, and E. Craig. 1996. Genomic libraries and a host strain designed for highly efficient two-hybrid selection in yeast. *Genetics* **144**:1425–1436.
30. Kallstrom, G., J. Hedges, and A. Johnson. 2003. The putative GTPases Nog1p and Lsg1p are required for 60S ribosomal subunit biogenesis and are localized to the nucleus and cytoplasm, respectively. *Mol. Cell Biol.* **23**: 4344–4355.
31. Kaser, A., E. Bogengruber, M. Hallegger, E. Doppler, G. Lepperding, M. Jantsch, M. Breitenbach, and G. Kreil. 2001. Brix from *Xenopus laevis* and Brx1p from yeast define a new family of proteins involved in the biogenesis of large ribosomal subunits. *Biol. Chem.* **382**:1637–1647.
32. Krogan, N. J., W. T. Peng, G. Cagney, M. D. Robinson, R. Haw, G. Zhong, X. Guo, X. Zhang, V. Canadien, D. P. Richards, B. K. Beattie, A. Lalev, W. Zhang, A. P. Davierwala, S. Mnaimneh, A. Starostine, A. P. Tikuisis, J. Grigull, N. Datta, J. E. Bray, T. R. Hughes, A. Emili, and J. F. Greenblatt. 2004. High-definition macromolecular composition of yeast RNA-processing complexes. *Mol. Cell* **13**:225–239.
33. Lambright, D. G., J. Sondke, A. Bohm, N. P. Skiba, H. E. Hamm, and P. B. Sigler. 1996. The 2Å crystal structure of a heterotrimeric G protein. *Nature* **379**:311–319.
34. Lapik, Y. R., C. J. Fernandes, L. F. Lau, and D. G. Pestov. 2004. Physical and functional interaction between Pes1 and Bop1 in mammalian ribosome biogenesis. *Mol. Cell* **15**:17–29.
35. Lerch-Gaggli, A., J. Haque, J. Li, G. Ning, P. Traktman, and S. A. Duncan. 2002. Pescadillo is essential for nucleolar assembly, ribosome biogenesis, and mammalian cell proliferation. *J. Biol. Chem.* **277**:45347–45355.
36. Matsuzaki, F., S. Matsumoto, and I. Yahara. 1988. Truncation of the carboxy-terminal domain of yeast β -tubulin causes temperature-sensitive growth and hypersensitivity to antimetabolic drugs. *J. Cell Biol.* **107**:1427–1435.
37. Milkereit, P., O. Gadal, A. Podtelejnikov, S. Trumtel, N. Gas, E. Petfalski, D. Tollervey, M. Mann, E. Hurt, and H. Tschochner. 2001. Maturation and intranuclear transport of pre-ribosomes requires Noc proteins. *Cell* **105**:499–509.
38. Milkereit, P., D. Strauss, J. Bassler, O. Gadal, H. Kuhn, S. Schutz, N. Gas, J. Lechner, E. Hurt, and H. Tschochner. 2003. A Noc complex specifically involved in the formation and nuclear export of ribosomal 40 S subunits. *J. Biol. Chem.* **278**:4072–4081.
39. Morita, D., K. Miyoshi, Y. Matsui, A. Toh-E, H. Shinkawa, T. Miyakawa, and K. Mizuta. 2002. Rpf2p, an evolutionarily conserved protein, interacts with ribosomal protein L11 and is essential for the processing of 27SB pre-rRNA to 25S rRNA and the 60S ribosomal subunit assembly in *Saccharomyces cerevisiae*. *J. Biol. Chem.* **277**:28780–28786.
40. Neer, E. J., and T. F. Smith. 1996. G protein heterodimers: new structures propel new questions. *Cell* **84**:175–178.
41. Nissan, T. A., J. Baßler, E. Petfalski, D. Tollervey, and E. Hurt. 2002. Pre-60S particles on the road to ribosomes. *EMBO J.* **21**:5539–5547.
42. Oeffinger, M., A. Leung, A. Lamond, and D. Tollervey. 2002. Yeast Pescadillo is required for multiple activities during 60S ribosomal subunit synthesis. *RNA* **8**:626–636.
43. Ouspenski, I., S. J. Elledge, and B. R. Brinkley. 1999. New yeast genes important for chromosome integrity and segregation identified by dosage effects on genome stability. *Nucleic Acids Res.* **27**:3001–3008.
44. Pestov, D. G., M. G. Stockelman, Z. Strezoska, and L. F. Lau. 2001. ERB1, the yeast homolog of mammalian Bop1, is an essential gene required for maturation of the 25S and 5.8S ribosomal RNAs. *Nucleic Acids Res.* **29**: 3621–3630.
45. Pestov, D. G., Z. Strezoska, and L. F. Lau. 2001. Evidence of p53-dependent cross-talk between ribosome biogenesis and the cell cycle: effects of nucleolar protein Bop1 on G₁/S transition. *Mol. Cell Biol.* **21**:4246–4255.
46. Raué, H. A. 2003. Pre-ribosomal RNA processing and assembly in *Saccharomyces cerevisiae*. The machine that makes the machine, p. 1–24. *In* M. O. J. Olson (ed.), *The nucleolus*. Kluwer Academic/Plenum Publishers, Dordrecht, The Netherlands.
47. Rigaut, G., A. Shevchenko, B. Rutz, M. Wilm, M. Mann, and B. Seraphin. 1999. A generic protein purification method for protein complex characterization and proteome exploration. *Nat. Biotechnol.* **17**:1030–1032.
48. Ripmaster, T. L., G. P. Vaughn, and J. L. Woolford, Jr. 1992. A putative ATP-dependent RNA helicase involved in *Saccharomyces cerevisiae* ribosome assembly. *Proc. Natl. Acad. Sci. USA* **89**:11131–11135.
49. Saveanu, C., D. Bienvenu, A. Namane, P. E. Gleizes, N. Gas, A. Jacquier, and M. Fromont-Racine. 2001. Nog2p, a putative GTPase associated with pre-60S subunits and required for late 60S maturation steps. *EMBO J.* **20**:6475–6484.
50. Saveanu, C., A. Namane, P.-E. Gleizes, A. Lebreton, J.-C. Rousselle, J. Noaillac-Depeyre, N. Gas, A. Jacquier, and M. Fromont-Racine. 2003. Sequential protein association with nascent 60S ribosomal particles. *Mol. Cell Biol.* **23**:4449–4460.

51. Schäfer, T., D. Strauß, E. Petfalski, D. Tollervey, and E. Hurt. 2003. The path from nucleolar 90S to cytoplasmic 40S pre-ribosomes. *EMBO J.* **22**: 1370–1380.
52. Scherl, A., Y. Coute, C. Deon, A. Calle, K. Kindbeiter, J.-C. Sanchez, A. Greco, D. Hochstrasser, and J. J. Diza. 2002. Functional proteomic analysis of the human nucleolus. *Mol. Biol. Cell* **13**:4100–4109.
53. Smith, T. F., C. Gaitatzes, K. Sacena, and E. J. Neer. 1999. The WD repeat: a common architecture for diverse functions. *Trends Biochem. Sci.* **24**:181–185.
54. Sondek, J., A. Böhm, D. G. Lambright, H. E. Hamm, and P. B. Sigler. 1996. Crystal structure of a G protein $\beta\gamma$ dimer at 2.1Å resolution. *Nature* **379**: 369–374.
55. Strezoska, Ž., D. G. Pestov, and L. F. Lau. 2000. Bop1 is a mouse WD40 repeat nucleolar protein involved in 28S and 5.8S rRNA processing and 60S ribosome biogenesis. *Mol. Cell. Biol.* **20**:5516–5528.
56. Strezoska, Z., D. G. Pestov, and L. F. Lau. 2002. Functional inactivation of the mouse nucleolar protein Bop1 inhibits multiple steps in pre-rRNA processing and blocks cell cycle progression. *J. Biol. Chem.* **277**:29617–29625.
57. Tsujii, R., K. Miyoshi, A. Tsuno, Y. Matsui, A. Toh-e, T. Miyakawa, and K. Mizuta. 2000. Ebp2p, yeast homologue of a human protein that interacts with Epstein-Barr virus nuclear antigen 1, is required for pre-rRNA processing and ribosomal subunit assembly. *Genes Cells* **5**:543–553.
58. Venema, J., R. J. Planta, and H. A. Raue. 1998. In vivo mutational analysis of ribosomal RNA in *Saccharomyces cerevisiae*. *Methods Mol. Biol.* **77**:257–270.
59. Wall, M. A., D. E. Coleman, E. Lee, J. A. Iniguez-Lluhi, B. A. Posner, A. G. Gilman, and S. R. Sprang. 1995. The structure of the G protein heterotrimer Gi alpha 1 beta 1 gamma 2. *Cell* **83**:1047–1058.
60. Wehner, K. A., and S. J. Baserga. 2002. The $\sigma 70$ -like motif: a eukaryotic RNA binding domain unique to a superfamily of proteins required for ribosome biogenesis. *Mol. Cell* **9**:329–339.
61. Zanchin, N. I., P. Roberts, A. DeSilva, F. Sherman, and D. S. Goldfarb. 1997. *Saccharomyces cerevisiae* Nip7p is required for efficient 60S ribosome subunit biogenesis. *Mol. Cell. Biol.* **17**:5001–5015.