

Yeast Repressor $\alpha 2$ Binds to Its Operator Cooperatively with Yeast Protein Mcm1

CYNTHIA A. KELEHER,¹ STEVEN PASSMORE,² AND ALEXANDER D. JOHNSON^{3*}

Department of Biochemistry and Biophysics¹ and Department of Microbiology,³ University of California, San Francisco, California 94143, and Section of Biochemistry, Molecular and Cell Biology, Cornell University, Ithaca, New York 14853²

Received 13 June 1989/Accepted 14 August 1989

To bring about repression of a family of genes in *Saccharomyces cerevisiae* called the a-specific genes, two transcriptional regulatory proteins, $\alpha 2$ and GRM (general regulator of mating type), bind cooperatively to an operator found upstream of each a-specific gene. To date, GRM has been defined only biochemically. In this communication we show that the product of a single yeast gene (*MCMI*) is sufficient to bind cooperatively with $\alpha 2$ to the operator. We also show that antiserum raised against the *MCMI* gene product recognizes GRM from yeast cells. These results, in combination with previous observations, provide strong evidence that *MCMI* encodes the GRM activity.

The expression of many eucaryotic genes appears to be controlled by combinations of regulatory proteins (for a recent review, see reference 2). An example of this phenomenon is found in the repression of a group of cell-type-specific genes (called the a-specific genes) in the α cell type of the yeast *Saccharomyces cerevisiae*. This repression requires the activity of two proteins that bind cooperatively to an operator located upstream of each a-specific gene. Either protein alone can recognize the operator, but protein-protein interactions between the two couple their affinities for the DNA and ensure tight binding by the complex. The proteins are $\alpha 2$, a cell-type-specific protein, and GRM, a non-cell-type-specific protein. GRM has been defined only in terms of biochemical function, as follows: GRM is an activity that is (i) capable of binding alone to the a-specific gene operator and (ii) able to bind the operator cooperatively with $\alpha 2$ (4).

Considerable evidence suggests that GRM is encoded by the *MCMI* gene. The *MCMI* gene was originally identified by the isolation of a mutation, *mcm1-1*, that rendered strains unable to maintain certain plasmids (hence its name, from minichromosome maintenance) (6; G. R. Maine, Ph.D. thesis, Cornell University, 1984). However, α cells that carried the *mcm1-1* allele exhibited an additional phenotype; they mated poorly (9), a property expected for a cell deficient in GRM. This connection was strengthened by the observation that GRM-binding activity in extracts derived from *mcm1-1* strains was lower than that in extracts from wild-type strains (4). Recently, the *MCMI* gene product (produced in *Escherichia coli*) has been shown to meet the first criterion for GRM activity, namely, it recognizes the a-specific gene operator (8).

To determine whether Mcm1 meets the second criterion of GRM activity, we tested the ability of the *MCMI* gene product produced in *E. coli* to bind to an a-specific gene operator cooperatively with $\alpha 2$. A portion of the *MCMI* coding sequence (amino acids 1 through 188 [total, 286]) fused to 13 amino acids of a bacteriophage protein was overexpressed in *E. coli* and partially purified (30% of the preparation was Mcm1 [see reference 8 for details]). The gel mobility shift experiment shown in Fig. 1 was set up with a ³²P-labeled DNA fragment that contained a synthetic copy of

the *STE6* a-specific gene operator. Partially purified GRM from yeast cells (purified approximately 1,300-fold [see reference 4 for details]) provided a positive control for cooperativity. Low levels of the *MCMI* gene product (from *E. coli*) and purified $\alpha 2$ (from *E. coli* as described in reference 10) were used such that added separately, each occupied only a small fraction of the operators (Fig. 1, lanes 2 and 3, respectively). Figure 1, lane 5 shows the result obtained when the same quantities of Mcm1 protein and $\alpha 2$ were added together to the operator. Here, virtually all of the operator migrates in new, slower-mobility complexes; the fraction of operators occupied by both Mcm1 and $\alpha 2$ is considerably greater than that occupied by either protein added singly. This enhanced binding of the operator by Mcm1 and $\alpha 2$ together is very similar to that obtained upon addition of bona fide GRM (from yeast cells) and $\alpha 2$ together (Fig. 1, lane 4). Also as in the case of GRM, formation of the Mcm1- $\alpha 2$ -operator complex required the GRM-binding sequences of the operator. Formation of the complex can be competed against only by operators containing a functional GRM-binding site (data not shown). Given these results, we conclude that Mcm1 and $\alpha 2$ bind to the operator cooperatively.

Using a standard statistical thermodynamic analysis (for an example, see reference 1) of data obtained from densitometric scans of Fig. 1 and of autoradiographs from other experiments, we estimate that the *MCMI* gene product raises the affinity of $\alpha 2$ for the operator approximately 500-fold, corresponding to a cooperative interaction energy of 3 to 4 kcal (12 to 16 kJ)/mol. Additional experiments (data not shown) demonstrate that the *MCMI* gene product, like GRM, cobinds but does not cooperate with the purified C-terminal DNA-binding domain of $\alpha 2$. Thus, cooperative binding of the *MCMI* gene product and $\alpha 2$ requires the amino-terminal domain of $\alpha 2$ and is not simply an indirect consequence of contiguous binding. Because a truncated *MCMI* gene product was used in this experiment, we also know that the C-terminal 98 amino acids of Mcm1 are not required for cooperative binding with $\alpha 2$. It is not known why the Mcm1 hybrid product used in this experiment gives rise to two minor DNA-protein complexes in addition to the major complex (Fig. 1, lane 2). One likely possibility is that the Mcm1 preparation contains minor degradation products.

Because the Mcm1 protein produced in *E. coli* can provide

* Corresponding author.




FIG. 1. Gel mobility shift assay of protein-DNA complexes formed by adding purified $\alpha 2$, partially purified Mcm1 (made in *E. coli*), partially pure GRM, or combinations of these proteins to labeled operator. The protein-DNA complexes were separated from free DNA by mobility shift on native polyacrylamide gels and visualized by autoradiography. Lanes: 1, 10 μg of partially pure GRM (total protein) per ml; 2, 24 ng of partially pure Mcm1 per ml; 3, 0.24 μg of purified $\alpha 2$ per ml; 4, 10 μg of GRM and 0.24 μg of $\alpha 2$ per ml; 5, 24 ng of Mcm1 and 0.24 μg of $\alpha 2$ per ml. All reactions contained 4 ng of labeled operator and 1 μg of sheared salmon sperm DNA per ml. No operator binding, alone or cooperatively with $\alpha 2$, was observed in extracts of *E. coli* that did not contain *MCMI* (not shown).

both of the functions that define GRM, we performed a gel mobility shift experiment (Fig. 2) to test for the presence of Mcm1 in the GRM-operator complex formed using partially pure GRM from yeast cells. Polyclonal antiserum that recognizes the *MCMI* gene product (8) was incubated with GRM prior to the addition of ^{32}P -labeled operator. If the anti-Mcm1 antibodies recognize GRM, they would be expected either to prevent GRM binding to the operator or to form an antibody-GRM-operator complex that migrates more slowly than the GRM-operator complex. The latter result was obtained (Fig. 2). Addition of anti-Mcm1 antiserum gave rise to a shifted complex (lane 8) which migrated more slowly than that given by GRM plus operator alone (lane 5). This recognition is specific, since addition of preimmune or anti- $\alpha 2$ serum (Fig. 2, lanes 6 and 7, respectively) had no effect on the mobility of the GRM-operator complex. Thus, the anti-Mcm1 antibodies do recognize GRM, showing that the *MCMI* gene product (or a related protein) is part of the GRM-binding activity from yeast.

We next tested whether the protein recognized by the anti-Mcm1 antiserum was present in the cooperative three-part complex made up of $\alpha 2$, GRM, and the operator. To this end, purified $\alpha 2$ and partially purified GRM were incubated alone or with anti-Mcm1, anti- $\alpha 2$, or preimmune antiserum before addition of labeled operator DNA. The cooperative GRM- $\alpha 2$ -operator complex is shown in Fig. 2, lane 13; it migrated more slowly than either GRM (lane 5) or $\alpha 2$ (lane 9) alone bound to the operator. As expected, addition of either anti- $\alpha 2$ antiserum (lane 15) or anti-Mcm1 antiserum (lane 16) but not preimmune antiserum (lane 14) further reduced the mobility of the GRM- $\alpha 2$ -operator complex. This indicates that both the $\alpha 2$ protein and the *MCMI* gene product are present in the ternary complex.




FIG. 2. Gel mobility shift assay of protein-DNA complexes formed by adding purified $\alpha 2$, partially pure GRM, or both to labeled operator in the presence or absence of anti-Mcm1 or anti- $\alpha 2$ antiserum. The protein-DNA complexes were separated from free DNA by mobility shift on native polyacrylamide gels and visualized by autoradiography. Lanes: 1 through 4, no protein; 5 through 8, 11 μg of partially pure GRM (total protein) per ml; 9 through 12, 0.8 μg of purified $\alpha 2$ per ml; 13 through 16, 5.6 μg of partially pure GRM and 0.4 μg of purified $\alpha 2$ per ml; 3, 7, 11, and 15, anti- $\alpha 2$ antiserum; 4, 8, 12, and 16, anti-Mcm antiserum; 2, 6, 10, and 14, preimmune serum; 1, 5, 9, and 13, no antiserum added. All reactions contain 10 ng of labeled operator per ml, and no nonspecific DNA was added.

In this paper, we show that Mcm1 binds cooperatively with $\alpha 2$ to the operator and that a protein recognized by anti-Mcm1 antiserum is present in the GRM activity isolated from yeast cells. In addition, preliminary evidence suggests that the *MCMI* gene product is required for repression by $\alpha 2$ in yeast cells: α cells expressing a truncated *MCMI* gene are deficient in repression of at least one α -specific gene (R. Elble and B.-K. Tye, unpublished results). The results presented in this paper add to those previously obtained (see above) (4, 8) to strongly suggest that GRM is encoded by *MCMI* and that Mcm1 and $\alpha 2$ function together in vivo to bring about repression of transcription. It should be noted that recent work (3) indicates that *MCMI* encodes pheromone-receptor transcription factor (PTRF) that, with $\alpha 1$, coactivates the α -specific genes (a family of genes transcribed only in the α cell type of *S. cerevisiae*). Thus, it appears that as previously suggested (3, 4, 8), *MCMI* encodes a protein involved in either repression or activation, depending on its interaction with $\alpha 2$ or $\alpha 1$, respectively.

Mcm1 and $\alpha 2$ are members of two general classes of eucaryotic regulatory proteins. A portion of $\alpha 2$ has been shown to have homology with the homeodomain, a region found in many proteins known to specify cell identity (5, 11). Mcm1 has a region of strong homology with the serum response factor, a mammalian transcription activator that appears to be present in other eucaryotes as well (7). Therefore, we feel that it is reasonable to expect that gene regulation by pairwise combinations of homeodomain proteins and SRF-like proteins will be found in other cases.

We are grateful to T. deLange, K. Yamamoto, N. Hollingsworth, and I. Herskowitz for their critical reading and helpful comments on

the manuscript. We are indebted to A. Vershon for the purified $\alpha 2$ protein and for his valuable advice on protein purification. We thank M. Redd for advances in gel mobility shift techniques.

LITERATURE CITED

1. Akers, G. K., A. D. Johnson, and M. A. Shea. 1982. Quantitative model for gene regulation by λ phage repressor. Proc. Natl. Acad. Sci. USA **79**:1129–1133.
2. Atchison, M. L. 1988. Enhancers: mechanisms of action and cell specificity. Annu. Rev. Cell Biol. **4**:127–153.
3. Jarvis, E. E., K. L. Clark, and G. F. Sprague, Jr. 1989. The yeast transcription activator PRTF, a homolog of the mammalian serum response factor, is encoded by the *MCM1* gene. Genes Dev. **3**:936–945.
4. Keleher, C. A., C. Goutte, and A. D. Johnson. 1988. The yeast cell-type-specific repressor $\alpha 2$ acts cooperatively with a non-cell-type-specific protein. Cell **53**:927–936.
5. Laughon, A., and M. P. Scott. 1984. The sequence of a *Drosophila* segmentation gene: protein structure homology with DNA binding proteins. Nature (London) **310**:25–30.
6. Maine, G. T., P. Sinha, and B.-K. Tye. 1984. Mutants of *S. cerevisiae* defective in the maintenance of minichromosomes. Genetics **106**:365–385.
7. Norman, C., M. Runswick, R. Pollock, and R. Treisman. 1988. Isolation and properties of cDNA clones encoding SRF, a transcription factor that binds to the *c-fos* serum response element. Cell **55**:989–1003.
8. Passmore, S., R. Elble, and B.-K. Tye. 1989. A protein involved in minichromosome maintenance in yeast binds a transcriptional enhancer conserved in eukaryotes. Genes Dev. **3**:921–935.
9. Passmore, S., G. T. Maine, R. Elble, C. Christ, and B.-K. Tye. 1988. A *Saccharomyces cerevisiae* protein involved in plasmid maintenance is necessary for mating of *MAT α* cells. J. Mol. Biol. **204**:593–606.
10. Sauer, R. T., D. L. Smith, and A. D. Johnson. 1988. Domain structure, subunit organization, and operator-binding features of the yeast repressor $\alpha 2$. Genes Dev. **2**:807–816.
11. Shepherd, J. C. W., W. McGinnis, A. E. Carrasco, E. M. DeRobertis, and W. J. Gehring. 1984. Fly and frog homeo domains show homologies with yeast mating type regulatory proteins. Nature (London) **310**:70–71.