

## Differential regulation of NF- $\kappa$ B by elongation factors is determined by core promoter type

Liat Amir-Zilberstein<sup>1</sup>, Elena Ainbinder<sup>1</sup>, Leanne Toubé<sup>1</sup>, Yuki Yamaguchi<sup>2</sup>, Hiroshi Handa<sup>2</sup>  
and Rivka Dikstein<sup>1,\*</sup>

<sup>1</sup>Dept. of Biological Chemistry,  
The Weizmann Institute of Science,  
Rehovot 76100, Israel.

<sup>2</sup>Dept. of Biological Information,  
Tokyo Institute of Technology,  
Yokohama 226-8501, Japan.

\*Corresponding author

Tel: 972-8-9342117

Fax: 971-8-9344118

E-mail: [rivka.dikstein@weizmann.ac.il](mailto:rivka.dikstein@weizmann.ac.il)

**Running title:** Regulation of NF- $\kappa$ B genes by elongation factors

No. of characters (with spaces): 52,399

## Summary

NF- $\kappa$ B transcription factors activate genes important for immune response, inflammation and cell survival. P-TEFb and DSIF, positive and negative transcription elongation factors respectively, both regulate NF- $\kappa$ B induced transcription but the mechanism underlying their recruitment to NF- $\kappa$ B target genes is unknown. We show here that upon induction of NF- $\kappa$ B a subset of target genes is regulated differentially by either P-TEFb or DSIF. The regulation of these genes and their occupancy by these elongation factors are dependent on the NF- $\kappa$ B enhancer and the core promoter type. Converting a TATA-less to a TATA promoter switches the regulation of NF- $\kappa$ B from DSIF to P-TEFb. Accumulation or displacement of DSIF and P-TEFb is dictated by the formation of distinct initiation complexes (TFIID-dependent or independent) on the two types of core promoter. The underlying mechanism for the dissociation of DSIF from TATA promoters upon NF- $\kappa$ B activation involves the phosphorylation of Pol II by P-TEFb. The results highlight a regulatory link between the initiation and the elongation phases of the transcription reaction and broaden our comprehension of the NF- $\kappa$ B pathway.

## Introduction

Transcription of protein encoding genes by RNA polymerase II (pol II) is a multi-step process, each step being a target for regulation and critical for the production of mature mRNA (27, 29). A number of factors that control RNA pol II elongation have been characterized in recent years. Among these are the positive elongation factor P-TEFb, which induces Pol II processivity by facilitating the transition from the early to the late elongation phase (24), and two negative elongation factors, DSIF (DRB sensitivity inducing factor) (31) and NELF (negative elongation factor) (37). *In vitro*, P-TEFb alleviates transcription inhibition by DSIF (25, 32).

NF- $\kappa$ B is a transcription factor central to the cellular response to a broad range of extracellular signals including inflammatory cytokines, tumor promoters and chemotherapeutic agents. In response to these agents NF- $\kappa$ B induces the expression of cell cycle regulators, pro- and anti-apoptotic factors, inflammatory cytokines, chemokines, adhesion molecules, and many other factors (22). In unstimulated cells NF- $\kappa$ B is retained in the cytoplasm by I $\kappa$ B proteins. NF- $\kappa$ B activating signals trigger degradation of I $\kappa$ B and nuclear translocation of NF- $\kappa$ B which result in activation of responsive genes (14). A subset of early response genes that includes I $\kappa$ B $\alpha$  and A20, are themselves negative regulators of the NF- $\kappa$ B pathway and so form a negative feedback loop. Transcriptional control of these genes is likely to influence the strength and the duration of the inflammatory signal.

Induction of NF- $\kappa$ B target genes is remarkably fast and the mechanism underlying their rapid transcriptional activation has been previously investigated. It was found that the promoters of NF- $\kappa$ B regulated genes are bound by the general transcription machinery prior to NF- $\kappa$ B activation NF- $\kappa$ B, which increases the rate of the transcription cycles (re-initiation) rather than promoting pre-initiation complex formation (2). Further experiments with the A20 NF- $\kappa$ B target gene revealed that both the basal and the NF- $\kappa$ B induced transcription are repressed at the level of elongation. We identified the inhibitory factor as DSIF, which in this system acts without NELF (1). On the other hand NF- $\kappa$ B induced transcription of the IL-8 but

not the  $\text{I}\kappa\text{B}\alpha$  gene was shown to be regulated by the positive elongation factor P-TEFb (4, 18). Thus, NF- $\kappa$ B target genes are subjected to regulation by both positive and negative transcription elongation factors. However the mechanism underlying the differential control and recruitment of these factors to NF- $\kappa$ B target genes is currently unknown.

Here we investigated the regulation of NF- $\kappa$ B-mediated transcription by DSIF and P-TEFb. Our data revealed that DSIF attenuation of NF- $\kappa$ B is promoter dependent and requires the NF- $\kappa$ B response element to be in a context of a TATA-less core promoter, which, in turn, enhances DSIF occupancy upon NF- $\kappa$ B induction. By contrast, TATA-box-containing NF- $\kappa$ B promoters are not targeted for inhibition by DSIF and in these genes NF- $\kappa$ B diminishes DSIF occupancy. Remarkably, the core promoter also influences regulation and recruitment of the positive elongation factor P-TEFb, but inversely to DSIF. We found that the two core promoter types dictate formation of distinct initiation complexes that are activated by NF- $\kappa$ B, thereby linking the initiation machinery to elongation control. Thus, the core promoter type, via the formation of distinct initiation complexes, affects the extent of NF- $\kappa$ B activation by reducing or facilitating transcription elongation rate.

## Materials and Methods

### Plasmid constructions

The A20, A20 mNF- $\kappa$ Bs, 2 $\kappa$ B-A20, 2 $\kappa$ B(A20)- $\alpha$ -actin and DSIF RNAi 1 have been previously described (1, 2). The DSIF RNAi 2 and CDK9 RNAi were constructed according to Brummelkamp et al (6), using pSUPER plasmid and a synthetic oligonucleotides targeting the 5'-GTTTCATTGCCTACCAGTTC and 5'-CCAAAGCTTCCCCCTATAA sequences corresponding to the 784-802 position of DSIF p160 and the 358-376 position of cdk9 mRNAs respectively. The A20-TATA was constructed by replacing a PmlI-XmaI fragment from the A20 promoter with a double stranded oligo containing the TATA mutation: 5'-CCTACAACCCGTATAAAACTGAAACGGGGC; reverse 5'-GCCCCGTTTCAGTTTTATACGGGTTGTAGG. The promoters of the IP-10, RANTES and I $\kappa$ B $\alpha$  were amplified by PCR from genomic DNA and cloned in the promoter-less reporter gene pGL2-basic (Promega). The primers used are: IP-10 5'- CAAGGCACTCATCTGATTTTC; 5'- GACAAAGCTTCGGGATGTCTCTCAGCGGTG; RANTES 5'- CCTATGACCAGGATGAAAGC; 5'- AGCCAAGCTTAGAGGCTGTGCGAGGTCCAC; I $\kappa$ B $\alpha$  5'- AAGGCTCACTTGCAGAGGG; 5'- GGACTGCTGTGGGCTCTG. All the constructs were verified by sequencing.

### Transient transfection assays and chromatin immunoprecipitation

293T cells (human embryonic kidney fibroblasts) were maintained in Dulbecco's modified Eagle's medium supplemented with 10 % fetal calf serum. Transfections were performed using the standard CaPO<sub>4</sub> method. To avoid basal NF- $\kappa$ B activity, cells were kept from reaching confluence, and re-plated no more than nine times. For reporter assays, subconfluent cells were transfected in a 24-well plate using 1.1  $\mu$ g pSuper or DSIF RNAi plasmid, and 20 ng of the reporter plasmids, 1 ng RSV-Renilla, 10 ng CMV-GFP and 1 ng p65/RelA. 48h after transfection cells were harvested and their luciferase and renilla activities were measured. For the ChIP assay subconfluent cells in 100 mm dishes were transfected

with 0.2  $\mu$ g of the reporter plasmids and 24 hours later cells were left untreated or treated with TNF $\alpha$  for 1h (20 ng/ml) and fixed with formaldehyde for 10 minutes. Chromatin extract was then prepared from the cells and used for immunoprecipitation as described in (2). To correct for differences in transfection efficiency, the amount of input DNA was measured by PCR prior to the analysis of the immunoprecipitated DNAs and differences in the total amount of transfected DNA in the samples were normalized by adjusting the amount of precipitated DNA taken for the PCR analysis. The forward primer for each reporter is derived from the promoter and the reverse primer is from the luciferase gene 5'- CCATCCTCTAGAGGATAGAATG. The primers from the 1 kb upstream region of the A20 promoter that was used as control for specificity are: 5'-GGTTAGCTCCTTCGGTCCTC; 5'-CCGTATTGACGCCGGGCAAG. For the endogenous genes the amount of input DNA was first measured by PCR and differences in the input DNA in the samples were normalized by adjusting the amount of precipitated DNA taken for the PCR analysis. The primers used are: A20 promoter 5'- CAGCCCGACCCAGAGAGTCAC; 5'- CTTGGCCCGCCACGAA; A20 coding the same as for the RT-PCR. I $\kappa$ B $\alpha$  promoter 5'- AAGGCTCACTTGACAGAGGG; 5'- GGACTGCTGTGGGCTCTG; I $\kappa$ B $\alpha$  coding 5'-TCCTGAGCTCCGAGACTTTC; 5'- GTAGTTGGTAGCCTTCAGG; RANTES promoter 5'- CTTATGATACCGCCAATGC; 5'- GTGCGAGGTCCACGTGCTGTC; RANTES coding 5'-CACAGGTGAGAGGCCCTTCG; 5'- CAGCTGAACTTCTTCTCGCCC;  $\beta$ -actin promoter 5'-AAAGGAGGGGAGAGGGGGTAA; 5'- AAAGGCGAGGCTCTGTGCTC; BLR1 promoter 5'-CATTACAAGTTGTGAGCC; 5'- CATCAGTGCTAGTCAAGC. Pol II antibodies are from BAbCo, p65/RelA, CDK9 and TAF1 antibodies are from Santa Cruz Biotechnology, Inc. DSIF antibodies were previously described (1). The ChIP data were quantified by densitometric analysis using the Quantity One 1-D Analysis software (Bio-Rad).

#### RNA preparation and quantitative RT-PCR analysis

Human 293T cells in 100 mm plates were left untreated or treated with TNF $\alpha$  for 1h (20 ng/ml). Total RNA was prepared using the TRIzol reagent (GibcoBRL), according to the manufacturer's instructions. RNA preparations were treated with RQ1 DNase I (Promega) to avoid contamination of genomic DNA. First strand cDNA was synthesized from 1 $\mu$ g of total RNA using Oligo(dT)<sub>15</sub> primer, SuperScript II reverse transcriptase (Invitrogen). The PCR was performed in 20 $\mu$ l glass capillary tubes using a LightCycler System (Roche Molecular Biochemicals), equipped with a thermal cycler and real-time detector of fluorescence. The total cDNAs were amplified using LightCycler–FastStart DNA Master SYBR Green I kit (Roche Molecular Biochemicals), according to manufacturer's instructions. The oligonucleotides used for real time PCR are: human GAPDH 5'-CTGAGCTGAACGGGAAGCTC; 5'- CACCTGGTGCTCAGTGTAGC; A20 5'- GCTGCTGCCTCAGGGAAAGTC; 5'- CTCTTCTGTCCTTTTGGCCTC; I $\kappa$ B $\alpha$  5'- CCTTCCTCAACTTCCAGAACAACC; 5'- GGCTAAGTGTAGGCAGGTGTGGC. The primers for the RT-PCR analysis of the ts13 hamster cells are: A20 5'-CAAATGCTAAGAAGTTTGG; 5'-CTCTGTTAACAAGTGGAACAG; I $\kappa$ B $\alpha$  is the same as the human;  $\beta$ -actin 5'-CCCTGGAGAAGAGCTACGAGCTGCC; 5'- GCTTGCTGATCCACATCTGCTGGAAGG.

## Results

### Core promoter context determines DSIF attenuation of NF- $\kappa$ B-mediated transcription

The A20 gene is highly responsive to the cytokine TNF $\alpha$  through its activation of NF- $\kappa$ B. Treatment of HEK 293T cells with TNF $\alpha$  for one hour resulted in induction of A20 mRNA as determined by quantitative real-time PCR analysis (Fig. 1A). Upon transfection of these cells with DSIF RNAi there was a significant increase in the TNF $\alpha$ -induced A20 mRNA levels, suggesting that DSIF attenuates A20 induced transcription.

Next we determined the effect of DSIF knockdown on a luciferase reporter gene driven by the wild type or NF- $\kappa$ B-sites-mutated A20 promoter in the presence of co-transfected NF- $\kappa$ B protein p65/RelA. As expected the A20 promoter was activated by NF- $\kappa$ B whereas no induction was seen with the mutant A20 promoter (Fig. 1B). Consistent with previous results (1) down regulation of DSIF by RNAi (Fig. 1B right panel) enhanced the NF- $\kappa$ B induced transcription of the wild type but had no effect on the NF- $\kappa$ B mutated A20 promoter (Fig. 1B left panel) suggesting that NF- $\kappa$ B itself is required for the inhibition by DSIF.

To examine whether DSIF inhibitory activity is common to other NF- $\kappa$ B target genes we determined the effect of DSIF knockdown on luciferase activity driven by RANTES and IP-10 promoters in the presence of co-transfected NF- $\kappa$ B protein p65/RelA (Fig. 1C). Unlike the A20 promoter, the NF- $\kappa$ B induced activity of these promoters was not significantly affected by DSIF depletion. These results suggest that even though the inhibition of NF- $\kappa$ B by DSIF requires NF- $\kappa$ B, the effect is not general but promoter specific.

To understand the basis for the promoter-mediated regulation of NF- $\kappa$ B transcription by DSIF we set out to determine the region in the A20 promoter that in conjunction with NF- $\kappa$ B generates the inhibitory effect of DSIF. The A20 promoter has 6 Sp1 binding sites followed by two NF- $\kappa$ B binding sites in front of a TATA-less core promoter (Fig. 2A). To determine whether Sp1 plays a role in DSIF activity, the Sp1 sites were deleted leaving only the two NF- $\kappa$ B binding sites and the core promoter. The results (Fig. 2A) show that the A20 minimal promoter (2 $\kappa$ B-A20) retained the DSIF inhibitory effect (presented as the ratio between the activities in



the presence and absence of DSIF RNAi), indicating that Sp1 is not required for DSIF activity. Thus the differential regulation of NF- $\kappa$ B transcription by DSIF could be dependent either on the specific sequence of the NF- $\kappa$ B sites or on the core promoter. To test this we replaced the A20 core promoter with the core promoter of the  $\alpha$ -actin gene and examined the effect of DSIF RNAi on NF- $\kappa$ B activity (Fig. 2A). While the minimal A20 promoter (2 $\kappa$ B-A20) was sensitive to DSIF knockdown, the construct bearing the same two NF- $\kappa$ B sites in front of the heterologous  $\alpha$ -actin core promoter (2 $\kappa$ B(A20)- $\alpha$ -actin) was unaffected by DSIF knockdown (Fig. 2A) (the amount of NF- $\kappa$ B used in these experiments is well below that required to achieve maximal activation). This suggests that it is the context of the core promoter that is important for attenuation of NF- $\kappa$ B by DSIF.

Analysis of the core promoters of the genes used in Fig.1 and Fig. 2A revealed that the DSIF responsive promoter A20 is TATA-less whereas the less sensitive promoters IP-10, RANTES and 2 $\kappa$ B- $\alpha$ -actin contain a canonical TATA box (TATAWA) at the appropriate location (-25 to -30 relative to the transcription start site). To confirm that it is indeed the nature of the core promoter that determines the DSIF inhibitory activity of NF- $\kappa$ B, the TATA-less A20 promoter was converted into a canonical TATA promoter by substituting two nucleotides in the core promoter as shown in Fig. 2B, and the effect of two distinct DSIF RNAis were analyzed in the presence of co-transfected p65/RelA (using a concentration well below that required to achieve maximal activation). The TATA box mutation (A20-TATA) increased the basal activity but not the activation fold by NF- $\kappa$ B (Fig. 2C). Expression of each DSIF p160 RNAi enhanced the activity of the wild type A20 but not of the A20-TATA mutant by ~2 fold. The difference in the effect of DSIF on A20 and A20-TATA is statistically significant ( $p=0.0025$ ). Thus the inhibitory effect of DSIF on NF- $\kappa$ B induced transcription requires NF- $\kappa$ B site(s) to be in the context of a TATA-less core promoter.

### **DSIF occupancy is enhanced by NF- $\kappa$ B in a core promoter dependent manner**

If the attenuation of NF- $\kappa$ B-induced transcription by DSIF is a direct effect it should involve association of DSIF with the promoter in a manner dependent on NF- $\kappa$ B and the core promoter. We therefore transfected 293T cells with the different A20 reporter genes, treated them with TNF $\alpha$  for 1 hour to induce NF- $\kappa$ B activity, and then subjected them to chromatin immunoprecipitation (ChIP) assay using antibodies against RNA Pol II, p65/RelA, the p160 subunit of DSIF and a non-relevant antibody as a control. After reverse cross-linking PCR reactions were performed with primers corresponding to the 5' end of A20 promoter and to the beginning of the luciferase gene. As shown in Fig. 3A under basal conditions Pol II and DSIF constitutively bind the A20 wild type, and the A20-TATA mutant. Upon TNF $\alpha$  treatment NF- $\kappa$ B p65/RelA associates with the A20 wild type and the A20-TATA mutant and there is a concomitant increase in Pol II occupancy. DSIF occupancy on the other hand, is enhanced only in the wild type A20 and the 2 $\kappa$ B-A20 but is reduced in the TATA containing promoters A20-TATA and 2 $\kappa$ B- $\alpha$ -actin. Enhancement of Pol II and DSIF occupancies is NF- $\kappa$ B dependent as an A20 promoter bearing mutations in the two NF- $\kappa$ B binding sites which prevented binding of p65/RelA after TNF $\alpha$  induction (Fig. 3A bottom panels), failed to enhance Pol II and DSIF binding. The association of Pol II, p65/RelA and DSIF with the promoter is specific as a non-promoter sequence located 1 kb upstream was not enriched by them (data not shown).

To confirm that the endogenous A20 gene is also directly regulated by DSIF upon NF- $\kappa$ B induction we performed ChIP assays on the native A20 gene. Fig. 3B shows that upon NF- $\kappa$ B induction by TNF $\alpha$ , DSIF occupancy of the A20 promoter is increased as found with the transfected promoter. Enhancement of DSIF occupancy is specific to NF- $\kappa$ B as DSIF occupancy at the non-inducible housekeeping gene  $\beta$ -actin is unaffected by NF- $\kappa$ B induction. To confirm the specificity of the interaction of Pol II, p65 and DSIF with the A20 promoter, a similar analysis was performed with the promoter of the BLR1 gene, a B-cell-specific NF- $\kappa$ B target gene (22) that is not expressed in Jurkat T cells. This promoter was not enriched by any

of these factors. Together the results suggest that the effect of DSIF on the A20 gene upon NF- $\kappa$ B induction correlates directly promoter occupancy in a manner dependent on NF- $\kappa$ B and the core promoter.

### **The core promoter has opposite effects on DSIF and P-TEFb occupancies upon NF- $\kappa$ B induction**

Previous studies demonstrated that the positive elongation factor P-TEFb alleviates inhibition of transcription elongation by DSIF *in vitro* (25, 32). To determine the interplay between DSIF and P-TEFb in the regulation of the A20 gene we compared P-TEFb and DSIF occupancy of the various A20 promoter derivatives before and after stimulation of NF- $\kappa$ B by TNF $\alpha$  (Fig. 4). The results revealed that on the TATA-less A20 reporter, P-TEFb occupancy is clearly detectable under basal conditions but in contrast to DSIF, it is down regulated upon NF- $\kappa$ B induction (A20 panels). This effect is NF- $\kappa$ B dependent since P-TEFb occupancy is unaffected when the NF- $\kappa$ B sites are mutated (A20-mNF- $\kappa$ Bs panels). P-TEFb associates specifically with the promoter region as in a non-promoter sequence located 1 kb upstream P-TEFb was not detected (data not shown). Notably, converting the A20 to a TATA promoter reverses the effect of NF- $\kappa$ B: P-TEFb occupancy is still clearly detected before stimulation as in the wild type but now its occupancy is unchanged after NF- $\kappa$ B induction as opposed to DSIF levels which are reduced (A20-TATA panels). Thus, in the NF- $\kappa$ B induced system switching the core promoter type inverts the ratio between positive and negative elongation factors. This suggests that the ratio between functionally opposing elongation factors is important for elongation control and can be modulated by core promoter type.

P-TEFb is known to phosphorylate serine 2 of Pol II CTD heptapeptides and to facilitate its processivity. We therefore assessed the occupancy of unphosphorylated and Serine 2 phosphorylated forms of Pol II as well as of P-TEFb and DSIF at the promoter and at downstream coding sequences of endogenous A20 gene in Jurkat T cells treated with TNF $\alpha$  for one hour (Fig. 4B). Consistent with the results from transfected promoters, P-TEFb and

serine 2 Pol II, which are detected under basal conditions, are clearly reduced in the A20 promoter upon  $\text{TNF}\alpha$  induction (A20 promoter panels) as opposed to the unphosphorylated Pol II and DSIF which are clearly increased. Within the A20 gene Pol II is present only in its Serine 2 phosphorylated form, along with P-TEFb and DSIF under basal conditions. Following  $\text{TNF}\alpha$  treatment substantial amounts of unphosphorylated Pol II are detected whereas P-TEFb is reduced and DSIF is increased. Surprisingly the Serine 2 phosphorylated form of pol II does not fall in spite of P-TEFb release, possibly because a small fraction of the large quantity of induced Pol II is phosphorylated by the residual P-TEFb. The effect of  $\text{NF-}\kappa\text{B}$  then, on the occupancies of DSIF and P-TEFb on the promoter continues into the coding region indicating its relevance to the elongation phase of the A20 gene.

#### **P-TEFb regulation of $\text{NF-}\kappa\text{B}$ is dependent on a TATA box**

The P-TEFb ChIP experiments (Fig. 4A) suggest that P-TEFb, like DSIF, regulates  $\text{NF-}\kappa\text{B}$  in a core promoter dependent manner. To examine this 293T cells were transfected with the A20 wild type and A20-TATA mutant reporter genes and 24 hours later were treated with either  $\text{TNF}\alpha$  alone or with  $\text{TNF}\alpha$  together with increasing amounts of the drug DRB, a potent inhibitor of P-TEFb (24). The results (Fig. 5) show that  $\text{TNF}\alpha$  stimulated the luciferase activity of both reporters by more than 3-fold but the induced activity of the A20-TATA is more sensitive to inhibition by DRB than the wild type A20.

To examine further the role of P-TEFb in  $\text{NF-}\kappa\text{B}$ -mediated transcription we used RNAi to reduce the endogenous P-TEFb levels. The wild type A20 and the A20-TATA mutant reporter genes were transfected into 293T cells together with a plasmid directing the expression of RNAi specific for the cdk9 subunit of P-TEFb. Analysis of cdk9 expression by immunoblot shows its specific depletion from cells transfected with cdk9 RNAi but not from cells transfected with the parental plasmid (Fig 5B). Akin to the effect of DRB, down-regulation of cdk9 had no effect on the  $\text{NF-}\kappa\text{B}$ -activated transcription of the wild A20 promoter but it significantly reduced  $\text{NF-}\kappa\text{B}$  induction of the A20-TATA promoter. These findings indicate that

P-TEFb is particularly important for TATA containing NF- $\kappa$ B target genes and are consistent with the ChIP assay data that the P-TEFb/DSIF ratio on these promoters is increased upon NF- $\kappa$ B induction. In agreement with this model, the IL-8 gene, previously shown to be regulated by P-TEFb upon NF- $\kappa$ B induction (4, 18) has a TATA-box promoter whereas the TATA-less I $\kappa$ B $\alpha$  is not regulated by P-TEFb (18).

**Differential occupancy of other NF- $\kappa$ B target genes by P-TEFb and DSIF is correlated with the core promoter type**

To gain further support for the dependency of DSIF and P-TEFb occupancies on the core promoter type upon NF- $\kappa$ B induction, we examined two additional NF- $\kappa$ B responsive genes of which one is TATA-less (I $\kappa$ B $\alpha$ ) and the second has a canonical TATA element (RANTES). These genes were analyzed by ChIP before and after one hour treatment with TNF $\alpha$  using antibodies against unphosphorylated (Pol II) and Serine 2 phosphorylated (Ser2) forms of Pol II, DSIF or non-relevant control antibodies. PCR reactions were performed with primers corresponding to the promoter region (promoter) and an internal region of the gene (coding). As shown in Fig. 6 in the TATA-less I $\kappa$ B $\alpha$  promoter and gene unphosphorylated Pol II and DSIF occupancies are enhanced but the promoter serine 2 Pol II is reduced upon TNF $\alpha$  induction, as observed with A20. With the TATA-containing RANTES promoter the effects are reversed, DSIF is released and serine 2 Pol II is enhanced by NF- $\kappa$ B. In the coding region of RANTES gene, both the unphosphorylated and serine 2 phosphorylated forms of Pol II are increased in the promoter and the coding sequences, but DSIF is not diminished from the coding sequences upon TNF $\alpha$  induction. The lack of correlation of DSIF occupancy between the promoter and the coding argues that in the RANTES gene DSIF regulation during elongation is not mediated by the promoter. This regulation is likely to involve additional elements of the elongating Pol II which we don't understand. We now refer to this observation in the text.

Together these results strengthen the notion that differential occupancy by DSIF and P-TEFb during the initial stages of transcription, and by and large during elongation, is dependent on core promoter type.

### **DSIF occupancy is counteracted by serine 2 phosphorylation of Pol II**

Given that upon NF- $\kappa$ B activation DSIF and P-TEFb occupancies are mutually exclusive and that in vitro P-TEFb relieves the inhibitory effect of DSIF (25, 32) it is possible that DSIF NF- $\kappa$ B-induced accumulation or dissociation upon NF- $\kappa$ B activation is the result of, respectively, P-TEFb release or recruitment. Alternatively, DSIF may be recruited or released by NF- $\kappa$ B itself. However whereas P-TEFb has been shown to interact directly with the NF- $\kappa$ B protein p65/RelA (4), we failed to detect any direct interaction between p65/RelA and DSIF in vitro or in cell extracts, suggesting that the latter possibility is less likely. To examine the role of P-TEFb in DSIF occupancy, we performed ChIP experiments on TNF $\alpha$  induced Jurkat T cells treated with the P-TEFb inhibitor DRB, using antibodies against DSIF and the serine 2 phosphorylated form of Pol II (Fig. 7). Analysis of the TATA promoter RANTES revealed that the TNF $\alpha$ -induced serine 2 phosphorylation by P-TEFb is diminished by DRB (compare TNF $\alpha$  to TNF $\alpha$ +DRB lanes). This inhibition of P-TEFb activity resulted in failure to displace DSIF from the promoter. By contrast DRB had no effect on DSIF accumulation in the I $\kappa$ B $\alpha$  promoter. Thus it is most likely that the reduced DSIF occupancy upon NF- $\kappa$ B activation in TATA-containing promoters is a consequence of P-TEFb recruitment.

Notably under basal conditions DSIF occupies the RANTES promoter in the absence of the unphosphorylated or serine 2 phosphorylated forms of Pol II (Fig. 6 and 7). As DSIF is recruited to genes via its association with Pol II we checked for the presence of another form of Pol II, the serine 5 phosphorylated form. Indeed we found that serine 5 phosphorylated Pol II occupies this promoter exclusively and that this form disappears upon stimulation by TNF $\alpha$  (data not shown).

## The core promoter type determines the nature of initiation complex activated by NF- $\kappa$ B

It is well established that enhancers and core promoters direct the initiation step of transcription. The experiments described above suggest that at least for NF- $\kappa$ B target genes, the core promoter type is also involved in regulation of transcription elongation. Given that the core promoter is the site on which the transcription initiation complex assembles, we reasoned that formation of different initiation complexes on the two types of core promoter may dictate recruitment or displacement of distinct elongation factors. Therefore we wanted to determine the nature of the initiation complex that is formed on the different NF- $\kappa$ B target genes. Our experiments were based on yeast studies in which, depending on the core promoter structure, initiation complex assembly is mediated by either TFIID or SAGA (5, 8, 13, 15, 20). To examine the role of the core promoter in determining the type of the transcription initiation complex activated by NF- $\kappa$ B, we analyzed the occupancy of A20 promoter derivatives by TAF1, a TFIID-specific subunit not present in the mammalian SAGA related complexes STAGA, TFTC and PCAF (12, 19, 21, 35). 293T cells were transfected with the A20 wild type or A20-TATA promoter and 24 hours later treated with TNF $\alpha$  for 1 hour to induce NF- $\kappa$ B, and then subjected to ChIP assay using antibodies against TAF1, RNA Pol II and a non-relevant antibody. The results (Fig. 8A) show that TAF1 occupies the the TATA-less A20 promoter before and after NF- $\kappa$ B stimulation whereas it is completely absent from the A20-TATA. The presence of pol II and TAF1 under basal conditions is consistent with our previous findings that the basal transcription machinery is already assembled in rapidly-induced NF- $\kappa$ B target genes (2).

Next, we determined TAF1 and Pol II occupancy of the endogenous TATA-less A20 and I $\kappa$ B $\alpha$  genes and of the TATA containing RANTES gene in Jurkat T cells that were treated with TNF $\alpha$  for 1 hour. TAF1 was found to be specifically associated with the A20 and I $\kappa$ B $\alpha$  promoters but not with the coding regions of these genes before and after TNF $\alpha$  induction (Fig. 8B). By contrast TAF1 was undetectable on the RANTES promoter under both basal and TNF $\alpha$  stimulated conditions. These findings suggest that the core promoter sequence of NF-

$\kappa$ B target genes controls the pathway of the transcription initiation complex assembly, which can be either TFIID-dependent (TATA-less) or TFIID-independent (TATA).

To gain further support for the idea of a the differential requirement for TFIID in NF- $\kappa$ B transcription we used the temperature sensitive hamster cell line ts13 in which TAF1 contains a point mutation that renders TFIID inactive at 39.5°C. These cells were co-transfected with NF- $\kappa$ B dependent reporter plasmids and the NF- $\kappa$ B protein p65/RelA. As shown in Fig. 8C, activation by NF- $\kappa$ B of the TATA-less A20 and I $\kappa$ B promoters decreased at the non-permissive temperature, 39.5°C, compared to the permissive temperature, 32°C, demonstrating TFIID-dependency. Activation of the TATA-containing RANTES promoter, on the other hand, was unaffected by changing from the permissive to the non-permissive temperature. Moreover, modifying the TATA-less sequence of the A20 promoter to canonical TATA (A20-TATA) abolished its dependency on TFIID.

We also assessed the involvement of TFIID in the transcription of endogenous NF- $\kappa$ B target genes by measuring the mRNA levels of the A20 and I $\kappa$ B $\alpha$  genes in the hamster ts13 before and after treatment with TNF $\alpha$  for one-hour (the endogenous RANTES gene, whose promoter was analyzed above, is not expressed in these cells). Quantitative RT-PCR measurements (Fig. 8D) showed a significant decrease in the TNF $\alpha$ -induced mRNA levels of A20 and I $\kappa$ B $\alpha$  at the non-permissive temperature. The mRNA of the  $\beta$ -actin gene was unchanged and we used it to normalized the results.

Considering the correlation between the core promoter structure and differential recruitment of DSIF and P-TEFb we can conclude that DSIF attenuation of NF- $\kappa$ B is dependent on the initiation complex assembled via TFIID whereas P-TEFb regulation of NF- $\kappa$ B requires TAF-independent assembly of the initiation complex.



## Discussion

Two types of DNA element, enhancers and the core promoter, regulate transcription. Several studies have indicated that specific combinations of these elements play a regulatory role in transcription (7, 9, 10, 15, 20, 28, 30, 33, 34). Such combinations are considered to be important for the initiation step of transcription. The data presented in this study demonstrate the importance of arrangement of enhancer with a core promoter type in gene specific effects of elongation regulatory factors. Our results thus reveal new links between the initiation and the elongation phases of the transcription reaction that are relevant to gene regulation. Thus it appears that the genetic information encoded by transcription regulatory regions affects more steps of the transcription cycle than initially believed. A role of the promoter in RNA Pol II processivity was reported for HIV-LTR and c-myc transcription (17, 38) where it was found that a paused but not processive transcription is dependent on an intact TATA box, though the basis for the TATA box requirement was not explored. Our findings extend those previous studies by showing that in the NF- $\kappa$ B pathway the TATA box is required for NF- $\kappa$ B-induced facilitation of elongation by P-TEFb upon NF- $\kappa$ B activation. Given the involvement of P-TEFb in TAT activation of HIV-LTR elongation, it remains to be seen whether their findings are linked to ours.

Specifically we demonstrated that attenuation of NF- $\kappa$ B induced transcription by the negative elongation factor DSIF occurs in genes carrying NF- $\kappa$ B binding sites in the context of a TATA-less core promoter. This specific configuration of the promoter is also responsible for enhanced recruitment of DSIF to these genes upon NF- $\kappa$ B induction. Contrary to TATA-less NF- $\kappa$ B target genes, transcription of TATA-box containing genes is not inhibited by DSIF and DSIF is actually lost from these genes upon NF- $\kappa$ B induction. In addition, the core promoter also influences gene occupancy and regulation by the positive elongation factor P-TEFb in a manner that correlates inversely with that of DSIF. The finding that activation of the A20 and I $\kappa$ B $\alpha$  genes results in a large increase in hypo-phosphorylated pol II within the coding region but no increase of ser-2 phosphorylated pol II was surprising and suggests that in

some genes transcription elongation occurs by a mechanism that does not involve ser-2 phosphorylation. This result adds to growing number of genes whose transcription is reported to be independent of P-TEFb (11, 18).

The mechanism by which the elongation factors DSIF and P-TEFb differentially regulate NF- $\kappa$ B target genes involves both NF- $\kappa$ B and the initiation complex that is formed on the different types of core promoter. When the initiation complex is assembled via TFIID on TATA-less promoters, P-TEFb is released upon NF- $\kappa$ B induction. As a consequence DSIF occupancy is enhanced along with the unphosphorylated form of Pol II. By contrast when the initiation complex is formed by a TFIID-independent pathway on TATA containing promoters, reminiscent of the SAGA complex in yeast (5, 8, 15, 20), NF- $\kappa$ B recruits or retains P-TEFb which then phosphorylates Pol II CTD on serine 2, resulting in DSIF displacement.

How does differential regulation of NF- $\kappa$ B target genes by positive and negative elongation factors affect its biological activity? DSIF has been implicated in the coordination of transcription with mRNA capping (16, 23). Whether this function of DSIF plays a role in activation of NF- $\kappa$ B target genes is yet to be determined. However the increase in the level of the luciferase protein in DSIF RNAi treated cells would seem to indicate that the luciferase mRNA has been capped. DSIF attenuates an important subset of NF- $\kappa$ B-induced genes, (A20 and I $\kappa$ B $\alpha$ ). Although the inhibitory effect of DSIF on these genes is moderate (~2 fold) it is likely to have significant biological impact. NF- $\kappa$ B is latent in the cytoplasm and translocates into the nucleus upon activation to induce target genes. Among those are A20 and I $\kappa$ B $\alpha$  which act to terminate NF- $\kappa$ B activity in a negative feedback loop. Thus, a two-fold inhibition of these genes by DSIF may alter the duration and the extent of the NF- $\kappa$ B activity.

In addition to the genes analyzed here there are other genes induced by NF- $\kappa$ B in different cells. We analyzed the type of core promoter of and expression pattern of 48 documented NF- $\kappa$ B target genes (22) for which the transcription initiation site is known. A TATA-box present in the core promoter of 32 of these genes (data not shown), the majority of which (26/32) are cell type specific (expressed in some but not all cell types). On the other

hand most of the TATA-less genes (14/16) are ubiquitously expressed (expressed in all cell types). The overrepresentation of TATA-less promoters among ubiquitously expressed target genes and TATA promoters among cell type specific target genes is in agreement with the genome-wide distribution of these two groups of genes (26). Interestingly, genes that are activated by NF- $\kappa$ B to modulate its own signaling pathway such as A20, I $\kappa$ B $\alpha$ , cIAP2, TRAF1, TRAF2, NF- $\kappa$ B1 (p105) and cRel are predominantly in the group of the TATA-less target genes. Given the dependency of DSIF and P-TEFb on the core promoter type, it can be predicted that most of the genes that mediate the physiological response of NF- $\kappa$ B including cytokines, chymokines and adhesion molecules are controlled by P-TEFb whereas genes involved in the signaling pathway of NF- $\kappa$ B are attenuated by DSIF.

DSIF is a positive and negative elongation regulatory factor and the mechanisms of these distinct functions seem to be different. The positive activity of DSIF is facilitated by P-TEFb which phosphorylates DSIF (36), and on DSIF positively regulated hsp70 and c-fos genes co-occupancy of P-TEFb and DSIF was observed (3, 36). On the other hand the negative activity of DSIF is actually counteracted by P-TEFb (25, 32) and this is consistent with our findings of their mutually exclusive regulation and occupancy in the NF- $\kappa$ B pathway. Moreover, DSIF inhibition is dependent on NF- $\kappa$ B and a TATA-less promoter in contrast to its positively regulated genes c-fos and hsp70 genes which are driven by a TATA promoter.

What then is the function of DSIF and P-TEFb in NF- $\kappa$ B-mediated transcription? NF- $\kappa$ B is a powerful transcription factor responsible for activation of many genes in response to a variety of external signals. For any particular biological response, the desired level of activation of different genes by the same activated NF- $\kappa$ B may not be equal. Reducing or enhancing the processivity of the activated Pol II via positive and negative elongation factors is a mechanism that can tune the level of activation by the same NF- $\kappa$ B up or down in a regulated fashion. The differential effect of positive and negative elongation factors on NF- $\kappa$ B target genes that belong to functionally distinct groups may be particularly important for modulating the strength and the duration of the inflammatory response.

## References

1. **Ainbinder, E., L. Amir-Zilberstein, Y. Yamaguchi, H. Handa, and R. Dikstein.** 2004. Elongation inhibition by DRB sensitivity-inducing factor is regulated by the A20 promoter via a novel negative element and NF-kappaB. *Mol Cell Biol* **24**:2444-54.
2. **Ainbinder, E., M. Revach, O. Wolstein, S. Moshonov, N. Diamant, and R. Dikstein.** 2002. Mechanism of rapid transcriptional induction of tumor necrosis factor alpha-responsive genes by NF-kappaB. *Mol Cell Biol* **22**:6354-62.
3. **Andrulis, E. D., E. Guzman, P. Doring, J. Werner, and J. T. Lis.** 2000. High-resolution localization of Drosophila Spt5 and Spt6 at heat shock genes in vivo: roles in promoter proximal pausing and transcription elongation. *Genes Dev.* **14**:2635-2649.
4. **Barboric, M., R. M. Nissen, S. Kanazawa, N. Jabrane-Ferrat, and B. M. Peterlin.** 2001. NF-kappaB binds P-TEFb to stimulate transcriptional elongation by RNA polymerase II. *Mol Cell* **8**:327-37.
5. **Basehoar, A. D., S. J. Zanton, and B. F. Pugh.** 2004. Identification and distinct regulation of yeast TATA box-containing genes. *Cell* **116**:699-709.
6. **Brummelkamp, T. R., R. Bernards, and R. Agami.** 2002. A system for stable expression of short interfering RNAs in mammalian cells. *Science* **296**:550-555.
7. **Butler, J. E., and J. T. Kadonaga.** 2001. Enhancer-promoter specificity mediated by DPE or TATA core promoter motifs. *Genes Dev* **15**:2515-9.
8. **Cheng, J. X., M. Floer, P. Ononaji, G. Bryant, and M. Ptashne.** 2002. Responses of four yeast genes to changes in the transcriptional machinery are determined by their promoters. *Curr Biol* **12**:1828-32.
9. **Conkright, M. D., E. Guzman, L. Flechner, A. I. Su, J. B. Hogenesch, and M. Montminy.** 2003. Genome-wide analysis of CREB target genes reveals a core promoter requirement for cAMP responsiveness. *Mol Cell* **11**:1101-8.
10. **Das, G., C. S. Hinkley, and W. Herr.** 1995. Basal promoter elements as a selective determinant of transcriptional activator function. *Nature* **374**:657-60.
11. **Gomes, N. P., G. Bjerke, B. Llorente, S. A. Szostek, B. M. Emerson, and J. M. Espinosa.** 2006. Gene-specific requirement for P-TEFb activity and RNA polymerase II phosphorylation within the p53 transcriptional program. *Genes Dev.* **20**:601-612
12. **Grant, P. A., D. Schieltz, M. G. Pray-Grant, D. J. Steger, J. C. Reese, J. R. Yates, 3rd, and J. L. Workman.** 1998. A subset of TAF(II)s are integral components of the SAGA complex required for nucleosome acetylation and transcriptional stimulation. *Cell* **94**:45-53.
13. **Huisinga, K. L., and B. F. Pugh.** 2004. A genome-wide housekeeping role for TFIID and a highly regulated stress-related role for SAGA in *Saccharomyces cerevisiae*. *Mol Cell* **13**:573-85.
14. **Karin, M., and A. Lin.** 2002. NF-kappaB at the crossroads of life and death. *Nat Immunol* **3**:221-7.
15. **Li, X. Y., S. R. Bhaumik, X. Zhu, L. Li, W. C. Shen, B. L. Dixit, and M. R. Green.** 2002. Selective recruitment of TAFs by yeast upstream activating sequences. Implications for eukaryotic promoter structure. *Curr Biol* **12**:1240-4.

16. **Lindstrom, D. L., S. L. Squazzo, N. Muster, T. A. Burckin, K. C. Wachter, C. A. Emigh, J. A. McCleery, J. R. Yates, 3rd, and G. A. Hartzog.** 2003. Dual roles for Spt5 in pre-mRNA processing and transcription elongation revealed by identification of Spt5-associated proteins. *Mol Cell Biol* **23**:1368-78.
17. **Lu, X., T. M. Welsh, and B. M. Peterlin.** 1993. The human immunodeficiency virus type 1 long terminal repeat specifies two different transcription complexes, only one of which is regulated by Tat. *J Virol* **67**:1752-60.
18. **Luecke, H. F., and K. R. Yamamoto.** 2005. The glucocorticoid receptor blocks P-TEFb recruitment by NFkappaB to effect promoter-specific transcriptional repression. *Genes Dev* **19**:1116-27.
19. **Martinez, E., T. K. Kundu, J. Fu, and R. G. Roeder.** 1998. A human SPT3-TAFII31-GCN5-L acetylase complex distinct from transcription factor IID. *J Biol Chem* **273**:23781-5.
20. **Mencia, M., Z. Moqtaderi, J. V. Geisberg, L. Kuras, and K. Struhl.** 2002. Activator-specific recruitment of TFIID and regulation of ribosomal protein genes in yeast. *Mol Cell* **9**:823-33.
21. **Ogryzko, V. V., T. Kotani, X. Zhang, R. L. Schiltz, T. Howard, X. J. Yang, B. H. Howard, J. Qin, and Y. Nakatani.** 1998. Histone-like TAFs within the PCAF histone acetylase complex. *Cell* **94**:35-44.
22. **Pahl, H. L.** 1999. Activators and target genes of Rel/NF-kappaB transcription factors. *Oncogene* **18**:6853-66.
23. **Pei, Y., and S. Shuman.** 2002. Interactions between fission yeast mRNA capping enzymes and elongation factor Spt5. *J Biol Chem* **277**:19639-48.
24. **Price, D. H.** 2000. P-TEFb, a cyclin-dependent kinase controlling elongation by RNA polymerase II. *Mol Cell Biol* **20**:2629-34.
25. **Renner, D. B., Y. Yamaguchi, T. Wada, H. Handa, and D. H. Price.** 2001. A highly purified RNA polymerase II elongation control system. *J Biol Chem* **276**:42601-9.
26. **Schug, J., W. P. Schuller, C. Kappen, J. M. Salbaum, M. Bucan, and C. J. Stoeckert, Jr.** 2005. Promoter features related to tissue specificity as measured by Shannon entropy. *Genome Biol* **6**:R33.
27. **Shilatifard, A., R. C. Conaway, and J. W. Conaway.** 2003. The RNA polymerase II elongation complex. *Annu Rev Biochem* **72**:693-715.
28. **Simon, M. C., T. M. Fisch, B. J. Benecke, J. R. Nevins, and N. Heintz.** 1988. Definition of multiple, functionally distinct TATA elements, one of which is a target in the hsp70 promoter for E1A regulation. *Cell* **52**:723-9.
29. **Sims, R. J., 3rd, R. Belotserkovskaya, and D. Reinberg.** 2004. Elongation by RNA polymerase II: the short and long of it. *Genes Dev* **18**:2437-68.
30. **Taylor, I. C., and R. E. Kingston.** 1990. Factor substitution in a human HSP70 gene promoter: TATA-dependent and TATA-independent interactions. *Mol Cell Biol* **10**:165-75.
31. **Wada, T., T. Takagi, Y. Yamaguchi, A. Ferdous, T. Imai, S. Hirose, S. Sugimoto, K. Yano, G. A. Hartzog, F. Winston, S. Buratowski, and H. Handa.** 1998. DSIF, a novel transcription elongation factor that regulates RNA polymerase II processivity, is composed of human Spt4 and Spt5 homologs. *Genes Dev* **12**:343-56.
32. **Wada, T., T. Takagi, Y. Yamaguchi, D. Watanabe, and H. Handa.** 1998. Evidence that P-TEFb alleviates the negative effect of DSIF on RNA polymerase II-dependent transcription in vitro. *Embo J* **17**:7395-403.

33. **Wang, W. D., and J. D. Gralla.** 1991. Differential ability of proximal and remote element pairs to cooperate in activating RNA polymerase II transcription. *Mol Cell Biol* **11**:4561-71.
34. **Wefald, F. C., B. H. Devlin, and R. S. Williams.** 1990. Functional heterogeneity of mammalian TATA-box sequences revealed by interaction with a cell-specific enhancer. *Nature* **344**:260-2.
35. **Wieczorek, E., M. Brand, X. Jacq, and L. Tora.** 1998. Function of TAF(II)-containing complex without TBP in transcription by RNA polymerase II. *Nature* **393**:187-91.
36. **Yamada, T., Y. Yamaguchi, N. Inukai, S. Okamoto, T. Mura, and H. A.Handa.** 2006. P-TEFb-mediated phosphorylation of hSpt5 C-terminal repeats is critical for processive transcription elongation. *Mol Cell* **21**:227-237.
37. **Yamaguchi, Y., T. Takagi, T. Wada, K. Yano, A. Furuya, S. Sugimoto, J. Hasegawa, and H. Handa.** 1999. NELF, a multisubunit complex containing RD, cooperates with DSIF to repress RNA polymerase II elongation. *Cell* **97**:41-51.
38. **Yankulov, K., J. Blau, T. Purton, S. Roberts, and D. L. Bentley.** 1994. Transcriptional elongation by RNA polymerase II is stimulated by transactivators. *Cell* **77**:749-59.

ACCEPTED

## **Acknowledgements**

We thank Dr. Sandra Moshonov for critical reading and editing of the manuscript and Nadav Bar and Rofa Elfakess for their assistance throughout the project. This work was supported by grants from the Israel Cancer Research Foundation and the Israel Science Foundation.

ACCEPTED

## Figure legends

**Figure 1: Attenuation of NF- $\kappa$ B mediated transcription by DSIF.** **A.** 293T cells were transfected with either pSuper or DSIF p160 RNAi and 48 hours post-transfection cells were treated with TNF $\alpha$  for 1 hour. Total RNA was extracted and subjected to a quantitative RT-PCR for the A20 and GAPDH genes mRNAs, using Light Cycler. The data represent the means and standard deviations of 3 independent experiments. Representative immunoblot verifying down regulation of p160 DSIF is shown in the right panel. **B.** 293T cells were transfected with either wild type A20 or A20-mNF- $\kappa$ B (in which the two NF- $\kappa$ B sites are mutated) promoter, with or without p65/RelA (p65) expression vector, and with either pSuper (parental vector) or DSIF p160 RNAi as indicated. Cells were harvested 48 hours post-transfection and luciferase activity was measured. RSV promoter-driven *Renilla* luciferase reporter plasmids served to normalize the transfection efficiency. Shown is the relative luciferase activity (luciferase units divided by the activity of co-transfected RSV promoter-driven *Renilla* reporter luciferase, RLU). The data represent the means and standard deviations of seven independent experiments each with independent duplicates (left panel). A representative immunoblot showing DSIF knockdown in transfected cells is shown in the right panel. **C.** 293T cells were co-transfected with p65/RelA expression plasmid together with luciferase reporter genes driven by the RANTES and IP-10 promoters and with either pSuper or DSIF p160 RNAi as indicated. The average RLU of seven independent transfection experiments is shown on the graphs. A representative immunoblot showing DSIF knockdown in cells transfected with the indicated reporter plasmids together with p65/RelA is shown in the right panel.

**Figure 2: DSIF attenuation of NF- $\kappa$ B is dependent on TATA-less core promoter.** **A.** 293T cells were co-transfected with A20 promoter mutants (schematically shown on the right) with sub-optimal concentration of p65/RelA expression vector, and with either pSuper (parental vector) or DSIF p160 RNAi as indicated, and analyzed as described in Fig. 1B. The effect of



DSIF knockdown on different A20 promoter mutants activated by sub-optimal p65/RelA concentration is presented in terms of inhibition fold (the ratio of the relative luciferase activity in the presence of DSIF RNAi to the activity in the presence of the parental vector pSuper). **B.** The effect of DSIF knockdown by two distinct RNAi on A20 promoter wild type and a mutant (A20-TATA) which was converted into a canonical TATA box promoter by substitution of two nucleotides (bold letters) in -30 and -26 positions relative to the transcription start site, in the presence or absence of p65/RelA (sub-optimal concentration). DSIF RNAi 1 is the same one used in Fig.1 and DSIF RNAi 2 is directed against a different region of the p160 subunit (see materials and methods). The results shown in the graphs are presented in terms of inhibition fold as in A and are the average of seven (DSIF RNAi 1) or six (DSIF RNAi 2) independent duplicate transfection experiments. Representative immunoblots showing down regulation of DSIF p160 by the two RNAis are shown on the right. **C.** The responsiveness of the A20 mutants reporter genes to p65/RelA NF- $\kappa$ B protein.

**Figure 3: The core promoter controls differential occupancy of DSIF on A20 gene upon NF- $\kappa$ B induction.** **A.** The wild type and mutant A20 promoters-driven reporter plasmids, as indicated, were each transfected into 293T cells. 24 hours later cells were treated with TNF $\alpha$  for 1 hour, and then subjected to chromatin immunoprecipitation (ChIP) with the indicated antibodies and a non relevant antibody as control, followed by PCR analysis after normalization to the input (see Materials and Methods). The forward primer is specific to each of the promoters and the reverse primer is derived from proximal region of the luciferase gene. Representative ChIP results are shown on the left of the figure (ChIP). 0.1% of the input DNA (Input) was subjected to an increasing number of PCR cycles in order to find the linear range to serve as a reference for the immunoprecipitation. Quantified results, normalized to the input, derived from two independent experiments are shown on the right. **B.** Jurkat T cells were untreated or treated with TNF $\alpha$  for 1 hour, followed by ChIP assay with the indicated antibodies and a non-relevant antibody. The immunoprecipitated DNAs were subjected to

PCR with primers specific to the promoter and beginning of each of the A20,  $\beta$ -actin and BLR1 endogenous genes. Quantified results, normalized to the input, derived from three independent experiments are shown on the right.

**Figure 4: P-TEFb occupancy is the inverse of DSIF upon NF- $\kappa$ B induction.** **A.** ChIP of transfected A20 wild type and mutant reporter genes, as described in Fig. 3A, using antibodies against Pol II, CDK9 subunit of P-TEFb, p160 subunit of DSIF and a non-relevant antibody as control. Representative results of ChIP and input are shown on the left and quantified results, normalized to the input, derived from two independent experiments are shown on the right. **B.** ChIP assay of endogenous A20 gene from control or 1 hour TNF $\alpha$  treated Jurkat T cells as described in Fig. 3B, using antibodies to unphosphorylated form of Pol II (Pol II), CDK9 subunit of P-TEFb (P-TEFb), serine 2 phosphorylated form of Pol II (Ser 2), p160 subunit of DSIF (DSIF) and a non-relevant antibody as control (Con). The immunoprecipitated DNAs were subjected to PCR with primers specific to the promoter (promoter) or to the internal region of these genes (coding). Representative results of ChIP and input are shown on the left and quantified results, normalized to the input, derived from two independent experiments are shown on the right.

**Figure 5: Differential regulation of NF- $\kappa$ B by P-TEFb is core promoter type dependent.** **A.** 293T cells were transfected with the indicated reporters and 24 hours later were treated either with TNF $\alpha$  or TNF $\alpha$  with increasing amounts of DRB for 4 hours. The luciferase activity was normalized to the values of the co-transfected RSV-*Renilla*. The graphs represent the mean and standard deviation of 3 duplicate experiments. **B.** 293T cells were transfected with either wild type A20 or A20-TATA mutant promoters together with p65/RelA expression vector, and with either pSuper (parental vector) or cdk9 RNAi as indicated. Cells were harvested 48 hours post-transfection and luciferase activity was measured. The data represent the means and standard deviations of six independent duplicated experiments (left

panel). A representative immunoblot showing cdk9 depletion in transfected cells is shown in the right panel.

**Figure 6: ChIP analysis of other NF- $\kappa$ B target genes.** The TATA-less I $\kappa$ B $\alpha$  and the TATA-containing RANTES genes were subjected to ChIP assay before or 1 hour after TNF $\alpha$  treatment as described in Fig. 3B. Antibodies to the unphosphorylated form of Pol II (Pol II), serine 2 phosphorylated form of Pol II (Ser 2), p160 subunit of DSIF (DSIF) and a non-relevant control antibody (Con) were used for immunoprecipitation. The precipitated DNAs were subjected to PCR with primers specific to the promoter or the coding region of the genes. Representative results of ChIP and input are shown on the left. Quantified results, normalized to the input, derived from two independent experiments are shown on the right.



**Figure 7: Serine 2 phosphorylation of Pol II on TATA promoter releases DSIF.** Jurkat cells were untreated or treated with TNF $\alpha$  without or with DRB (20 $\mu$ M) for one hour and then subjected to ChIP assay with antibodies to serine 2 phosphorylated form of Pol II (Ser 2), p160 subunit of DSIF (DSIF) and a non-relevant antibody as control (Con) followed by PCR analysis of the TATA-less I $\kappa$ B $\alpha$  and the TATA-containing RANTES promoters. Representative results from two experiments are shown. Quantified results, normalized to the input, derived from two independent experiments are shown on the right.

**Figure 8: Dependency of the different NF- $\kappa$ B responsive promoters on TAF1. A.** Wild type A20 or A20-TATA mutant promoters were each transfected into 293T and analyzed by ChIP as described in Fig. 3A, using antibodies against TAF1, Pol II and a non-relevant antibody as control. Representative results of ChIP and input are shown on the left and quantified results, normalized to the input, derived from two independent experiments are shown on the right. **B.** Fixed chromatin extract was prepared from Jurkat T cells induced by TNF $\alpha$  for 1 hour. ChIP assay was performed as described in Fig. 3B using antibodies to Pol II,



TAF1 and a non-relevant antibody as control. Representative results are shown on the left and quantified results, normalized to the input (the same as in Fig. 4B) derived from two independent experiments on the right. **C.** Hamster ts13 temperature sensitive cells were co-transfected with the indicated reporter plasmids together with the p65/RelA expression plasmid. The cells were incubated at the permissive temperature (32°C) for 6 hours, washed and then separated into two groups. One was grown at 32°C and the second at the non-permissive temperature (39.5°C) for an additional 48 hours, after which the cells were harvested and luciferase activity measured. RSV promoter-driven *Renilla* reporter luciferase plasmids served to normalize transfection efficiency within each group and *c-fos*-luciferase (TAF independent promoter) served to normalize transfection efficiency between the groups. Results are the mean of four independent experiments, each with independent duplicates. **D.** Hamster ts13 cells were grown at 32°C and then shifted to the non-permissive temperature 39.5°C for 6 hours followed by 1 hour treatment with mouse TNF $\alpha$ . Total RNA was extracted and subjected to a quantitative RT-PCR assay to measure the mRNAs of A20, I $\kappa$ B $\alpha$ , and  $\beta$ -actin genes, using Light Cycler. The data represent the means and standard deviations of 2 independent experiments.

**Fig. 1**




**A.**




**B.**




**C.**




**A.**




**B.**



**C.**





**Fig. 3**



**Fig. 3**


**B.**






**Fig. 4**

**A.**




**Fig. 4**

**B.**




**Fig. 5**


**A.**




**B.**



**Fig. 6**




**Fig. 7**




**Fig. 8**

**A.**




**Fig. 8**

**B.**



**Fig. 8**

**C.**



**D.**

